

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/00, 7/021, 7/025, 7/06, 7/09, 7/16, 7/32, 7/42, 31/74		A1	(11) International Publication Number: WO 98/48768
			(43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PC17US98/08931		63 Webster Street, Whitman, MA 02382 (US). LUCZAK, Scott [US/US]; 3 Remsen Avenue, Medfield, MA 02052 (US). MENDUM, Thomas, H., E. [US/US]; 45 Columbus Avenue #1, Somerville, MA 02143 (US). (74) Agents: KREBS, Robert, E. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US). (81) Designated States: A1, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NI, PT, SE), OAPI patent (BF, BJ, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 1 May 1998 (01.05.98)			
(30) Priority Data: 08/846,883 1 May 1997 (01.05.97) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/846,883 (CON) Filed on 1 May 1997 (01.05.97)			
(71) Applicant (for all designated States except US): MEDLOGIC GLOBAL CORPORATION [US/US]; 4815 List Drive, Colorado Springs, CO 80919 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): RON, Eyal, S. [US/US]; 7 Coach Road, Lexington, MA 02173 (US). HAND, Barry, J. [US/US]; 145 Butternut Hollow, Acton, MA 01718 (US). BROMBERG, Lev, S. [US/US]; 17 Sherwood Road, Swampscott, MA 01907 (US). KEARNEY, Marie [US/US]; 342 Faneuil Street #1, Brighton, MA 02135 (US). SCHILLER, Matthew, E. [US/US]; 23C Sagamore Way, Waltham, MA 02154 (US). AHEARN, Peter, M. [US/US];		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: COMPOSITIONS FOR COSMETIC APPLICATIONS

(57) Abstract

A cosmetic composition is described having a cosmetically acceptable carrier, comprising a reverse thermal viscosifying polymer network comprising at least one poloxamer component capable of aggregation in response to a change in temperature randomly bonded to at least one poly(acrylic acid) component; and a cosmetically active agent which imparts a preselected cosmetic effect, said carrier and said agent disposed within an aqueous-based medium.



BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOSITIONS FOR COSMETIC APPLICATIONS

This application is a continuation-in-part application of copending application
5 U.S.S.N. 60/034,805 filed January 2, 1997, and entitled "Responsive Polymer
Networks and Methods of Their Use", which is a continuation-in-part application of
copending application PCT/US96/10376 filed June 14, 1996, designating the United
States, and entitled "Responsive Polymer Networks and Methods of Their Use", which
is a continuation-in-part application of copending application U.S.S.N. 08/580,986 filed
10 January 3, 1996, and entitled "Responsive Polymer Networks and Methods of Their
Use", each of which is incorporated entirely by reference.

Field of the Invention

The present invention relates to a cosmetic composition useful in a variety of
15 topical and personal care products, including treatments of disorders and imperfections
of the skin or other areas of the body. More particularly, the present invention is
directed to a cosmetic composition comprising a poloxamer:poly(acrylic acid)
polymer network that can be designed to reversibly gel over a wide range of
conditions to provide a composition having a controllable range of viscosities, making
20 it useful in a variety of cosmetic and personal care applications.

Background of the Invention

Many examples are known of cosmetic compositions intended for treatment of
the skin or elsewhere on the body, where it is desired to have certain properties of
25 viscosity. Hydrogels, such as cellulose, have been included as thickeners in cosmetic
compositions. A hydrogel is a polymer network which absorbs a large quantity of
water without the polymer dissolving in water. The hydrophilic areas of the polymer
chain absorb water and form a gel region. The extent of gelation depends upon the
volume of the solution which the gel region occupies.

30 Reversibly gelling solutions are known in which the solution viscosity increases

and decreases with an increase and decrease in temperature, respectively. Such reversibly gelling systems are useful wherever it is desirable to handle a material in a fluid state, but performance is preferably in a gelled or more viscous state.

A known material with these properties is a thermal setting gel using block copolymer polyols, available commercially as Pluronic® polyols (BASF, Ludwigshafen, Germany), which is described in U.S. Patent No. 4,188,373. Adjusting the concentration of the polymer gives the desired liquid-gel transition. However, concentrations of the polyol polymer of at least 18-20 % by weight are needed to produce a composition which exhibits such a transition at commercially or physiologically useful temperatures. Also, solutions containing 18-20 % by weight of responsive polymer are typically very viscous even in the "liquid" phase, so that these solutions can not function under conditions where low viscosity, free-flowing is required prior to transition. In addition, these polymer concentrations are so high that the material itself may cause unfavorable interactions during use.

Another known system which is liquid at room temperature, but forms a semi-solid when warmed to about body temperature is formed from tetrafunctional block polymers of polyoxyethylene and polyoxypropylene condensed with ethylenediamine, commercially available as Tetronic® polyols. These compositions are formed from approximately 10% to 50% by weight of the polyol in an aqueous medium. See, U.S. Patent No. 5,252,318.

Joshi *et al.* in U.S. Patent No. 5,252,318 reports reversible gelling compositions which are made up of a physical blend of a pH-sensitive gelling polymer (such as a cross-linked poly(acrylic acid) and a temperature-sensitive gelling polymer (such as methyl cellulose or block copolymers of poly(ethylene glycol) and poly(propylene glycol)). In compositions including methylcellulose, 5- to 8-fold increases in viscosity are observed upon a simultaneous change in temperature and pH for very low methylcellulose levels (1-4% by weight). See, Figs. 1 and 2 of Joshi *et al.* In compositions including Pluronic® and Tetronic® polyols, commercially available forms of poly(ethylene glycol)/poly(propylene glycol) block copolymers, significant increases in viscosity (5- to 8-fold) upon a simultaneous change in temperature and pH

are observed only at much higher polymer levels. See, Figs. 3-6 of Joshi *et al.*

Hoffman *et al.* in WO 95/24430 disclose block and graft copolymers comprising a pH-sensitive polymer component and a temperature-sensitive polymer component. The block and graft copolymers are well-ordered and contain regularly repeating units of the pH-sensitive and temperature-sensitive polymer components. The copolymers are described as having a lower critical solution temperature (LCST), at which both solution-to-gel transition and precipitation phase transition occur. Thus, the transition to a gel is accompanied by the clouding and opacification of the solution. Light transmission is reduced, which may be undesirable in many applications, where the aesthetic characteristics of the composition are of some concern.

Thus, the known systems which exhibit reversible gelation are limited in that they require large solids content and/or in that the increase in viscosity less than 10-fold. In addition, some known systems exhibit an increase in viscosity which is accompanied with the undesirable opacification of the composite.

Summary of the Invention

It is an object of the present invention to provide a cosmetic composition which is includes a component capable of reversible gelation or viscosification.

It is a further object of the invention to provide a cosmetic composition which includes an ingredient capable of gelation or viscosification at very low solids content.

It is another object of the present invention to provide a cosmetic composition which possesses improved flow and gelation characteristics as compared to properties possessed by conventional reversible gelation compositions.

It is a further object of the invention to provide a polymer network composition for use in cosmetic compositions useful as a surfactant or emulsifier in the solubilization of additives and, in particular, hydrophobic additives.

It is a further object of the invention to provide a cosmetic composition which possesses the appropriate thickness, emolliency and cosmetic effect with a minimum of solids content.

It is a further object of the invention to provide a polymer network for use in

cosmetic compositions useful as a suspending agent for otherwise insoluble additives.

It is yet a further object of the present invention to provide a composition capable of solubilizing emulsions at elevated temperatures.

It is yet a further object of the invention to provide new and useful cosmetic compositions incorporating the reversibly gelling polymer network composition of the present invention, which take advantage of its unique advantageous properties.

It is yet another object of the present invention to provide reversibly gelling polymer network compositions which are composed of biocompatible polymers.

These and other objects of the invention are achieved with a cosmetic composition which incorporates a poloxamer:poly(acrylic acid) polymer network as a cosmetically acceptable carrier. The polymer network comprises a poloxamer component randomly bonded to a poly(acrylic acid), or P.A.A. component in an aqueous-based medium, the polymer network being capable of aggregating in response to an increase in temperature. The reverse thermal viscifying poloxamer:poly(acrylic acid) polymer network includes random covalent bonding between the poly(acrylic acid) component and the poloxamer component of the network. The polymer network may also include some unbound or "free" poloxamer or other additives which contribute to or modify the characteristic properties of the polymer composition.

In addition, the cosmetic composition includes a cosmetic agent selected to provide a preselected cosmetic effect. By "cosmetic agent", as that term is used herein, it is meant that the additive imparts a cosmetic effect. A cosmetic effect is distinguishable from a pharmaceutical effect in that a cosmetic effect relates to the promoting bodily attractiveness or masking the physical manifestations of a disorder or disease. In contrast, a pharmaceutical seeks to treat the source or symptom of a disease or physical disorder. It is noted however, that the same additives may have either a cosmetic or pharmaceutical effect, depending upon the amounts used and the manner of administration.

By "cosmetic" as that term is used herein, it is meant the cosmetic and

personal-care applications intended to promote bodily attractiveness or to cover or mask the physical manifestations of a disorder or disease. Cosmetics include those products subject to regulation under the FDA cosmetic guidelines, as well as sunscreen products, acne products, skin protectant products, anti-dandruff products, and deodorant and antiperspirant products.

By "gelation" or viscosification, as that term is used herein, it is meant a drastic increase in the viscosity of the polymer network solution. Gelation is dependent on the initial viscosity of the solution, but typically a viscosity increase in the range of preferably 2- to 100-fold, and preferably 5- to 50-fold, and more preferably 10- to 20-fold is observed in the polymer network which is used in the preparation of the cosmetic compositions of the invention. Such effects are observed in a simple polymer network solution and the effect may be modified by the presence of other components in the cosmetic composition.

By "reversibly gelling" as that term is used herein, it is meant that the process of gelation takes place upon an *increase* in temperature rather than a decrease in temperature. This is counter-intuitive, since it is generally known that solution viscosity *decreases* with an increase in temperature.

As used herein, "poloxamer" is a triblock copolymer derived from poly(ethylene glycol)-poly(propylene glycol)-poly(ethylene glycol) blocks. The poloxamer is capable of responding to a change in temperature by altering its degree of association and/or agglomeration. The aggregation may be in the form of micelle formation, precipitation, labile crosslinking or other factors. The poloxamer has the general formula of a triad ABA block copolymer, $(P_1)_a(P_2)_b(P_1)_a$, where P_1 = poly(ethylene glycol) and P_2 = poly(propylene glycol) blocks, where a is in the range of 10-50 and where b is in the range of 50-70.

The poly(acrylic acid) component includes poly(acrylic acid) and its salts. The poly(acrylic acid) supports and interacts with the poloxamer component so that a multi-material, responsive polymer network is formed. The interaction of the poloxamer and poly(acrylic acid) exhibits a synergistic effect, which magnifies the effect of the poloxamer component in viscosifying and/or gelling the solution.

The novel interaction between the constituent polymers components of the polymer network permits formation of gels at very low solids content. Gelation and/or viscosification is observed in aqueous solutions having about 0.01 to 20 wt% of the poloxamer component and about 0.01 to 20 wt% of the poly(acrylic acid) component.

5 A typical reversibly gelling polymer network may be comprised of less than about 4 wt% of total polymer solids (e.g., poloxamer and poly(acrylic acid)) and even less than 1 wt% total polymer solids while still exhibiting reverse thermal viscosification. Of course, the total solids content including additives of a reversibly gelling polymer network composition may be much higher. The viscosity of the gel increases at least
10 ten-fold with an increase in temperature of about 5°C at pH 7 and 1 wt% polymer. Viscosity increases may be even greater over a larger temperature range at pH 7 and 1% polymer network content.

The relative proportion of poloxamer and poly(acrylic acid) may vary dependent upon the desired properties of the polymer composition. In one
15 embodiment, the poloxamer is present in a range of about 1 to 20 wt% and the poly(acrylic acid) is present in a range about of 99 to 80 wt%. In another embodiment, the poloxamer component is present in a range of about 21 to 40 wt% and the poly(acrylic acid) component is present in a range of about 79 to 60 wt%. In another embodiment, the poloxamer component is present in a range of about 41 to 50
20 wt% and the poly(acrylic acid) component is present in a range of about 59 to 50 wt%. In another embodiment, the poloxamer component is present in a range of about 51 to 60 wt% and the poly(acrylic acid) component is present in a range of about 49 to 40 wt%. In yet another embodiment, the poloxamer component is present in a range of about 61 to 90 wt% and the poly(acrylic acid) component is present in a
25 range of about 39 to 20 wt%. In another embodiment, the poloxamer component is present in a range of about 81 to 99 wt% and the poly(acrylic acid) component is present in a range of about 19 to 1 wt%.

The poloxamer:poly(acrylic acid) polymer network described above is included
30 in a cosmetic composition to improve the flow characteristics, thickness and other properties of the composition. The composition includes additional cosmetic agents.

such as are needed for the cosmetic purpose of the composition. Additives also may be included to modify the polymer network performance, such as to increase or decrease the temperature of the liquid-to-gel transition and/or to increase or decrease the viscosity of the responsive polymer composition.

5 In one aspect of the invention, the poloxamer:poly(acrylic acid) polymer network is incorporated into a cosmetic compositions to impart thickening properties to the cosmetic composition at the use and/or application temperature. Such thickening properties include enhanced overall viscosity, as well as a desirable viscosity response with temperature. The polymer network may be useful as a thickener in pH ranges
10 where other thickeners are not effective.

In another aspect of the invention, the poloxamer:poly(acrylic acid) polymer network is incorporated into a cosmetic composition to stabilize and solubilize hydrophobic agents in the cosmetic composition. The polymer network may be included to increase emulsion stability. Many emulsions, i.e., suspension of small
15 droplets or particles of a first material in a second material, lose viscosity upon heating. As will be demonstrated herein, the poloxamer:poly(acrylic acid) polymer network retains its emulsifying properties even with temperature increase.

In addition, it may be included in the composition to impart emolliency to the composition. The composition may also act as a film-forming agent after it has been
20 applied to the skin. This film-forming agent may be used as a barrier to prevent water loss from the skin which contributes to the moisturization of the skin.

In another aspect of the invention, the poloxamer:poly(acrylic acid) polymer network may be included as an additive in cosmetic applications to prevent viscosity loss at elevated temperatures.

Brief Description of the Drawing

The invention is described with reference to the Drawing, which is presented for the purpose of illustration and is in no way intended to be limiting, and in which:

Figure 1 is a graph of viscosity vs. temperature for a 1 wt%, 2 wt% and 3 wt%
30 responsive polymer network aqueous composition of a poloxamer/poly(acrylic acid)

(1:1) at pH 7.0 measured at a shear rate of 0.44 sec^{-1} ;

Figure 2 is a graph of viscosity vs. temperature for a 1 wt% poloxamer: poly(acrylic acid) polymer network composition demonstrating reversibility of the viscosity response;

5 Figure 3 shows the viscosity response of a 2 wt% poloxamer:poly(acrylic acid) polymer composition at various shear rates;

Figure 4 shows a viscosity response curve for a 2 wt% poloxamer: poly(acrylic acid) polymer network composition prepared with nominal mixing and stirring and prepared using high shear homogenization (8000 rpm, 30 min);

10 Figure 5 is a graph of viscosity vs. temperature for a 1 wt% poloxamer: poly(acrylic acid) polymer network composition at various pHs;

Figure 6 is a graph of viscosity vs. temperature for a 1 wt% poloxamer: poly(acrylic acid) polymer network composition with and without addition of 0.25 wt% KCl;

15 Figure 7 is a graph of viscosity vs. temperature for a 1 wt% poloxamer: poly(acrylic acid) polymer network composition with and without addition of 0.5 wt% acetamide MEA;

Figure 8 is a graph of viscosity vs. temperature for a 1 wt% poloxamer: poly(acrylic acid) polymer network composition without and with 5 wt%, 10 wt% and 20 wt% added ethanol, respectively;

Figure 9 is an illustration of a reversibly gelling polymer network used as an emulsifier and stabilizer for a hydrophobic agent;

Figure 10 is a schematic illustration of the poloxamer:poly(acrylic acid) polymer network below and above the transition temperature illustrating the aggregation of the hydrophobic poloxamer regions;

25 Figure 11 is a graph of viscosity vs. pH for a 1 wt% responsive polymer network aqueous composition of a poloxamer/poly(acrylic acid) (1:1) measured at a shear rate of 0.44 sec^{-1} ;

Figure 12 is a plot of viscosity vs. temperature for (a) a 1 wt% responsive polymer network aqueous composition of Pluronic® F127 poloxamer/poly(acrylic acid)

30

(1:1) and (b) a 1 wt% physical blend of Pluronic® F127 poloxamer/poly(acrylic acid) (1:1) at pH 7.0 measured at a shear rate 0.22 sec^{-1} ;

Figure 13 is a plot of viscosity vs. temperature for a 1 wt% responsive polymer network aqueous composition of Pluronic® F88 poloxamer/poly(acrylic acid) (1:1) at pH 7.0 measured at a shear rate 2.64 sec^{-1} ;

Figure 14 is a graph of the viscosity vs. temperature effect for a responsive polymer network composition of 2 wt% Pluronic® P104 poloxamer/poly(acrylic acid) (1:1) in deionized water at pH 7.0 measured at shear rate of 22 sec^{-1} ;

Figure 15 is plot of viscosity vs. temperature for a responsive polymer network composition of 2 wt% Pluronic® F123 poloxamer/poly(acrylic acid) (1:1) at pH 7.0 measured at a shear rate of 22 sec^{-1} ;

Figure 16 is a plot of viscosity vs. temperature for 1 wt% made of series of poloxamers and poly(acrylic acid) (1:1) in deionized water at a shear rate of 132 sec^{-1} ;

Figure 17 is a plot showing release of hemoglobin from a poloxamer/poly(acrylic acid) polymer network of the invention;

Figure 18 is a plot showing the release of lysozyme from the poloxamer/poly(acrylic acid) polymer complex of the invention;

Figure 19 is a plot showing release of insulin from a poloxamer/poly(acrylic acid) polymer network composition of the invention;

Figure 20 is a plot of viscosity vs. temperature for a poloxamer/poly(acrylic acid) polymer network composition (a) before and (b) after sterilization by autoclave;

Figure 21 is a plot of viscosity vs. temperature for an oil-free moisturizing formulation prepared from (a) a responsive polymer network composition of the invention and (b) a conventional oil-in-water formulation;

Figure 22 is a plot of equilibrium solubility of estradiol (A, B) and progesterone (C, D) in aqueous solutions (pH 7) of Pluronic® F127 (A, C) and responsive polymer network (B, D) vs. temperature;

Figure 23 is a plot of the ratio of equilibrium solubilities of estradiol in responsive polymer network and water vs. polymer concentration in the responsive polymer network solutions;

Figure 24 is a plot of the effect of loading fluorescein on the onset of gelation of responsive polymer network vs. total polymer concentration in responsive polymer network solution (pH 7.0);

Figure 25 is a plot of the percentage of a) estradiol and b) progesterone release
5 from responsive polymer network vs. time;

Figure 26 is a plot of the rate of progesterone release and macroscopic viscosity vs. polymer concentration;

Figure 27 is a plot of the percentage of progesterone release vs. polymer concentration in responsive polymer network and,

10 Figure 28 is a plot of the relative diffusivity of poly(styrene) latex particles in water and responsive polymer network.

Detailed Description of the Invention

The present invention is directed to a cosmetic composition comprising a
15 cosmetically acceptable carrier comprising a novel poloxamer:poly(acrylic acid) polymer network. The polymer network functions as a temperature sensitive thickening agent, and in addition possesses surfactant and emulsifying capabilities which may be beneficial to the cosmetic composition. The polymer network composition according to the invention includes a poloxamer component randomly
20 bonded to a poly(acrylic acid) component. The two polymer components may interact with one another on a molecular level. The polymer network contains about 0.01-20 wt% each of poloxamer and poly(acrylic acid). Exemplary polymer network-compositions range from about 1:10 to about 10:1 poloxamer:poly(acrylic acid). Polymer network gel compositions which exhibit a reversible gelation at body
25 temperature (25-40°C) and/or at physiological pH (ca. pH 3.0-9.0) and even in basic environments up to pH 13 (hair care) are particularly preferred for cosmetic applications.

In one embodiment of the invention, a 1:1 poloxamer:poly(acrylic acid) polymer network at appropriate pH exhibits flow properties of a liquid at about room
30 temperature, yet rapidly thickens into a gel consistency of at least about five times

greater, preferably at least about 10 times greater, and even more preferably at least about 30 times and up to 100 times greater, viscosity upon increase in temperature of about 10 °C and preferably about 5 °C. The reversibly gelling polymer network of the present invention exhibit gelation even at very low polymer concentrations. For

5 example, polymer network compositions at pH 7 comprising about 0.5 wt% poloxamer component and about 0.5 wt% PAA exhibits a significant increase in viscosity from a free-flowing liquid (50 cps) to a gel (6000 cps). The observed gelation takes place at low solids contents, such as less than 20 wt% or preferably less than about 10 wt%, or more preferably less than about 2.5 wt% or most preferably less than about 0.1 wt%.

10 Thus, only a small amount by weight of the polymer network need be incorporated into a cosmetic composition in order to provide the desired thickening or viscosifying effect.

The reverse viscosification effect at low polymer concentrations provides clear, colorless gels which are particularly well-suited to cosmetic applications. For example, very little residue is formed upon dehydration which may be important in some

15 applications, such as in topically applied cosmetics. An additional advantage of the polymer network of the invention is that it remains clear and translucent above and below the critical temperature or pH. These characteristics of the reversibly gelling polymer network make it well suited for use in cosmetic compositions.

20 The polymer network of the present invention technology may be added to cosmetic formulations to increase the thickness and viscosity of the composition. The poloxamer:poly(acrylic acid) polymer network possesses hydrophobic regions capable of aggregation. Unlike conventional thickeners, the aggregation of the polymer network of the present invention is temperature sensitive. Thus, the inventive polymer network of the present invention may have a transition temperature (i.e. temperature of aggregation) above room temperature so that the cosmetic composition is of low

25 viscosity at or below room temperature and is of high viscosity at or around body temperature (body temperature includes both surface and internal body temperature). Thus, a composition may be prepared at low temperatures while the polymer network is in a low viscosity state. Mixing of ingredients under low viscosity is expected to be

30

easier, thus simplifying the manufacturing process. Yet, the resultant mixture would be of increased viscosity at use temperatures. As a further advantage, a cosmetic composition comprising poloxamer:poly(acrylic acid) polymer network may be spread thinly to allow for even application, due to its low viscosity at room temperature, but will thicken and "fill" the skin contours upon warming up to body surface temperature.

In another aspect of the invention, the composition may be applied through a nozzle that provides high shear to reduce viscosity, yet the composition regains its viscosity after application to the skin. This contrasts with conventional formulations which permanently lose viscosity after being subjected to high shear.

In another aspect of the invention, the composition may be formulated and applied as a liquid, spray, semi-solid gel, cream, ointment, lotion, stick, roll-on formulation, mousse, pad-applied formulation, and film-forming formulation.

The poloxamer:poly(acrylic acid) polymer network may also be included in a cosmetic composition for use as a stabilizing, solubilizing or emulsifying agent for a hydrophobic component of the cosmetic formulation. The strong hydrophilic regions of the poloxamer resulting from aggregation and micelle formation create hydrophobic domains which may be used to solubilize and control release of hydrophobic agents. Similar micelle-based systems have been shown to protect trapped peptides against enzymatic degradation from surface enzymes.

The reversibly gelling polymer network of the present invention is a unique polymer composition designed to abruptly change its physical characteristics or the characteristics and properties of materials mixed therewith with a change in temperature. Without intending to be bound by any particular mechanism or chemical structure, it is believed that the structure of the polymer network involves a random bonding of the poloxamer onto the backbone of the poly(acrylic acid). A portion of the poloxamer which is present during the polymerization reaction which forms the poly(acrylic acid) is bonded to the backbone of the forming poly(acrylic acid) through hydrogen abstraction and subsequent reaction. See detailed discussion of the mechanism, below. The combination of the poly(acrylic acid) and randomly bonded poloxamer gives the composition its unique properties. Any free poloxamer remaining

after polymerization of PAA remains associated with the random co-polymer, resulting in a miscible composition. Free poloxamer may also be present in the polymer network composition; however, its presence is not required in order to observe reverse thermal viscosification.

5 The poly(acrylic acid) may be linear, branched and/or crosslinked. Poly(acrylic acid) is capable of ionization with a change in pH of the solution. By ionization, as that term is used with respect to poly(acrylic acid), it is meant the formation of the conjugate base of the acrylic acid, namely acrylate. As used herein, poly(acrylic acid) includes both ionized and non-ionized versions of the polymer. Changes in ionic
10 strength may be accomplished by a change in pH or by a change in salt concentration. The viscosifying effect of the polymer network is partly a function of the ionization of the poly(acrylic acid); however, reverse thermal gelling may occur without ionization. Changes to the ionic state of the polymer causes the polymer to experience attractive (collapsing) or repulsive (expanding) forces. Where there is no need or desire for the
15 composition to be applied in a high viscosity state, it may be possible to prepare the composition as non-ionized poly(acrylic acid). The body's natural buffering ability will adjust the pH of the applied composition to ionize the poly(acrylic acid) and thereby develop its characteristic viscosity.

 The poloxamer possesses regions of hydrophobic character, e.g., poly(propylene glycol) blocks, and hydrophilic character, e.g., poly(ethylene glycol) blocks. The
20 poloxamer may be linear or branched. Suitable poloxamers include triad block copolymers of poly(ethylene glycol) and poly(propylene glycol) having the general formula $(P_1)_a(P_2)_b(P_1)_a$, where P_1 = poly(ethylene glycol) and P_2 = poly(propylene glycol) blocks, where a is in the range of 10-50 and where b is in the range of 50-70.
25 where poly(propylene glycol) represents the hydrophobic portion of the polymer and poly(ethylene glycol) represents the hydrophilic portion of the polymer. Pluronic® polymers (BASF) are commercially available for a in the range of 16 to 48 and b ranging from 54-62. One or more poloxamers may be used in the reversibly gelling polymer network composition of the present invention.

30 The reversibly gelling responsive polymer networks compositions of the present

invention are highly stable and do not exhibit any phase separation upon standing or upon repeated cycling between a liquid and a gel state. Samples have stood at room temperature for more than three months without any noticeable decomposition, clouding, phase separation or degradation of gelation properties. This is in direct contrast to polymer blends and aqueous mixed polymer solutions, where phase stability and phase separation is a problem, particularly where the constituent polymers are immiscible in one another.

An example of the dramatic increase in viscosity and of the gelation of the reversibly gelling polymer network compositions of the invention is shown in Figure 1. Figure 1 is a graph of viscosity vs. temperature for 1 wt%, 2 wt% and 3 wt% polymer network compositions comprising 1:1 poloxamer:poly(acrylic acid), hydrated and neutralized. The viscosity measurements were taken on a Brookfield viscometer at a shear rate of 0.44 sec^{-1} at pH 7.0. All solutions had an initial viscosity of about 1080 cP and exhibited a dramatic increase in viscosity to gel point at about 35°C . This is not typical of all polymer network compositions since polymerization condition will affect initial viscosity. Final viscosities were approximately 33,000 cP, 100,000 cP and 155,000 cP for the 1 wt%, 2 wt% and 3 wt% compositions, respectively. This represents viscosity increases of about 30-, 90- and 140-fold, respectively. This effect is entirely reversible. Upon cooling, the composition regains its initial viscosity. This is demonstrated in Figure 2, where a 1 wt% poloxamer:poly(acrylic acid) composition is warmed through the transition temperature up to 35°C (simple curve), cooled to room temperature (24°C , ticked curve) and then warmed again to up above the transition temperature (open box curve). The viscosity response was virtually identical in all three instances.

As would be expected with a non-Newtonian system, the solution viscosity differs with different shear rates. Figure 3 shows the viscosity response of a 2 wt% poloxamer:poly(acrylic acid) polymer composition at various shear rates. The viscosity response is consistent between 24°C and 34°C ; however, the final viscosity is reduced with increasing shear rate.

However, unlike many prior art hydrogels, e.g., carbomers, the

poloxamer:poly(acrylic acid) polymer network composition does not permanently lose viscosity after being subjected to high shear conditions. The poloxamer:poly(acrylic acid) polymer network composition remains unaffected by such shear conditions as homogenization. Figure 4 compares the viscosity response curve of a 2 wt%
5 poloxamer:poly(acrylic acid) polymer composition prepared with nominal mixing (simple line) and stirring with that of a polymer composition of similar composition prepared using high shear homogenization designated by a ticked line (8000 rpm, 30 min). No significant decrease in viscosity is observed.

A number of factors influence the viscosity and transition temperature of the
10 composition. The more important factors include polymer concentration, pH and presence and nature of additives.

The effect of pH on the viscosity of reversibly gelling polymer networks is shown in Figure 5. Increasing pH from the starting pH has a lesser effect on the viscosity than decreasing the pH. This may relate to the extent of ionization of the
15 poly(acrylic acid) component of the polymer network as discussed above. This may be clearly seen in Figure 5 when comparing the viscosity response of a 1 wt% poloxamer:poly(acrylic acid) polymer composition at pH 5 and pH 11. Satisfactory viscosities can be obtained at high pHs indicating the potential value of the reversibly gelling polymer network in products such as depilatories, hair straighteners and hair
20 relaxers.

The responsive polymer network may also include additives for influencing the performance of the polymer composition, such as the transition temperature and the viscosity of the polymer composition above the transition temperature. The following
25 list is not intended to be exhaustive but rather illustrative of the broad variety of additives which can be used.

These materials include solvents (e.g., 2-propanol, ethanol, acetone, 1,2-pyrrolidinone, N-methylpyrrolidinone), salts (e.g., calcium chloride, sodium chloride, potassium chloride, sodium or potassium phosphates, borate buffers, sodium citrate),
preservatives (benzalkonium chloride, phenoxyethanol, sodium
30 hydroxymethylglycinate, ethylparaben, benzoyl alcohol, methylparaben, propylparaben,

butylparaben, Germaben II), humectant/moisturizers (acetamide MEA, lactimide MEA, hydrolyzed collagen, mannitol, panthenol, glycerin), lubricants (hyaluronic acid, mineral oil, PEG-60-lanolin, PPG-12-PEG-50-lanolin, PPG-2 myristyl ether propionate) and surfactants.

5 Surfactants may be divided into three classes: cationic, anionic, and nonionics. An example of a cationic surfactant used is ricinoleamidopropyl ethyldimonium ethosulfate (Lipoquat R). Anionic surfactants include sodium dodecyl sulfate and ether sulfates such as Rhodapex CO-436. Nonionic surfactants include Surfynol CT-111, TG, polyoxyethylene sorbitan fatty acid esters such as Tween 65 and 80, sorbitan fatty
10 acid esters such as Span 65, alkylphenol ethoxylates such as Igepal CO-210 and 430, dimethicone copolyols such as Dow Corning 190, 193, and Silwet L7001.

 The addition of polymers has been studied including xanthan gum, celluloses such as hydroxyethylcellulose (HEC), carbomethoxycellulose (CMC), lauryldimonium hydroxypropyl oxyethyl cellulose (Crodacel QL), hydroxypropylcellulose (HPC), and
15 hydroxypropylmethylcellulose (HPMC), poly(acrylic acid), cyclodextrins, methyl acrylamido propyl triammonium chloride (MAPTAC), polyethylene oxide, polyvinylpyrrolidone, polyvinyl alcohol, and propylene oxide/ethylene oxide random copolymers. Poloxamers may also be used as additives. Examples include both the Pluronic® polyols having an $(P_1)_a(P_2)_b(P_1)_a$ structure such as Pluronic® F38, L44, P65, F68, F88, L92, P103, P104, P105, F108, L122 and F127, as well as the reverse
20 Pluronic® R series $(P_2)_a(P_1)_b(P_2)_a$ structure such as Pluronic® 17R2 and 25R8. Other miscellaneous materials include propylene glycol, urea, triethanolamine, alkylphenol ethoxylates (Iconol series), and linear alcohol alkoxyates (Plurafac series).

 Additives affect the viscosity of the compositions differently depending upon
25 the nature of the additive and its concentration. Some additives will affect the initial or final viscosity, whereas others will affect the temperature range of the viscosity response, or both.

 Potassium chloride and acetamide MEA are two examples of additives which decrease the final viscosity of the composition (see, Example 30). KCl (0.25%) added
30 to a 1 wt% reversibly gelling polymer composition reduces the viscosity by about 3000

cps. See, Figure 6. The humectant, acetamide MEA, lowers the viscosity of a 1 wt% solution by approximately 1,500 cps (see, Figure 7).

Glycerin, ethanol and dimethicone copolymer have been shown to affect the temperature range over which the viscosity response occurs. Glycerin shifts the transition temperature to a slightly lower range from an initial 24-34 °C to about 24-30 °C, but does not affect the final viscosity (see, Example 44). The effect of ethanol on the viscosity is different at different concentration levels. At 5 wt% and 10 wt% added ethanol, the transition temperature is shifted to lower ranges, e.g., 24-29 °C and 20-29 °C, respectively. At 20 wt% added ethanol, the composition not only exhibits a lowering of the transition temperature, but also a marked increase in initial and final viscosity. See, Figure 8. Dimethicone copolymer (1 wt%) also changed the transition temperature, but in this instance the transition temperature range was raised to 28-41 °C. Thus, proper selection of additives permits the formulator to adjust the transition temperature to various ranges.

Those skilled in the art will appreciate that the polymer network compositions of the present invention may be utilized for a wide variety of cosmetic and personal care applications. To prepare a cosmetic composition, an effective amount of cosmetically active agent(s) which imparts the desirable cosmetic effect is incorporated into the reversibly gelling polymer network composition of the present invention.

Preferably the selected agent is water soluble, which will readily lend itself to a homogeneous dispersion through out the reversibly gelling polymer network composition; however, the polymer network has been demonstrated to significantly solubilize or suspend hydrophilic agents in order to improve formulation homogeneity (see, Example 36). It is also preferred that the agent(s) is nonreactive with the polymer network composition. For materials which are not water soluble, it is also within the scope of the invention to disperse or suspend powders or oil (lipophilic materials) throughout the polymer network composition. It will also be appreciated that some applications may require a sterile environment. It is contemplated as within the scope of the invention that the reversibly gelling polymer network compositions of the present invention may be prepared under sterile conditions. An additional feature

of the reversibly gelling polymer composition is that is prepared from constituent polymers that have known accepted toxicological profiles.

The poloxamer:poly(acrylic acid) polymer network has been evaluated under Good Laboratory Practice (GLP) standard protocols known in the art for toxicity in animal models and found to exhibit no toxic effects. The results of the toxicity study are summarized in the following Table 1. The non-toxicity of the polymer network makes it an ideal candidate for use in cosmetic compositions.

Table 1. Toxicity data for 6% poloxamer:poly(acrylic acid) solution at pH 7.

Reaction testes	mode of testing	results
Skin sensitization	guinea pig - topical	not a sensitizer
eye irritation	rabbit eye instillation	negative
primary dermal irritation	rabbit - topical	very slight edema (1 on a scale of 1-8)
acute dermal toxicity	rat - single dose (2g/kg)	no toxicity
acute oral toxicity	rat - single dose (5g/kg)	no toxicity
AMES test		negative

Exemplary cosmetic and personal care applications, for which the reversibly gelling polymer network composition may be used include, but are not limited to, baby products, such as baby shampoos, lotions, powders and creams; bath preparations, such as bath oils, tablet and salts, bubble baths, bath fragrances and bath capsules; eye makeup preparations, such as eyebrow pencil, eyeliner, eye shadow, eye lotion, eye makeup remover and mascara; fragrance preparations, such as colognes and toilet waters, powders and sachets; noncoloring hair preparations, such as hair conditioner, hair spray, hair straighteners, permanent waves, rinses shampoos, tonics, dressings and other grooming aids; color cosmetics; hair coloring preparations such as hair dye, hair tints, hair shampoos, hair color sprays, hair lighteners and hair bleaches; makeup preparations such as face powders, foundations, leg and body paints, lipstick, makeup bases, rouges and makeup fixatives; manicuring preparations such as basecoats and

undercoats, cuticle softeners, nail creams and lotions, nail extenders, nail polish and enamel, and nail polish and enamel remover; oral hygiene products such as dentrifices and mouthwashes; personal cleanliness, such as bath soaps and detergents, deodorants, douches and feminine hygiene product; shaving preparations such as aftershave lotion, beard softeners, men's talcum, shaving cream, shaving soap and preshave lotions; skin care preparations such as cleansing preparations, skin antiseptics, depilatories, face and neck cleansers, body and hand cleansers, foot powders and sprays, moisturizers, night preparations, paste masks, and skin fresheners; and suntan preparations such as suntan creams, gels and lotions, indoor tanning preparations.

Preparation of the above-named cosmetic compositions and others may be accomplished with reference to any of the cosmetic formulation guidebooks and industry journals which are available in the cosmetic industry. These references supply standard formulations which may be modified by the addition or substitution of the reversible viscosifying polymer network of the present invention into the formulation. Suitable guidebooks include Cosmetics and Toiletries Magazine, Vol. 111 (March, 1996); Formularv: Ideas for Personal Care; Croda, Inc. Parsippany, NJ (1993); and Cosmeticon: Cosmetic Formularv, BASF, which are hereby incorporated in their entirety by reference.

The cosmetic composition may be in any form. Suitable forms include but are not limited to lotions, creams, sticks, roll-ons formulations, mousses, aerosol sprays, pad-applied formulations, and film-forming formulations.

As those skilled in the art will appreciate, the foregoing list is exemplary only. Because the reversibly gelling polymer network composition of the present invention is suited for application under a variety of physiological conditions, a wide variety of cosmetically active agents may be incorporated into and administered from the polymer network composition. In addition to the poloxamer:poly(acrylic acid) polymer network, additional cosmetically acceptable carriers may be included in the composition, such as by way of example only, emollients, surfactants, humectants, powders and other solvents. By way of example only, the cosmetic composition also may include additional components, which serve to provide additional aspects of the

cosmetic affect or to improve the stability and/or administration of the cosmetic. Such additional components include, but are not limited to, preservatives, abrasives, acidulents, antiacne agents, anti-aging agents, antibacterials, anticaking, anticaries agents, anticellulites, antidandruff, antifungal, anti-inflammatories, anti-irritants, antimicrobials, antioxidants, astringents, antiperspirants, antiseptics, antistatic agents, astringents, binders, buffers, additional carriers, chelators, cell stimulants, cleansing agents, conditioners, deodorants, depilatories, detergents, dispersants, emollients, emulsifiers, enzymes, essential oils, exfoliants, fibers, film forming agents, fixatives, foaming agents, foam stabilizers, foam boosters, fungicides, gellants, glosser, hair conditioner, hair set resins, hair sheen agents, hair waving agents, humectants, lubricants, moisture barrier agents, moisturizers, ointment bases, opacifier, plasticizer, polish, polymers, powders, propellant, protein, refatting agents, sequestrant, silicones, skin calming agents, skin cleansers, skin conditioners, skin healing, skin lightening agents, skin protectants, skin smoothing agents, skin softening agents, skin soothing agents, stabilizers, sunscreen agents, surfactants, suspending agents, tanning accelerators, thickeners, vitamins, waxes, wetting agents, liquefiers, colors, flavors and/or fragrances. Suitable materials which serve the additive functions listed here are well known in the cosmetic industry. A listing of the additive function and materials suitable for incorporation into the cosmetic composition may be found in Appendix A, which is appended hereto at the end of the specification. Further information may be obtained by reference to The Cosmetic Bench Handbook, Cosmetics & Toiletries; C.C. Urbano, editor, Allured Publ. Corp., 1996, which is hereby incorporated in its entirety by reference.

A brief description of some preferred additives and cosmetically active agents follows. The compositions of the invention include a safe and effective amount of a cosmetically active agent. "Safe and effective", as it is used herein, means an amount high enough to significantly positively modify the condition to be treated or the cosmetic effect to be obtained, but low enough to avoid serious side effects.

Preservatives can be desirably incorporated into the cosmetic compositions of the invention to protect against the growth of potentially harmful microorganisms.

Suitable preservatives include, but are not limited to, alkyl esters of para-hydroxybenzoic acid, hydantoin derivatives, parabens, propionate salts, triclosan tricarbanilide, tea tree oil, alcohols, farnesol, farnesol acetate, hexachlorophene and quaternary ammonium salts, such as benzolconjure, and a variety of zinc and
5 aluminum salts. Cosmetic chemists are familiar with appropriate preservatives and may select that which provides the required product stability. Preservatives are preferably employed in amounts ranging from about 0.0001% to 2% by weight of the composition.

Emollients can be desirably incorporated into the cosmetic compositions of the
10 invention to provide lubricity to the formulation. Suitable emollients may be in the form of volatile and nonvolatile silicone oil, highly branched hydrocarbons and synthetic esters. Amounts of emollients may be in the range of about 0.1-30 wt%, and preferably about 1-20 wt%. By way of example only, suitable silicones include cyclic or linear polydimethylsiloxanes, polyalkylsiloxanes, polyalkylarylsiloxanes and
15 polyether siloxanes. By way of example only, suitable ester emollients include alkenyl esters of fatty acids, polyhydric alcohols, such as ethylene glycol mono and di-fatty acid esters, polyethylene glycol and the like, ether-esters, such as fatty acid esters of ethoxylated fatty alcohols, wax esters, such as beeswax, spermaceti, myristyl myristate and stearyl stearate, and sterol esters, such as cholesterol fatty acids.

A variety of oily emollients may be employed in the compositions of this
20 invention. These emollients may be selected from one or more of the following classes: 1. Triglyceride esters such as vegetable and animal fats and oils. Examples include castor oil, cocoa butter, safflower oil, cottonseed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, squalene, Kikui oil and
25 soybean oil; 2. Acetoglyceride esters, such as acetylated monoglycerides; 3. Ethoxylated glycerides, such as ethoxylated glyceryl monostearate; 4. Alkyl esters of fatty acids having 10 to 20 carbon atoms, such as, methyl, isopropyl, and butyl esters of fatty acids, and including hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate decyl stearate,
30 isopropyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate,

diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate; 5. alkenyl esters of fatty acids having 10 to 20 carbon atoms, such as oleyl myristate, oleyl stearate, and oleyl oleate and the like; 6. fatty acids having 10 to 20 carbon atoms, such as pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic, and erucic acids and the like; 7. fatty alcohols having 10 to 20 carbon atoms, such as, lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecanyl alcohols are examples of satisfactory fatty alcohols and the like; 8. fatty alcohol ethers, such as ethoxylated fatty alcohols of 10 to 20 carbon atoms including the lauryl, cetyl, stearyl, isostearyl, oleyl, and cholesterol alcohols, having attached thereto from 1 to 50 ethylene oxide groups or 1 to 50 propylene oxide groups; 9. ether-esters such as fatty acid esters of ethoxylated fatty alcohols; 10. Lanolin and derivatives, such as lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated lanolin, ethoxylated lanolin alcohols, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin alcohols, lanolin alcohols linoleate, lanolin alcohols ricinoleate, acetate of lanolin alcohols ricinoleate, acetate of ethoxylated alcohols-esters, hydrogenolysis of lanolin, ethoxylated hydrogenated lanolin, ethoxylated sorbitol lanolin, and liquid and semisolid lanolin absorption bases and the like; 11. polyhydric alcohol esters, such as, ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol polyfatty esters, ethoxylated glyceryl monostearate, 1,2-butylene glycol monostearate, 1,2-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters; 12. wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate; 13. beeswax derivatives, e.g. polyoxyethylene sorbitol beeswax; 14. vegetable waxes including carnauba and candelilla waxes; 15. phospholipids such as lecithin and derivatives; 16. sterol including cholesterol and cholesterol fatty acid

esters; 17. amides such as fatty acid amides, ethoxylated fatty acid amides, solid fatty acid alkanolamides.

Humectants may be added to the composition to increase the effectiveness of the emollient, to reduce scaling, to stimulate removal of built-up scale and improve skin feel. By way of example only, suitable humectants include polyhydric alcohols, such as glycerol, polyalkylene glycols, alkylene polyols their derivatives, propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,3-butylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol and the like. The amount of humectant may be in the range of about 0.5-30 wt% and preferably between 1-15 wt%.

In topical skin care applications, a variety of active substances may be advantageously employed. By way of example only suitable active agents which may be incorporated into the cosmetic composition include anti-aging active substances, anti-wrinkle active substances, hydrating or moisturizing or slimming active substances, depigmenting active substances, substances active against free radicals, anti-irritation active substances, sun protective active substances, anti-acne active substances, firming-up active substances, exfoliating active substances, emollient active substances, and active substances for the treating of skin disorders such as dermatitis and the like.

By way of example only, in the case of hydration, one or more moisturizers may be used, such as glycerin or urea, in combination with one or more precursor agents for the biosynthesis of structural proteins, such as hydroxyproline, collagen peptides and the like.

By the way of example only, in case of slimming, at least one ketolytic agent or an alpha-hydroxyacid such a salicylic acid or 5-n-octanoicsalicylic acid may be used in combination with at least on liporegulating agent such as caffeine.

By way of example only, in the case of depigmentation, at least one keratolytic agent is used in combination with a depigmenting agent such as hydroquinone, tyrosinasee inhibitor (kolic acid), ascorbic acid, kojic acid and sodium metabisulfite an the like.

By way of example only, in the case of protection against free radical agents, vitamin E (against COO^\cdot radicals), superoxide dismutase (against O_2^\cdot free radicals) and sugar and caffeine (against OH^\cdot free radicals).

By way of example only, in the case of anti-aging, moisturizers, sunscreens,
5 alpha-hydroxyacids, salicylic acid or surface restructuring agents may be used in combination with enzymes for the repair of DNA, vascular protective agents or phospholipids rich in oligoelements and polyunsaturated fatty acids.

By way of example only, in the case of anti-acne agents, keratolytics, such as salicylic acid, sulfur, lactic acid, glycolic, pyruvic acid, urea, resorcinol and N-
10 acetylcysteine, and retinoids, such as retinoic acid and its derivatives may be used.

By way of example only, in the case of anti-inflammation, non-steroidal anti-inflammatory agents (NSAIDS) may be used, such as propionic acid derivatives, acetic acid, fenamic acid derivatives, biphenylcarboxylic acid derivatives, oxicams, including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, benoxaprofen,
15 flurbiprofen, fenbufen, ketoprofen, indoprofen, piroprofen, carprofen, and bucloxic acid and the like.

By way of example only, in the case of antibiotics and antimicrobials may be included in the composition of the invention. Antimicrobial drugs preferred for inclusion in compositions of the present invention include salts of β -lactam drugs,
20 quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, triclosan, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, hexamidine isethionate, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole and
25 amantadine and the like.

By way of example only, in the case of sunscreen protection, suitable agents include 2-ethylhexyl p-methoxycinnamate, 2-ethylhexyl N,N-dimethyl-p-aminobenzoate, p-aminobenzoic acid, 2-phenyl p-methoxycinnamate, 2-ethylhexyl octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-t-
30 butyldibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-

5 methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof and the like. The sunscreens disclosed therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range. These sunscreens provide higher efficacy, broader UV absorption, lower skin penetration and longer lasting efficacy relative to conventional sunscreens. Generally, the sunscreens can comprise from about 0.5% to about 20% of the compositions useful herein. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF). SPF is a commonly used measure of photoprotection of a sunscreen against erythema.

By way of example only, in the case of sunless tanning agents include dihydroxyacetone, glyceraldehyde, indoles and their derivatives, and the like.

15 The composition may include cleansing surfactants. Cleansing surfactants are cationic, anionic, amphoteric or non-ionic surfactants which are water-soluble and produce a consumer-acceptable amount of foam. Nonionic surfactants are well-known materials and have been used in cleansing compositions. Therefore, suitable nonionic surfactants include, but are not limited to, compounds in the classes known as alkanolamides, block copolymers of ethylene and propylene, ethoxylated alcohols, 20 ethoxylated alkylphenols, alkyl polyglycosides and mixtures thereof. In particular, the nonionic surfactant can be an ethoxylated alkylphenol, i.e., a condensation product of an alkylphenol having an alkyl group containing from about 6 to about 12 carbon atoms in either a straight chain or branched chain configuration with ethylene oxide, the ethylene oxide being present in an amount equal to at least about 8 moles ethylene oxide per mole of alkylphenol. Examples of compounds of this type include 25 nonylphenol condensed with about 9.5 moles of ethylene oxide per mole of phenol; dodecylphenol condensed with about 12 moles of ethylene oxide per mole of phenol; dinonylphenol condensed with about 15 moles of ethylene oxide per mole of phenol; octylphenol condensed with about ten moles of ethylene oxide per mole of phenol; and 30 diisooctyl phenol condensed with about 15 moles of ethylene oxide per mole of

phenol.

A wide variety of acids, bases, buffers, and sequestrants can be utilized to adjust and/or maintain the pH and ionic strength of the compositions useful in the instant invention. Materials useful for adjusting and/or maintaining the pH and/or the ionic strength include sodium carbonate, sodium hydroxide, hydrochloric acid, phosphoric acid, sulfuric acid, acetic acid, sodium acetate, sodium hydrogen phosphate, sodium dihydrogen phosphate, citric acid, sodium citrate, sodium bicarbonate, triethanolamine, EDTA, disodium EDTA, tetrasodium EDTA, and the like.

The polymer network may be useful as a solubilization agent in cosmetic and personal care applications. A self-assembling system comprising the reversibly gelling polymer network exhibits thermogelation, pH sensitivity, and the ability to solubilize hydrophobic agents in aqueous media. When poloxamer is copolymerized with poly(acrylic acid) (PAA) according to the invention, the resulting copolymer network is bioadhesive and can be applied in a number of therapies. The materials described in this invention combine "reverse" thermoviscosification mucoadhesion, solubilization of hydrophobic and difficult to manage moieties, easy formulation, and protection of agents from degradation to provide a superior medium for cosmetic and personal care products.

The reversible viscification of the polymer network at elevated temperatures makes the materials ideal for use as thickening agents in cosmetic and personal care products at any temperature above the transition. Another use of the "thickening" of solutions containing the polymer network as a thickener supplement in emulsions. Currently emulsifiers are often negatively effected by increased temperatures. An additive with reverse thermal viscification properties, however, would react in exactly the opposite way, increasing its ability to emulsify as it gained three-dimensional structure upon heating above its transition temperature.

In the applications where the reversibly gelling polymer composition can act as a surfactant, the polymer network will have the ability to act as a primary emulsifier without any (or with very little) addition of traditional surfactant. The responsive polymer network will also act as a stabilizer for oil-soluble ingredients that would

conventionally need to be solubilized by oils in formulation. The hydrophobic portion of the polymer network (PPO) forms domains which act as reservoirs for an oil-soluble or hydrophobic additive, such as an oil droplet, as is illustrated in Figure 9.

5 These two features of the material of the invention would enable it to be used as a base in a cosmetic formulation that would be non-greasy due to lack of oils, such as petrolatum and mineral oil. The increase in viscosity above the transition temperature adds structure and yield value to the water phase and results in a highly stable emulsion.

10 Thus, poloxamer:poly(acrylic acid) polymer network compositions are valuable materials in the formulation of cosmetic and personal care products. In particular, they may be useful as rheology modifiers, provide a cushioning effect on the skin, offer barrier properties and controlled release of actives. In addition, the polymer composition may serve as a surfactant and is compatible with most ingredients used in the cosmetic industry.







15 The above properties of the poloxamer:poly(acrylic acid) polymer network provides a cosmetic composition that spreads evenly and smoothly and which leaves a lubricious feel to the skin. A sensory evaluation was conducted with seven random volunteers in order to determine the sensory effect of a cream formulation on the skin. An oil-free cosmetic formulation was prepared substantially as set forth in Example 20 33(b) and was compared to Nivea Oil Free, a product of Beiersdorf of Germany. Volunteers placed unmarked samples on the skin and evaluated the formulation based upon its feel and texture. The samples were rated on a scale of 1 (bad) to 5 (good). The oil-free cosmetic formulation of the present invention scored equally to the Nivea Oil Free moisturizing product. Both samples scored a 3.5 on the rating scale.

25 The observed thermal behavior of the reversibly gelling polymer network suggests that the increase in viscosity is due to aggregation of the hydrophobic portion of the poloxamer at the transition temperature which, because of bonding with the poly(acrylic acid) component, serve as temporary cross-links which physically bridge adjacent chains of poly(acrylic acid) to provide a viscous gel-like extended polymer 30 structure. The aggregation process may be understood as occurring as shown in Figure

10, in which a backbone 20 represent poly(acrylic acid), a thin band 24 represents the hydrophobic poly(propylene) glycol region of the poloxamer and a thick band 26 represents the hydrophilic poly(ethylene glycol) region of the poloxamer. Below the transition temperature, the polymer network is randomly arranged, as is shown in Figure 10(a). At or above the transition temperature, the hydrophobic regions 24 associate to form aggregations or micelles 28, as is shown in Figure 10(b). The association increases the effective molecular weight of the polymer network composition with the corresponding increase in viscosity.

A general method of making the poloxamer:PAA polymer network compositions of the present invention comprises solubilization of the poloxamer in acrylic acid monomer, followed by polymerization of the monomer to PAA. Polymerization may be accomplished by addition of a polymerization initiator or by irradiation techniques. The initiator may be a free radical initiator, such as chemical free radical initiators and uv or gamma radiation initiators. Conventional free radical initiators may be used according to the invention, including, but in no way limited to ammonium persulfate, benzoin ethyl ether, benzyl peroxide, 1,2'-azobis(2,4-dimethylpentanitrile) (Vazo 52) and azobisisobutyronitrile (AIBN). Initiation may also be accomplished using cationic or ionic initiators. Many variations of this methods will be apparent to one skilled in the art and are contemplated as within the scope of the invention. For example, the poloxamer component may be dissolved in an acrylic acid/water mixture instead of pure monomer. It may be desirable to remove unreacted monomer and/or free poloxamer from the resultant polymer network. This may be accomplished using conventional techniques, such as, by way of example, dialysis or sohxlet extraction.

Without intending to be bound by a particular mechanism or structure, the following scheme represents a possible chemical mechanism for the formation of the system here described. These mechanisms are presented by way of explanation and are no way limiting of the invention. It is contemplated that these or other mechanistic routes may in fact occur in the formation of the polymer network of the present invention.

I. InitiationII. Hydrogen AbstractionIII. Chain TransferIV. PropagationV. Side Chain Branching Off AA BackboneVI. AA Branching off Poloxamer BackboneVII. Homogenous TerminationVIII. Heterogenous Termination with bonding of Pluronic to PAA

The scheme for bonding of poloxamer to acrylic acid may involve initiation (eq 1), hydrogen abstraction from the propylene or ethylene moiety of the poloxamer (eq 3), and attachment to acrylic acid via addition across the unsaturated bond (eq 10). Propagation (eq 8) leads to the final PAA.

Alternatively, the mechanism may proceed by initiation according to eqs. (1) and (2), propagation to form PAA (eq.8), a chain transfer reaction to generate a reactive poloxamer moiety (eq. 5), followed by addition of the reactive poloxamer

moiety to the unsaturated bond of acrylic acid (eq. 10) and subsequent propagation of the PAA chain.

Thus the polymer network may include a plurality of poly(acrylic acid)) units bonded to a single poloxamer unit or, alternatively, a plurality of poloxamer units bound to a single PAA backbone. Combinations of these alternatives are also a possibility.

Reverse phase polymerization may be used to prepare polymer network beads by dispersion of the poloxamer and acrylic acid monomer mixture in a nonpolar solvent such as hexane or heptane. The aggregating polymer/monomer solution is dispersed with agitation in the nonpolar solvent in order to suspend droplets of the solution. Polymerization of the monomer is initiated by conventional means (i.e., addition of a initiator or irradiation) in order to polymerize the monomer and form responsive polymer network beads. See, U.S.S.N. 08/276,532 filed July 18, 1995 and entitled "Useful Responsive Polymer Gel Beads" for further information on the preparation of polymer gel beads, herein incorporated by reference. Such a method may be particularly desirable to provide a heat sink for the heat generated in the exothermic polymerization reaction.

The polymer network complexes and aqueous gelling solutions of the present invention may be understood with reference to the following examples, which are provided for the purposes of illustration and which are in no way limiting of the invention.

Example 1 This example describes the synthesis of a polymer network and an aqueous responsive polymer network solution prepared using a triblock polymer of poly(ethylene glycol) and poly(propylene glycol), Pluronic® F27 polyol, and poly(acrylic acid). This example also characterizes the gelation and the physical properties of the resultant polymer network.

Synthesis. Block copolymer of poly(propylene glycol) (PPG) and poly(ethylene glycol) (PEG) having triad ABA structure $(\text{PEG})_A(\text{PPG})_B(\text{PEG})_A$ (Pluronic® F127 NF polyol, Poloxamer 407 NF polyol, where "F" means Flakes, "12" means 12X300=3600 - MW of the PPG section of the block copolymer, "7" PEG in

the copolymer is 70 wt%, and nominal molecular weight is 12,600) from BASF (3.0 g) was dissolved in 3.0 g acrylic acid (Aldrich). This represents a substantially 1:1 weight ratio of Pluronic® F127 polyol and poly(acrylic acid). The solution was deaerated by N₂ bubbling for 0.5 h and following addition of 100 ml of freshly prepared saturated solution of ammonium persulfate (Kodak) in deionized water was kept at 70 °C for 16 h resulting in a transparent polymer.

Viscosity measurements. A known amount of the resultant polymer was suspended in 100 ml deionized water into which NaOH was added. Following swelling for 3 days while stirring, the pH of the resulting fine suspension was adjusted to 7. Samples of 15 ml each were taken, and pH in each vial was adjusted to desired value by addition of 1 M HCl or NaOH. Samples were then kept overnight and their viscosities were measured at different temperatures using Brookfield viscometer using either an SC4-18 or an SC4-25 spindle.

A control experiment was done with a physical blend of Pluronic® F127 polyol and poly(acrylic acid) (MW 450,000) available from Aldrich. Pluronic® F127 polyol and poly(acrylic acid) were dissolved together in deionized water at 1 wt% total polymer concentration and the resultant solution was adjusted to pH 7, stirred and kept in refrigerator. The responsiveness of the polymer network composition and the physical blend to temperature and pH is illustrated in Figs. 1, 11 and 12. Figs. 1 and 2 clearly demonstrate that the synthetic route outlined above resulted in a polymer network system that is sensitive to pH and temperature of the environment. Note that the liquid-gel transition is very sharp, occurring over a very small temperature change or pH (see, Figure 11). Figure 12 is a viscosity vs. temperature graph comparing the gelling characteristics of the responsive polymer network composition and the physical blend. The blend prepared by physically mixing of the triblock PEG/PPG/PEG polymer and poly(acrylic acid) did not exhibit viscosifying effect either as a function of temperature or pH.

It was generally observed that 0.5-5 wt% polymer network compositions made of Pluronic® F127 polyol and poly(acrylic acid) viscosify at temperatures of around 30 °C and higher if pH is adjusted to 6 or higher. The gelling effect was observed in

polymer network compositions standing 3 months or longer. Repeated heating and cooling of responsive polymer network compositions did not cause deterioration of the polymer network or the gelling effect. Solutions of either Pluronic® F127 polyol or poly(acrylic acid) (1-5 w% in water, adjusted to pH 6 or higher) or physical blends of the two lacked the reverse thermal gelling effects found for polymer network compositions.

Example 2. This example describes a standard operating procedure for the manufacture of the reversible gelling polymer network.

The procedure is based upon a 50 liter production. A NaOH solution was prepared by dissolving 131.8 g NaOH pellets in 131.8 mL DI water (50% solution). The NaOH was allowed to dissolve completely. The NaOH solution will be used to convert a percentage of the acrylic acid to sodium acrylate in situ. Acrylic acid monomer (4 kg) is charged into a monomer feed tank and agitated at 250 rpm. NaOH is added slowly. The precipitate formed as the acrylic acid is neutralized to sodium acrylate is allowed to dissolve. Pluronic® F127 (3.5 kg) is slowly added to the monomer feed tank. Pluronic® F127 is dissolved under continued agitation. Norpar 12 (a refined C-12 alkane) is added to the reaction vessel (37 L). The mixture is agitated at 100 rpm. Stabilizer solution of Ganex V-126 is prepared in 2L Norpar 12 and added to the reactor under agitation.

A reaction vessel was degassed using a nitrogen sparge introduced from the bottom of reactor and was continued throughout the reaction. Initiator (13.63 g Lauryl peroxide and 4.23 g Vazo 52 in 0.7 kg acrylic acid monomer) is introduced into the monomer solution. The monomer solution was transferred to the reaction vessel. Agitation was increased to 150 rpm. Nitrogen sparging continued for an additional 20 minutes and then heating began. Heating began at a rate of 0.5-1.0 °C/min up to 75 °C. The reaction began to exotherm at about 45-50 °C and is allowed to continue without cooling until a maximum is reached. It is then cooled to 75 °C using forced cooling. The reaction continued for 12 hours and was then cooled to 35 °C. The slurry was transferred into pails and the polymer beads were allowed to settle.

The slurry was filtered through Buchner Funnels with filter paper (11 µm pore

size) until the bulk of the Norpar had been removed from the beads. The beads were washed three times with heptane. The filtered beads were transferred to a Pyrex drying tray and spread on the tray in a uniform layer. The beads were dried under vacuum for 4 hours at 40-50 °C. The dried beads were analyzed as follows.

5 Elemental analysis. The elemental analysis was performed by Quantitative Technologies, Inc., Whitehouse, NJ using a Perkin Elmer 2400 CHN Elemental Analyzer. Analysis provided C (52.49%), H (7.50%), N (< 0.05%), the balance assumed to be oxygen (39.96%).

10 Thermal Gravimetric Analysis (TGA). The TGA method was performed by Massachusetts Material Research, Inc., West Boylston, MA using a Dupont TGA model 295. The assay was run using a temperature ramp from 30 to 500 °C/min. The resolution for the system was set to 4 (1.0 °C/min for all slope changes). The data was analyzed using the first derivative of the curve and using maxima and minima to mark transitions. The moisture content was also calculated in this manner. The first
15 derivative yielded three maxima. The first transition (moisture) was 3.0% by weight, the second transition was 14.0% by weight and the third was 67.02% by weight. Residue (15.98% remained).

20 Molecular weight determination by gel permeation chromatography (GPC). The molecular weight was determined by GPC on a Hewlett Packard 1100 Liquid Chromatography system with a Viscotech T60 Triple Detector system. Three Waters Ultrahydrogel columns, 1000, 500 and 250 Å, were used for the separation. The
25 mobile phase was 0.1M NaNO₃ and 0.01M K₂HPO₄ salt solution, pH adjusted with phosphoric acid to a pH of 8.0 ± 0.1. The flow rate for the separation was 0.9 mL/min. The column temperature was maintained at 15 °C. The injection volume for the assay was 50 µL. A PEG molecular weight standard of 23,000 Daltons was used
30 to align the detectors. The result for the assay were:

M_n: 341,700 Daltons

M_p: 1,607,000 Daltons

M_w: 2,996,000 Daltons

30 Free poloxamer determination by GPC. The amount of free (unbound)

poloxamer in the polymer matrix was determined using the above GPC method and comparing the poloxamer peaks to that of a standard poloxamer solution. The typical result is approximately 18-22% free poloxamer by weight.

5 The effect of both the bonded and non-bonded poloxamer on the gelation properties of the responsive polymer network has been determined by extraction of the non-bonded poloxamer from the material. Such extraction studies have established that the graft co-polymer alone exhibits the characteristic reverse thermal gelation of the composition; however, the presence of non-bonded poloxamer component modulates the gelation process. The non-bonded poloxamer component
10 can affect the temperature of transition (from liquid to gel) and the degree of transition and assists in a more controlled and reproducible transition.

Bound poloxamer determination by ethylene oxide (EO) titration. The EO titration was performed as follows. A 5 gm sample of the product polymer was extracted in dichloroethane for three hours at reflux temperatures. The solid is
15 removed and dried under a vacuum for 12 hours at room temperature. The dry material is then analyzed using ASTM method D 2959-95, "Standard Test Method for Ethylene Oxide Content". The amount of EO in the sample is related to the amount of poloxamer bound to the polymer. The typical result is approximately 15 % by weight of EO.

20 The relative amount of free poloxamer may be varied dependent upon the relative proportions of starting materials and the method of polymerization. Although the residual solids presumably contain only poloxamer which is bonded to the poly(acrylic acid), i.e., a graft co-polymer, the material still shows strong viscosification when it is neutralized and dissolved in water. However, the
25 temperature of viscosification is increased substantially and the degree of viscosification per gram of total solids is increased by removal of free poloxamer. Thus, the free poloxamer plays a role in modifying the extent and temperature of viscosification. The poloxamer undergoes conformational changes and changes to the critical micelle concentration as a function of temperature. The poloxamer will
30 change from an open, non-aggregated form to a micellular, aggregated form with

changes in temperature.

Residual acrylic monomer determination by gas chromatography (GC). The residual acrylic acid monomer was determined by GC analysis using a Hewlett Packard GC 5890A, using a HP-FFDAP-TPA 10 m x 0.53 mm x 1 μ m column. The sample was extracted and run in methanol. Using an internal standard ratio, the sample was compared to a one point calibration. The typical results for this assay were below 70 ppm acrylic acid monomer.

Residual Norpar solvent by GC. The residual Norpar in the sample was determined by GC using the above method and comparing the Norpar peaks to that of a standard. The typical results were below 1.5 wt%.

UV-vis spectrum. Optical clarity data of UV-vis spectrophotometer was obtained. A 1.0% solution in water was prepared and measured at 420 nm. Transmittance (%) was typically greater than 90%.

Differential scanning calorimetry (DSC). The DSC was performed by Massachusetts Material Research, Inc., West Boylston, MA using a temperature ramp from 30 to 350 °C at 5 °C/min. The resolution for the system was set to 4 (1.0°C/min for all slope changes). The assay yielded one endothermic event at 265 °C, typically 270 J/g.

Examples 3-9. This example describes the synthesis of a several reversible thermal gelling polymer network prepared using a variety of poloxamers and poly(acrylic acid). The gelation and the physical properties of the resultant polymer network compositions are reported in Table 2.

Table 2.

example	poloxamer	poloxamer composition	polox- amer: PAA	trans. temp.	comments
3	Pluronic® F88 Prill polyol	2400 MW PPG; 80 wt% PEG; nominal MW 11,400	1:1	48 °C	viscosity response curve shown in Figure 13
4	Pluronic® F127 NF polyol	3600 MW PPG; 70 wt% PEG; nominal MW 12,600	1:1	30 °C	pentaerythritol triallyl ether crosslink agent used
5	Pluronic® P104 polyol	3000 MW PPG; 40 wt% PEG; nominal MW 5,900	1:1	28 °C	viscosity response curve shown in Figure 14
6	Pluronic® P123 polyol	3600 MW PPG; 30 wt% PEG; nominal MW 5,750	1:1	25 °C	viscosity response curve shown in Figure 15
7	Pluronic® F127/Pluronic® F108 polyol blend (1:1)	as above	1:1.7	42 °C	polymer solid formed. dried; resolubilized in neutralizing solution
8	Pluronic® F88 polyol	as above	1:1.7	80 °C	polymer solid formed. dried; resolubilized in neutralizing solution
9	Pluronic® F127/Pluronic® F88 polyol blend (1:1)	as above	1:1.7	85 °C	polymer solid formed. dried; resolubilized in neutralizing solution

Example 10. The following example demonstrates the effect of hydrophilic/hydrophobic ratio on the gelling temperature. Polymer network compositions were prepared from the following poloxamers shown in Table 3.

Table 3. Composition of poloxamers investigated.

triblock polyol polymer composition	MW of PPG block	wt% of PEG block
PT03 (PEG) ₁₇ (PPG) ₃₆ (PEG) ₁₇	3250	50
PT04 (PEG) ₂₅ (PPG) ₄₆ (PEG) ₂₅	3250	40
PT05 (PEG) ₁₆ (PPG) ₄₆ (PEG) ₁₆	3250	30

Table 3 shows that in this series, the fraction of PEG is reduced when the molecular weight of the PPG block is kept constant. Linse (*Macromol.* 26:4437-4449 (1993)) report phase diagrams for these copolymers in water were calculated and it was shown that two-phase boundaries corresponding to the beginning of aggregation are almost unaffected by the molecular mass, given a constant PEG/PPG ratio, whereas these boundaries shifted to lower temperature as the PEG content of the polymer is reduced at constant mass. The strong dependence of the PEG/PPG ratio is a consequence of the differing solubilities of PEG and PPG in water at the elevated temperatures. Thus one would suppose that aggregation that causes viscosification in the responsive polymer network composition should shift to lower temperature as PEG fraction decreases.

The poloxamer (3.0 g) was dissolved in 3.0 g acrylic acid. The solution was deaerated by N₂ bubbling for 20 min. and following addition of the 100 :1 of freshly prepared saturated solution of ammonium persulfate in deionized water was kept at 70°C for 16 h resulting in a strong whitish polymer. A sample of the polymer obtained (0.4 g) was suspended in 40 ml deionized water into which NaOH was added. Suspended responsive polymer network particles were allowed to dissolve under constant stirring. The resulting 1 wt% polymer network solutions were subjected to the viscosity measurement at shear rate of 132 or 13.2 sec⁻¹ using a SC4-18 spindle. It can be seen from Figure 16 that, firstly, viscosity of the 1 wt%

responsive polymer network solutions before viscosification (at 20-24°C) decreases in the series $(\text{PEG})_{37}(\text{PPG})_{56}(\text{PEG})_{17}(\text{F103}) > (\text{PEG})_{25}(\text{PPG})_{56}(\text{PEG})_{25}(\text{F104}) > (\text{PEG})_{16}(\text{PPG})_{56}(\text{PEG})_{16}(\text{F105})$ and, secondly, the temperature at which gelation shifts from about 45°C for $(\text{PEG})_{37}(\text{PPG})_{56}(\text{PEG})_{17}$ to about 35°C for

5 $(\text{PEG})_{25}(\text{PPG})_{56}(\text{PEG})_{25}$ and $(\text{PEG})_{16}(\text{PPG})_{56}(\text{PEG})_{16}$. Both results are in excellent agreement with the theory set forth in Linse.

Example 11. The following example is related to release of an active agent from a poloxamer:poly(acrylic acid) polymer network. Drug loading and kinetics of release of the protein hemoglobin from poloxamer:poly(acrylic acid) polymer network is described.

10 Synthesis. Pluronic® F127 (3.0 g) was dissolved in 3.0 g acrylic acid. The solution was deaerated by N₂ bubbling for 0.5 h and following addition of 100 Fl of freshly prepared saturated solution of ammonium persulfate (Kodak) in deionized water was kept at 70°C for 16 h resulting in a transparent polymer. The resultant responsive polymer network obtained (5 g) was suspended in 95 ml deionized water into which NaOH was added. The resulting suspension was allowed to swell for 7 days.

Hemoglobin loading and release. A 5 wt% responsive polymer network composition (3 g) was allowed to swell for 16 h in 10 ml of 0.25 mg/ml solution of human hemoglobin (Sigma) in deionized water adjusted to pH 8. The resulting mixture was well shaken and placed into the feed chambers of customized vertical, static, Franz-like diffusion cells made of Teflon. The feed and receiver chambers of the diffusion cells were separated by mesh screens (# 2063). The receiver chamber was continuously stirred by a magnetic bar. The cells were allowed to equilibrate to either 25 or 37°C (in an oven). The feed and receiver phases consisted of 1 g of the hemoglobin-loaded responsive polymer network and 6 ml of phosphate-buffered saline (pH 7.4), respectively. In the control experiment, the feed phase was made of 1 g of 0.25 mg/ml hemoglobin solution. After the feed solution had been loaded into the cell, the kinetic time commenced. Samples of the receiver phase were withdrawn from time to time and their absorbance was measured spectrophotometrically at 400 nm.

To calculate hemoglobin concentrations, corresponding calibration curves (absorbance in PBS versus hemoglobin concentration) were generated. The results of the kinetic experiment are presented in Figure 17. It can be seen that the rate of hemoglobin release from the polymer network was substantially lowered at 37°C when compared to that at 25°C, because of viscosity increase in the polymer network at elevated temperatures (see Figure 1). The protein released from the polymer network composition still retained its native structure, as was determined by comparison of uv-vis spectra of release hemoglobin and natural hemoglobin.

Example 12. The following example is related to release of an active agent from a poloxamer:poly(acrylic acid) polymer network. Drug loading and kinetics of release of the protein lysozyme from a polymer network is reported.

Lysozyme loading and release. A 5 wt% responsive polymer network composition (3 g) was allowed to swell for 16 h in 10 ml of 1 mg/ml solution of chicken egg-white lysozyme (Sigma) and 1.5 mg/ml sodium dodecyl sulfate (Aldrich) in deionized water adjusted to pH 8.5. The resulting mixture was well shaken and placed into the feed chambers of customized vertical, static, Franz-like diffusion cells made of Teflon. The feed and receiver chambers of the diffusion cells were separated by mesh screens (# 2063). The receiver chamber was continuously stirred by a magnetic bar. The cells were allowed to equilibrate to either 25 or 37°C (in an oven). The feed and receiver phases consisted of 1 g of the lysozyme-loaded responsive polymer network and 6 ml of phosphate-buffered saline (pH 7.4), respectively. In the control experiment, the feed phase was made of 1 g of 1 mg/ml lysozyme solution. After the feed solution had been loaded into the cell, the kinetic time commenced. Samples were withdrawn and their absorbance measured spectrophotometrically at 280 nm. A calibration curve was prepared for lysozyme concentration ranging from 0 mg/ml to 0.5 mg/ml in phosphate buffered saline. The results of the kinetic experiment are presented in Figure 18. It can be seen that the rate of lysozyme release from the responsive polymer network composition was substantially lowered at 37°C when compared to that at 25°C, because of viscosity increase in responsive polymer network at elevated temperatures (see Figure 1).

In order to demonstrate the retention of the enzymatic activity of lysozyme, the lysozyme released from the responsive polymer network composition was assayed using *Micrococcus lysodeikticus* cells and compared to that of original lysozyme. The enzymatic activity of lysozyme was the same, within the error of the assay (15%), as that of the original lysozyme. Control without lysozyme in presence of sodium dodecyl sulfate did not show any appreciable lysis of the cells.

Example 13. The following example is related to release of an active agent from a poloxamer:poly(acrylic acid) polymer network. Drug loading and kinetics of release of insulin from a responsive polymer network composition is reported.

Insulin loading and release. A 5 wt% responsive polymer network composition (3 g) was allowed to swell for 16 h in 10 ml of 5 mg/ml solution of bovine Zn^{2+} -insulin (Sigma) in deionized water adjusted to pH 7. The resulting mixture was well shaken and placed into the feed chambers of customized vertical, static, Franz-like diffusion cells made of Teflon. The feed and receiver chambers of the diffusion cells were separated by mesh screens (# 2063). The receiver chamber was continuously stirred by a magnetic bar. The cells were allowed to equilibrate to either 25 or 37°C (in an oven). The feed and receiver phases consisted of 1 g of the insulin-loaded responsive polymer network and 6 ml of phosphate-buffered saline (pH 7.4), respectively. In the control experiment, the feed phase was made of 1 g of 5 mg/ml insulin solution. After the feed solution had been loaded into the cell, the timing commenced. Samples were withdrawn and their absorbance was measured spectrophotometrically at 280 nm. A calibration curve was prepared for insulin concentration ranging from 0 mg/ml to 1.25 mg/ml in phosphate buffered saline. The results of the kinetic experiment are presented in Figure 19. The rate of insulin release from responsive polymer network was substantially lowered at 37°C when compared to that at 25°C, because of viscosity increase in responsive polymer network at elevated temperatures (see Figure 1).

Example 14. This example demonstrates the preparation of a sterile reversibly gelling polymer network aqueous composition and the stability of the composition to sterilization. The polymer network is prepared as described in Example 1, except that

the composition is prepared at 2 wt% Pluronic® F127 polyol/poly(acrylic acid). After dissolution of the 2 wt% polymer network in water, the viscosity is measured. The composition then is sterilized by autoclaving at 121°C, 16 psi for 30 minutes.

5 Viscosity is determined after sterilization. The corresponding curves for viscosity (a) before and (b) after sterilization are shown in Figure 20 and establish that minimal change in the viscosity profile of the material has occurred with sterilization.

Examples 15-30. These examples show additives which may be used to affect the transition temperature overall viscosification of the polymer network composition.

10 A 1 wt% polymer network was prepared in deionized water at pH 7 in which a variety of additives were included in the composition. The effect of the additive was determined by generation of a Brookfield viscosification curve. Results are reported in Table 4.

Table 4.

Example No.	Additive (wt%)	Effect of additive on:	
		transition temp. (°C)	final viscosity (% change)
15	1,2-methyl pyrrolidone (5)	I (1.8)	N
16	Rhodapex CO-436 (2)	I (1.6)	N
17	Dow Corning 190 (2)	I (5)	I (150)
18	isopropyl alcohol (0.5)	I (3.1)	I (45)
19	Pluronic® L122 (1)	D (4.4)	D (13)
20	Pluronic® F88 (1)	N	I (41)
21	Tween 80 (0.5)	N	I (18)
22	Germaben® II (1)	D (9)	I (100)
23	Iconol NP-6 (1)	D (9)	I (500)
24	Plurafac C-17 (0.5)	I (5.2)	D (36)
25	Dow Corning 193 (0.75)	I (4.1)	D (12)
26	glycerin (5)	D (2)	N
27	UC 50-HB- 170/EO/PO random copolymer (0.5)	N	N
28	PVP K15 (1)	N	N
29	MAPTAC (1)	N	D (8)
30	potassium chloride (0.25)	N	D (34)

I = increase; D = decrease; and N = no change

Example 31. Because of the surfactant nature of the polymer network composition coupled with the gelation effect of the polymer network composition, it is possible to prepare formulation which are 100% water-based, but which are lubricous and thick.

Formulations including a nonionic surfactant formulation: An O/W (oil-in-water) emulsion was made by combining the following ingredients utilizing conventional mixing techniques:

Table 5.

Ingredient	% w/w
10 % wt. 1:1 responsive polymer network as prepared in Example 1	20.0
Emulsifying Wax NF ¹	2.5
Mineral Oil	5.0

¹ Polowax available from Croda

Into a vessel equipped with a high efficiency homogenizer, the formula amount of all ingredients is added, water is added to 100% w/w and allowed to mix to homogeneity. This formulation contains a nonionic surfactant and gives an emulsion that is fluid at room temperature but viscifies above 32°C.

Formulations including a cationic surfactant formulation: An O/W (oil-in-water) emulsion was made by combining the following ingredients utilizing conventional mixing techniques:

Table 6.

Ingredient	% w/w
10 % wt. 1:1 responsive polymer network as prepared in Example 1	20.0
Behentrimonium Methosulfate (and) Cetearyl alcohol ¹	2.5
Mineral Oil	5.0

¹ Incroquat Behenyl TMS available from Croda

Into a vessel equipped with a high efficiency homogenizer, the formula amount

of all ingredients is added and allowed to mix to homogeneity. This formulation contains a cationic surfactant and gives an emulsion that is fluid at room temperature but viscifies above 32°C.

Formulations including an anionic surfactant formulation: An O/W (oil-in-water) emulsion was made by combining the following ingredients utilizing conventional mixing techniques:

Table 7.

Ingredient	% w/w
10 % wt. 1:1 responsive polymer network as prepared in Example 1	20.0
Cetearyl Phosphate (and) Cetearyl alcohol ¹	2.5
Mineral Oil	5.0

¹ Crodafos CES available from Croda

Into a vessel equipped with a high efficiency homogenizer, the formula amount of all ingredients is added, water is added to 100% w/w and allowed to mix to homogeneity. This formulation contains a anionic surfactant and gives an emulsion that is fluid at room temperature but viscifies above 32°C.

Example 32. Acne Medication: An oil-free, clear, anti-acne treatment is made by combining the following ingredients utilizing conventional mixing techniques:

Table 8.

Ingredient	% w/w
10 % wt. 1:1 responsive polymer network prepared as in Example 1	20.0
Glycerin USP	5.0
Salicylic Acid	2.0
DL-Panthenol	0.5
Germaben® II ¹	0.1
Disodium EDTA	0.2
USP Purified Water	72.2

¹ Germaben®II available from Sutton Laboratories

To one vessel, equipped with a Lightnin' Mixer with a 3 blade paddle prop,

the full amount of USP Purified Water to 100% w/w is added. While maintaining the temperature, with moderate to vigorous mixing, the formula amount of Disodium EDTA, Citric Acid, DL-Panthenol, Glycerin, Salicylic Acid, and Germaben® II is added. These materials are allowed to dissolve at 50°C. After dissolution, the vessel
5 is then cooled to 20°C. To another vessel, equipped with a high efficiency homogenizer, the formula amount of responsive polymer network is added. The responsive polymer network vessel is then cooled to 4°C. After cooling, while vigorously homogenizing, the contents of the first vessel is added to the second vessel, and allowed to mix to homogeneity.

10 The composition displays a flowable clear jelly appearance with excellent spreadability and absorption characteristics at room temperature, and after heating the formulation to 32°C, the composition thickens to a gel-like consistency.

Example 33. (a) Oil-free Moisturizer (formulation I): An oil-free, lubricous
15 moisturizer was made by combining the following ingredients utilizing conventional mixing techniques:

Table 9.

Ingredient	% w/w
10% wt 1:1 responsive polymer network as prepared in Example 1	20.0
Glycerin USP	5.0
PPG-2 Myristyl Ether Propionate	3.0
DL-Panthenol	0.5
Germaben® II ¹	0.1
Disodium EDTA	0.2
Citric Acid	0.01
USP Purified Water	71.19

¹ Germaben® II available from Sutton Laboratories

The above ingredients were added and processed as described above for the acne composition. The composition displayed a flowable creamy lotion appearance with excellent emolliency, spreadability and absorption characteristics at room temperature. After heating the formulation to above 26°C, the composition thickened to a gel-like consistency. The viscosity vs. temperature curve is shown in Figure 21 and demonstrates that addition of adjuvants to the composition significantly enhances the responsive polymer network maximum viscosity (> 900,000 cps). The use of the poloxamer:poly(acrylic acid) polymer network in the formulation also imparts a unique viscosification effect after application to the skin, which is not evident in typical commercial O/W emulsion formulations (See, Figure 21b).

(b) Oil-free Moisturizer (formulation II): An oil-free, lubricious moisturizer was made by combining the following ingredients utilizing conventional mixing techniques:

Table 10.

Ingredient	% w/w
1:1 polymer network as prepared in Example 1	2.0
Glycerin USP	5.0
Carbopol 980	1.0
D-panthenol, propylene glycol	1.0
Preservative	1.0
Hydrolyzed protein (and) hyaluronic acid	0.5
Sodium hydroxide	0.2
USP Purified Water	90

The above ingredients were added and processed as described above for the acne composition. The composition displayed a flowable creamy lotion appearance with excellent emolliency, spreadability and absorption characteristics at room temperature. After heating the formulation to 26°C, the composition thickens to a gel-like consistency. The addition of adjuvants to the composition significantly enhances the polymer network maximum viscosity.

Example 34. Sunscreen Lotion. An oil-free, lubricious sunscreen lotion was made by combining the following ingredients utilizing conventional mixing techniques:

Table 11.

Ingredient	% w/w
1:1 polymer network as prepared in Example 1	2.0
Glycerin USP	8.0
Carbopol 980	1.0
Parsol MCX	7.0
Myristyl Ether Propionate	5.0
Preservative	1.0
Cyclomethicone	1.0
Sodium hydroxide	0.2
USP Purified Water	74

The above ingredients were added and processed as described above for the acne composition. The composition displayed a flowable creamy lotion appearance with excellent emolliency, spreadability and absorption characteristics at room temperature. After heating the formulation to above 26°C, the composition thickened to a gel-like consistency. The addition of adjuvants to the composition significantly enhances the polymer network maximum viscosity.

Example 35. Facial mask. A face mask was made by combining the following ingredients utilizing conventional mixing techniques:

Table 12.

Ingredient	% w/w
1:1 polymer network as prepared in Example 1	1.0
Polyvinyl alcohol	6.0
Polyvinylpyrrolidone (20%)	5.0
D-panthenol, propylene glycol	1.25
Propylene glycol	1.25
USP Purified Water	85.5

The above ingredients were added and processed as described above for the acne composition. The composition displayed a flowable creamy lotion appearance with excellent emolliency, spreadability and absorption characteristics at room temperature. After heating the formulation to above 26°C, the composition thickened to a gel-like consistency. The addition of adjuvants to the composition significantly enhances the polymer network maximum viscosity.

Example 36. Facial toner. A face mask was made by combining the following ingredients utilizing conventional mixing techniques:

Table 13.

Ingredient	% w/w
1:1 polymer network as prepared in Example 1	0.01
Hydroxyethyl cetyldimonium phosphate	1.00
PEG-40 hydrogenated castor oil	2.00
D-panthenol, propylene glycol	0.50
Glycerin	2.00
Witch hazel extract	5.00
USP Purified Water	88.49

The above ingredients were added and processed as described above for the acne composition. The composition displayed a flowable appearance with excellent emolliency, spreadability and absorption characteristics at room temperature. After heating the formulation to above 26°C, the composition thickened to a gel-like consistency. The addition of adjuvants to the composition significantly enhances the polymer network maximum viscosity.

Example 36. Solubilization studies of model hydrophobic agents in the poloxamer: poly(acrylic acid) polymer network: estradiol and progesterone. This example is presented to demonstrate the solubilization of a hydrophobic agent in the polymeric network. Progesterone and estradiol were used as the hydrophobic agents in this model solubilization study.

Acrylic acid (99%), fluorescein (98%), β -estradiol (98%), and progesterone (98%) were all obtained from Aldrich and used as received. Pluronic® F127 NF was obtained from BASF. Poly(oxyethylene-b-oxypropylene-b-oxyethylene)-g-poly(acrylic acid) copolymers (responsive polymer network) were synthesized by free-radical polymerization of acrylic acid in the presence of poloxamer as described above. The polymer network copolymers discussed here were composed of about 1:1 ratio of PAA to poloxamer. The rheological properties of polymer network were assessed using LVDV-II+ and RVDV-II+ Brookfield viscometers. The microscopic light scattering of 21 nm poly(styrene) latex particles in deionized water and 1 w% reversibly gelling polymer network was measured using He-Ne laser as described previously (See, Matsuo, E.S., Orkisz, M., Sun, S.-T., Li, Y., Tanaka, T., Macromolecules, 1994, 27, 6791). The solubility of fluorescein and hormones in aqueous solutions was measured by the equilibration of excess solubilizate with the corresponding solution following removal of undissolved species by centrifugation and filtration. Hydrophobic agents were assayed spectrophotometrically at 240 (progesterone) or 280 nm (estradiol), or by using 70/30 w/w H₂SO₄/MeOH (Tsilifonis-Chafetz reagent). In vitro hormone release studies were conducted using thermostatted, vertical Franz cells. Spunbonded polypropylene microfilters (micron retention, 15-20) were used as a membrane separating feed and receiver phases in

Franz cells. The responsive polymer network, water, ethanol, and 20% PEG in water were observed to wet the membrane. The receiver solutions consisted of 20 w% PEG in water (pH 7) and were stirred by magnetic bars. The feed phases composed of responsive polymer network were loaded with either estradiol or progesterone. Each hormone was dissolved in ethanol and the resulting solution was added into the responsive polymer network.

Equilibrium solubility vs. temperature plots for estradiol and progesterone (partition coefficient octanol/water (P) 7200 and 5888, respectively, in aqueous solutions of Pluronic® F127 polyol and responsive polymer network are presented in Figure 22. It can be seen that increasing temperature and concentration (C) of polymers in the solution raises the amount of the hormone dissolved. In Figure 22a, vertical lines represent critical micellar temperatures (CMT) for corresponding Pluronic F127 polyol solutions. It is interesting to note that the slope of the solubility-temperature plots increased as temperature reached CMT, indicating that solubilization in the Pluronic solutions was predominantly due to the formation of micelles. Similar trend was observed in the responsive polymer network solutions. The S values in 5% aqueous solutions of branched PAA did not exceed 15 and 40 $\mu\text{g/mL}$ at 60 °C for estradiol and progesterone, respectively. The solubility values found for responsive polymer network were the same as S in parent Pluronic solutions of equivalent concentrations. Therefore, it may be suggested that solubilization behaviors of the responsive polymer network are governed by the properties of the poloxamer incorporated into it. Thermodynamic parameters of the solubilization process with responsive polymer network were calculated using the same approximations as in the micellar solubilization with Pluronic polyols. See, Saito, Y., Kondo, Y., Abe, M., Sato, T., Chem.Pharm.Bull., 1994, 42, 1348. Namely, partition coefficient P was estimated from equilibrium solubilities of estradiol in responsive polymer network and water:

$$P = S_{SH}/S_w \quad (13)$$

by extrapolating the solubility plots of the steroid in Figure 22 to 100 % responsive polymer network. Using P values obtained from data in Figure 23, we calculated the

standard free energy change (ΔG), standard enthalpy of solubilization (ΔH), and standard entropy of solubilization (ΔS) using the following expressions:

$$\Delta G = -RT \ln P; \Delta H = -R \Delta \ln P / \Delta(1/T); \Delta S = (\Delta H - \Delta G)/T \quad (14)$$

Thermodynamic parameters obtained along with P values are given in Table 13.

- 5 Apparent partition coefficients and thermodynamic parameters for solubilization of estradiol by responsive polymer network.

Table 13.

T, K	P=SSH/S	ΔG kJ/mol	ΔH kJ/mol	ΔS J/mol
277	490	-14.3	4.72	68.6
293	520	-15.2		52.0
310	660	-16.7		53.9
323	660	-17.4		54.0
333	660	-18.0		54.0

Negative ΔG values indicate spontaneous solubilization at all temperatures, whereas positive ΔH shows that the solubilization was endothermic, similar to the solubilization of estriol, as well as indomethacin, by the poloxamer. Notably, ΔS of solubilization was always positive, suggesting that the more ordered water molecules surrounding hydrophobic estradiol molecules moved to the less ordered bulk phase when the estradiol was transferred to the hydrophobic core of PPG segments in responsive polymer network. The aggregation of the PPG segments at elevated temperatures provides not only temporary cross-linking in the gel, but also a thermodynamically "friendly" environment for the hydrophobic drugs. Indeed, one can express the free energy of formation of the aggregate core-water interface in responsive polymer network as:

$$\Delta G = [\sigma P_w(1 - \phi) + \sigma W_D \phi](4\pi R^2/n) \quad (15)$$

where σP_w and σW_D are the interfacial tensions between pure PPO polymer and water and between water and the drug, respectively; ϕ is the volume fraction of the drug within PPO core; R is the effective radius of the core, and n is the aggregation number.

Equation (3) shows that solubilization of a hydrophobic drug of high σ_{WD} should increase the stability of the aggregate. The solubilization process was found to decrease the critical micellization concentration and substantially increase the micellar core radius in Pluronic surfactants (Hurter, P.N. *et al.*, "In Solubilization in Surfactant Aggregates", Christian, S.D., Ed., Marcel Dekker, New York, 1995). A similar trend is indicated by the lowering the onset of gelation of the responsive polymer network upon solubilization of fluorescein (LogP 2.1) (Figure 24). The solubilization of hydrophobic drugs by responsive polymer network, analogous to the micellar solubilization of drugs by poloxamer, suggests that the responsive polymer network can be an effective vehicle in drug delivery.

Our *in vitro* study of hormone release from responsive polymer network shows an increase in the initial transport rate with either decreasing total polymer concentration in the formulation or decreasing temperature (Figure 25). These effects are related to the changes in macroscopic viscosity of the responsive polymer network, which erodes more rapidly from the feed phase through the membrane into the receiver compartment as the viscosity decreases (Figure 26). The degree of the responsive polymer network erosion was measured by weighing hormone-loaded responsive polymer network before and after kinetic experiment.

Figure 27 shows that the relative amount of progesterone penetrating into the receiver phase decreased 4-fold with the increase of total polymer concentration, whereas the total relative amount of progesterone stayed almost constant as total polymer concentration in the responsive polymer network increased. This result shows the existence of two routes of transport of hydrophobic drugs in our model system. Firstly, the drug incorporated into aggregates within the responsive polymer network system can flow through the membrane along with the erosion of the responsive polymer network; secondly, the drug not associated with the responsive polymer network aggregates can diffuse out of the responsive polymer network in the feed phase. The second process should not be related to the viscosity of the responsive polymer network. Indeed, the dynamic light scattering experiment shows no dramatic change of diffusivity of poly(styrene) latex particles in the responsive polymer

network as temperature rises thereby increasing macroscopic viscosity more than 10-fold (Figure 28). This result indicates that the viscosity of the responsive polymer network is essentially unaffected on the microscopic scale.

5 Appendix A attached.

APPENDIX A

Cosmetic Bench Reference Function Definitions

Abrasive: abrades, smoothes, polishes	Emollient: softens, smoothes skin
Absorbent powder: takes up liquids, sponge-like action	Emulsifier: a surface-active agent (surfactant) that promotes the formation of water-in-oil or oil-in-water emulsions
Absorption base: forms water-in-oil emulsions	Enzymes: complex proteins produced by living cells that catalyze biochemical reactions at body temperature
Acidulent: acidifies, lowers pH, neutralizes alkalis	Fiber: strands of natural or synthetic polymers; for instance, cotton, wool, silk, nylon, polyester
Amphoteric: capable of reacting chemically either as an acid or a base; amphoteric surfactants are compatible with anionic and cationic surfactants	Film former: solution of a polymer that forms films when the solvent evaporates after application to a surface
Analgesic: relieves pain	Fixative: fixes or sets perfumes; retards evaporation; promotes longer lasting aroma
Antacid: neutralizes stomach acidity	Flavor: imparts a characteristic taste (and aroma) to edible foods and drinks; sometimes used in lip products
Antibacterial: destroys/inhibits the growth/reproduction of bacteria	Foam booster: enhances quality and quantity of lather of shampoo
Anti-caking: prevents or retards caking of powders; keeps powders free-flowing	Foamer: a surface-active agent (surfactant) that produces foam; an emulsion of air-in-water
Anti-dandruff: retards or eliminates dandruff	Foam stabilizer: see Foam booster
Antifoam: suppresses foam during mixing	Fungicide: inhibits or destroys growth of fungi
Anti-inflammatory: reduces, suppresses, counteracts inflammation	Gellant: a gelling agent; forms gels; includes a wide variety of materials such as polymers, clays and soaps
Anti-irritant: reduces, suppresses or prevents irritation	Glosser: furnishes a surface luster or brightness; usually used in lip or hair products
Antimicrobial: destroys, inhibits or suppresses the growth of microorganisms	Hair colorant: see Colorant
Antioxidant: inhibits oxidation and rancidity	Hair conditioner: see Conditioner
Antiperspirant: reduces or inhibits perspiration	Hair dye: imparts a new permanent or semi-permanent color to hair
Antipruritic: reduces or prevents itching	Hair-set polymer: polymer and/or resins used to maintain desired hair shape
Antiseptic: inhibits the growth of microorganisms on the skin or on living tissue	Hair-set resin: see Hair-set polymer
Antistat: reduces static by neutralizing electrical charge on a surface	Hair waving: see Reducing agent and Neutralizer
Astringent: contracts organic tissue after application	Humectant: absorbs, holds and retains moisture
Binder: promotes cohesion of powders	Hydrotrope: enhances water solubility
Bleaching agent: lightens color, oxidizing agent	Intermediate: basic chemicals which are chemically modified to obtain the desired function
Botanical: natural plant derivative	Lathering agent: a surface active agent (surfactant) that forms a foam or lather on mixing with air in solution; see also Foamer
Buffer: helps maintain original pH (acidity or basicity) of a preparation	Lubricant: reduces friction, smoothes, adds slip
Carrier: a vehicle or base used for a preparation	Moisture barrier: retards passage of moisture or water
Chelate: form a complex with trace-metal impurities, usually calcium or iron	Moisturizer: aids in increasing the moisture content of the skin through humectant or barrier action
Colorant: adds color, may be a soluble dye or an insoluble pigment	Neutralizer: an oxidizing agent used in hair waving that stops the action of the reducing agent and re-establishes the disulfide linkages in hair
Conditioner: improves condition of skin and hair	Oil absorbent: see Absorbent powder
Coupling agent: aids in solubilization or emulsification of incompatible components	Ointment base: an anhydrous mixture of oleaginous components used as a vehicle for medicaments
Decolorant: removes color by adsorption, bleaching or oxidation	Opacifier: opacifies clear liquids or solids
Denaturant: used to denature ethyl alcohol	Oxidant: oxidizing agent, neutralizes reducing agents, bleaching agent
Dental powder: powdered dentifrice	Pearlant: imparts a pearlescent texture and luster
Deodorant: desroys, masks or inhibits formation of unpleasant odors	Perfume solvent: see Solvent and Solubilizer
Depilatory: removes hair chemically	
Detergent: a surface-active agent (surfactant) that cleans by emulsifying oils and suspends particulate soil	
Disinfectant: destroys pathogenic microorganisms	
Dispersant: promotes the formation and stabilization of a dispersion or suspension	
Dye stabilizer: see Stabilizer	

Peroxide stabilizer: see Stabilizer

Pigment: a finely powdered insoluble substance used to impart color, luster or opacity

Plasticizer: plasticizes (makes more flexible) polymeric films or fibers

Polish: smoothest; adds gloss and luster

Polymer: a very high molecular weight compound consisting of repeating structural units

Powder: a solid in the form of fine particles

Preservative: protects products from spoilage by microorganisms

Propellant: pressurized gas in a container used to expel the contents when pressure is released by opening a valve

Protein: naturally occurring complex combinations of amino acids

Reducing agent: reduces a chemical compound usually by donating electrons; neutralizes oxidizing agents

Refatting agent: adds oily materials to the surface of substrates, e.g., skin and hair

Resin: nonvolatile solid or semisolid organic substances obtained from plants as exudates or prepared by polymerization of simple molecules

Sequestrant: forms coordination complexes with multivalent positive ions

Silicone: polymeric organic silicon compounds which are water resistant

Skin protectant: protects skin from environmental

Solubilizer: solubilizes, usually into aqueous vehicles, normally insoluble materials, such as fragrances, flavors, oils, etc.

Solvent: usually liquids capable of dissolving other substances

Stabilizer: added to stabilize emulsions and/or suspensions

Stimulant: produces a temporary increase in the functional activity of an organism or any of its parts

Surfactant (surface-active agent): lowers surface tension between two or more incompatible phases: soaps, detergents, wetting agents, solubilizing agents and emulsifying agents are typical surfactants; surfactants are classified as anionic, cationic, nonionic and amphoteric; anionic surfactants are negatively charged, cationic surfactants have no electrical charge

Suspending agent: keeps finely divided solid particles in suspension

Sweetener: sweetens to provide a more pleasant taste

Tanning accelerator: accelerates the tanning of skin

Thickener: thickens or increases viscosity/consistency

Thixotrope: the property of certain gels and emulsions of becoming more fluid or less viscous when shaken or stirred

UV absorber: used as a sunscreen and to protect preparations from degradation by UV radiation


UVA absorber: absorbs in the range 320-400 nanometers (nm)

UVB absorber: absorbs in the range 290-320 nanometers (nm)

Wax: any of numerous substances of plant, animal or synthetic origin that contain principally esters of higher fatty acids and higher fatty alcohols; free fatty alcohols, fatty acids and hydrocarbons may also be present; waxes derived from petroleum products are mainly high-molecular-weight hydrocarbons

Wetting agent: a surface-active agent (surfactant) that lowers the surface and interfacial tension, facilitating the wetting of surfaces

From the Editors of *Cosmetics & Toiletries* magazine



Hair Care

Adsorption of cationic polymers

— D. Coddard and R. Schmitt

Ceramides — D. Braidà et al

Melanins — K.C. Brown and G. Prota

Men's hair coloring — S. Cusperton

Skin permeation of hair dyes — H. Beck et al

African-American hair — A. Syed et al

Ethnic hair care — A. Syed

Hair curl relaxers — P. Obukowho and M. Birman

Cysteine waving lotions — A. Iwasaki

Keeping VOCs under control — S. Price

Soft cover

69 pages

Published 1996

ISBN: 0-931710-47-2

International airmail: \$6

Price: \$64



A New Book Compiles The Very Best in Hair Research

The Applied Research Series pulls together the very best papers on scientific advances and research. Featuring the latest technical information, formulating advice and ingredient reviews, this book is an ideal addition to your research library.

I would like _____ copies of the Hair Care book (Payment required with order)

☐ Payment enclosed, US\$64 per copy. (US funds drawn on a US bank only.)

Please add US\$6 per copy for all international airmail orders.

☐ Charge my



Card Number

Expiration Date

Name as it appears on card (Please Print)

Signature

Name (Please Print)

Company

Address

City/State/Postal Code

Country

Telephone (include area code)

Fax (include area code)

e-mail address

CBR

All publications sold on a non-refundable basis • Prices subject to change

Send your order with payment to: Allured Publishing Corp.

362 S. Schmale Road • Carol Stream IL 60188 USA • Phone: 708/653-2155 • Fax: 708/653-2192

Functions

Abrasive

Adzuki beans
Almond (Prunus amygdalus) meal, shell granules
Aluminum silicate
Apricot (Prunus armeniaca) kernel powder, shells
Hydrated silica
Jojoba (Buxus chinensis) seed powder
Luffa cylindrica
Olive stone granules
Oyster shell powder
Peach (Prunus persica) pit powder
Peach (Prunus persica) stone granules
Polyethylene
Polyethylene HEC granules
Polyethylene (oxidized, P. spheres)
Polystyrene
Pumice
Rice (Oryza sativa) bran
Silica and S. colloidal
Sodium chloride
Walnut (Juglans regia) shell powder

Absorption base

1,2,6-Hexanetriol
Kaolin
Petrolatum
Rice (Oryza sativa) starch
Soy (Glycine soja) sterol
Zeolite

Absorbent powder

Corn (Zea mays) starch
Maltodextrin
Nylon-12
Oat (Avena sativa) bran, flour, meal
Zeolite

Acidulant

Acetic acid
Citric acid
Fumaric acid
Glutaric acid
Glycolic acid

Hydrochloric acid
Lactic acid
Nitric acid
Phosphoric acid
Sodium bisulfate
Sulfuric acid
Tartaric acid

AHA

Apple (Pyrus malus) extract
Apricot (Prunus armeniaca) kernel powder
Citric acid
Ethyl lactate
Glycolic acid
Lactic acid
Malic acid
Sodium lactate
Tartaric acid

Antiacne

Clays (white, yellow, red, green, pink)
Perfluorodecalin
Salicylic acid
Sulfur

Anti-aging

Basil (Ocimum basilicum) extract
Carrot (Daucus carota) extract
Catalpa kaempferia extract
Ceramide 33 (liquid soy extract)
Crataegus cuneata extract
Eugenia jambolana extract
Fomes fomentarius extract
Fomitopsis pinicola extract
Ganoderma lucidum oil
Ginseng (Panax ginseng) extract
Hyaluronic acid
Hydrolyzed serum protein
Hydrolyzed soy flour
Isachne pulchella extract
Lactoferrin
Lady's Thistle (Silybum marianum) extract
Ligusticum jeholense extract

Manne collagen
Mushroom (Coniopus versicolor) extract
Musk rose (Rosa moschata) oil
Perfluorodecalin
Quaternium-51
Rubus thunbergii extract
Serum protein
Senecalia mucallii extract
Tricholoma matsutake extract

Antibacterial

Ammonium iodide
Chlorhexidine
Chlorhexidine diacetate, C. digluconate
Chlorhexidine dihydrochloride
Chlorphenesin
Hexamidine diisethionate
Hexidine
Iceland moss (Cetraria islandica) extract
Lactoferrin
Lauralkonium bromide, L. chloride
Laurtrimonium chloride
Laurylpyridinium chloride
Mauriella armata extract
Mushroom (Cordyceps sabolifera) extract
Orange blossom extract
Orange (Citrus aurantium dulcis) peel extract
PEG-42 Ebiniko ceramides extract
Peppermint (Mentha pipenta) extract
Phlodonendron (Phellodendron amurense) extract
Pine (Pinus sylvestris) needle extract
Polymethoxy bicyclic oxazolidine
Quaternium 73
Rubus thunbergii extract
Tea tree (Melaleuca alternifolia) oil
Triclocarban
Undecylenic acid

Anticaking

Aluminum starch octenylsuccinate
Calcium stearate
Distarch phosphate
Hydrated silica



The Only Form of Gentle Efficacy



Natural • Odorless • Biodegradable • Custom colors • Mild or aggressive particles

CORPORATE OFFICES • 1151 N. Fiesta Blvd. • Gilbert, AZ 85233 • (602) 545-7000 • Fax (602) 892-3000 • New York • Canada • Europe • Asia

FLORATECH
The Natural Solution

BROOKS INDUSTRIES inc.

Cosmetic Ingredients & Ideas®

70 Tyler Place, South Plainfield, NJ 07080 USA
Tel 908-561-5200 Fax 908-561-9174

Multifruit®

Alpha Hydroxy Acids



**UNIVERSAL
PRESERV-A-CHEM INC.**

297 North 7th Street, Brooklyn, New York 11211
Phone: 718-782-7429
Fax: 718-782-8109

*We Deliver
Quality*

"ALPHA HYDROXY ACIDS"
Glycolic Malic
Lactic Citric
Tartaric Ascorbic
Allantoin Aloe Vera

Functions

Kaolin
Magnesium myristate, M. silicate
Polyethylene, micronized
Silica silylate
Sodium aluminum silicate
Zinc stearate

Anticaries agent

Cetylamine hydrofluoride
Olaflur
Sodium fluoride
Stearyl trihydroxyethyl propylenediamine dihydrofluoride

Anticellulite

Aminophylline
Bladderwrack (*Fucus vesiculosus*) extract
Butcherbroom (*Ruscus aculeatus*) extract
Carcinia cambogia extract
Fomes tomentarius extract
Fomistopsis pinicola extract
Ivy extract
Mushroom (*Coniolum versicolor*) extract
TEA-hydroiodide
Tricholoma matsutake extract

Antidandruff

Burdock (*Arcium lappa*) extract
Chloroxylenol
Corydalis ambigua extract
Disodium undecylenamido MEA-sulfosuccinate
Ginger root extract
Inga edulis extract
Mauritiella armata extract
Myristalkonium saccharinate
PEG-6 undecylenate
Piroctone olamine
Resorcinol
Rosemary (*Rosmarinus officinalis*) extract
Sodium shale oil sulfonate
Stenocalyx micalii extract
Undecylenamide DEA
Willow (*Salix alba*) bark extract
Zinc pyrithione

Antifungal

Black walnut (*Juglans nigra*) extract
Coneflower (*Echinacea angustifolia*) extract
Orange blossom extract
Pfaffia paniculata extract

Anti-inflammatory

Allantoin polygalacturonic acid
Bisabolol
Black poplar (*Populus nigra*) extract
Brassica rapa-depressa extract
Butcherbroom (*Ruscus aculeatus*) extract
Calendula officinalis extract
Catalpa kaempferia extract
Celastrum paniculata extract
Ceramide 33 (liquid soy extract)
Chaparral (*Larrea mexicana*) extract
Coneflower (*Echinacea angustifolia*) extract
Coneflower (*Echinacea cyanus*) extract
Dipotassium glycyrrhizinate
Euphorbium fortunei extract
Euphrasia officinalis extract
Ficus racemosa extract
Golden seal (*Hydrastis canadensis*) root extract
Guaiaculene
Horse chestnut (*Aesculus hippocastanum*) extract
Jujube (*Zizyphus jujuba*) extract
Laminaria japonica extract
Licorice (*Glycyrrhiza glabra*) extract
Ligusticum jeholense, L. lucidum extract
Matricaria (*Chamomilla recutita*) extract
Melaleuca uncinata extract
Melia azadirachta extract

Mulberry (*Morus nigra*) extract
Niacinamide ascorbate
Orange (*Citrus aurantium dulcis*) peel extract
Orange blossom extract
Palmetto extract
Palmitoyl collagen amino acids
Passion flower (*Passiflora laurifolia*) fruit extract
Paulownia impenalis extract
Salicylic acid
Shea butter (*Butyrospermum parkii*)
Sodium carboxymethyl beta-glucan
Soy (*Glycine soja*) protein
Stearyl glycyrrhizinate
Stenocalyx micalii extract
Tocopheryl acetate, T. nicotinate
Trichomonas japonica extract
Willow (*Salix alba*) extract
Witch hazel (*Hamamelis virginiana*) extract
Withania somniferum extract
Yarrow (*Achillea millefolium*) extract
Zinc lactate

Anti-irritant

Acetyl monoethanolamine
Allantoin
Allantoin acetyl methionine, A. glycylmethionine acid
Azelaic acid
Betaine
Calendula officinalis extract
Cocamidopropyl betaine
Coceth-7 carboxylic acid
Cornflower (*Centaurea cyanus*) extract
Diisostearyl dimer diisoleate
Dipalmitoyl cysteine
Green tea extract
Hydrolyzed sweet almond protein
Hydroxypropyltrimonium gelatin
Lauroyl collagen amino acids
L-Lysine lauroyl methionine
Mallow extract
Matricaria (*Chamomilla recutita*) extract
Palmitoyl hydrolyzed milk protein
Palmitoyl hydrolyzed wheat protein
Palmitoyl keratin amino acids
PEG-12 palm kernel glycerides
PEG-28 glyceryl tallowate
PEG-30 glyceryl monococoate
PEG-60 almond glycerides
PEG-78 glyceryl cocoate
PEG-82 glyceryl tallowate
PEG-200 glyceryl tallowate
Propionyl collagen amino acids
PVP
Saccharomyces lysate extract
Sodium C12-15 parabens-15 sulfonate
Sodium lauroamphoacetate
Soy (*Glycine soja*) protein
Undecylenoyl collagen amino acids
Valerian (*Valeriana officinalis*) extract

Antimicrobial

Benzalkonium chloride
Benzoic acid
Benzyl alcohol
Bromochlorophene
2-Bromo-2-nitropropane-1,3-diol
Butylparaben
Capryloyl collagen amino acids
Capryloyl glycine, C. keratin amino acids
Caplan
Cetethyldimonium bromide
Cetyl pyridinium chloride
Chlorothymol
Chloroxylenol
Citron oil
Copper PCA
Dichlorobenzyl alcohol
Dilauryldimonium chloride

Domiphen bromide
Ethylparaben
Eucalyptus (*Eucalyptus globulus*) extract
Fennel (*Foeniculum vulgare*) extract
Garlic (*Allium sativum*) extract
Glyceryl caprylate, G. laurate
Hexamidine diisethionate
Hinokitiol
Honeysuckle (*Lonicera caprifolium*) extract
Lichen (*Usnea barbata*) extract
Myristalkonium chloride
Pentylene glycol
Phenethyl alcohol
Phenol
Phenoxyethanol
Phenoxyisopropanol
Phenyl mercuric acetate, P.m. benzoate, P.m. borate
o-Phenylphenol
Polymethoxy bicyclic oxazolidine
Potassium sorbate
Propylparaben
Ricinoleamodopropyltrimonium ethosulfate
Sage (*Salvia officinalis*) extract
Sodium benzoate, S. pyrithione
Sodium ricinoleate, S. shale oil sulfonate
Thimerosal
Thyme (*Thymus vulgaris*) extract
Thymol
Triclocarban
Triclosan
Undecylenamido propyltrimonium methosulfate
Undecylenic acid
Zinc oxide, Z. PCA
Zinc pyrithione, Z. undecylenate

Antioxidant

Ascorbic acid
A. polypeptide
Ascorbyl oleate, A. palmitate
Beta-carotene
BHA
BHT
t-Butyl hydroquinone
Dilauryl thiodipropionate
Dimyristyl thiodipropionate
Disodium EDTA
Distearyl thiodipropionate
Dodecyl gallate
EDTA
Erythorbic acid
Ferulic acid
Grape (*Vitis vinifera*) seed extract
Green tea extract
HEDTA
Hydroquinone
Hydroquinone-beta-D-glucopyranoside
p-Hydroxyanisole
Lactoferrin
Lysine PCA
Melanin
Methyl gallate
Niacinamide ascorbate
Nordihydroguaiaretic acid
Oat (*Avena sativa*) extract
Oryzanol
Pentasodium pentetate
Pentetic acid
Propyl gallate
Retinyl palmitate polypeptide
Rosemary (*Rosmarinus officinalis*) extract
Saccharomyces lysate extract
Sage (*Salvia officinalis*) extract
Sodium ascorbate, S. erythorbate
Sodium metabisulfite
Sodium selenate, S. sulfite
Superoxide dismutase
Tea (*Camellia sinensis*) extract
Tetrasodium EDTA
Tocopherol

Functions

Tocopherol acetate, E. linoleate
Wild marjoram (*Origanum vulgare*) extract
Yeast (*Saccharomyces cerevisiae*) extract (Faex)

Antiperspirant

Allantoin-aluminum chlorhydrate
Aluminum caprylyl hydrolyzed collagen
Aluminum chlorohydrate-gly, A. chloride
Aluminum chlorohydrate, A. chlorohydrate
Aluminum PCA, A. sesquichlorohydrate
Aluminum undecylenyl collagen amino acids
Aluminum zirconium pentachlorohydrate
Aluminum zirconium tetrachlorohydrate
Aluminum zirconium tetrachlorohydrate GLY
Aluminum zirconium trichlorohydrate
Aluminum-zirconium-glycine powder
Sage (*Salvia officinalis*) extract
Tomentil (*Potentilla erecta*) extract
Zirconium chlorohydrate

Antiseptic

Aluminum PCA
Azadirachta indica extract
2-Bromo-2-nitropropane-1,3-diol
Calendula amurensis extract
p-Chloro-m-cresol
Clove (*Eugenia caryophyllus*) oil
Crataegus cuneata extract
Dichlorobenzyl alcohol
Eniada phaseoloides extract
Eucalyptus (*Eucalyptus globulus*) extract
Gulden seal (*Hydrastis canadensis*) root extract
Hexachlorophene
Melia australasica, M. azadirachta extract
Methyl salicylate
Orange (*Citrus aurantium dulcis*) peel extract
Oxyquinoline sulfate
Pfafia paniculata extract
Potassium abietoyl hydrolyzed collagen
PVP-iodine
Silver nitrate
Sodium salicylate
Sterculia platantolia extract
Tea tree (*Melaleuca alternifolia*) oil
Tomentil (*Potentilla erecta*) extract
Xanthozylum bungeanum extract

Antistat

Acetamide MEA
Acetamidopropyl trimonium chloride
6-(N-Acetylamino)-4-oxyhexyltrimonium chloride
Alkyl dimethyl betaine
Babassuamidopropylalkonium chloride
Behenamidopropyl ethyldimonium ethosulfate
Behenamidopropyl hydroxyethyl dimonium chloride
Carboxymethyl chitin
Cetethyl morpholinium ethosulfate
Cetrimonium chloride
Chitin
Chitosan
Cocamidopropyl ethyldimonium ethosulfate
Cocodimonium hydroxypropyl hydrolyzed rice protein
Cocodimonium hydroxypropyl hydrolyzed soy protein
Dimethicone hydroxypropyl trimonium chloride
Dimethyl behenamine, D. cocamine
Dimethyl palmitamine, D. soyamine
Dimethyl tallowamine
Diolelamidodethyl hydroxyethylmonium methosulfate
Dipalmitoylethyl hydroxyethylmonium methosulfate
N-Dodecyl-N,N-dimethyl-N-(dodecyl acetate) ammonium chloride
Erucamidopropyl hydroxysulfate
Glyceryl monopyroglytate
Hydrogenated tallowamine oxide
Isostearamidopropyl dimethylamine

Lactamidopropyl trimonium chloride
Lauryldimonium hydroxypropyl hydrolyzed collagen
Linoleamidopropyl dimethylamine dimer diinoleate
Olealkonium chloride
PEG-2 cocamine
PEG-2 cocomonium chloride
PEG-2 oleammonium chloride
PEG-8 caprylic/capric glycerides
PEG-10 cocamine
PEG-15 soyamine
PPG-9 diethylmonium chloride
PPG-25 diethylmonium chloride
PPG-40 diethylmonium chloride
Propylene glycol stearate
Quaternium-26, -27, -53, -62, -72
Rapeseedamidopropyl benzylidimonium chloride
Rapeseedamidopropyl epoxypromyl dimonium chloride
Silica, colloidal
Sorbitan caprylate
N-Soy-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate
Soyethyl morpholinium ethosulfate
Soyethylidimonium ethosulfate
Stearalkonium chloride
Stearamidopropyl benzyl dimonium chloride
Stearamidopropyl ethyldimonium ethosulfate
Steatrimonium chloride
N-Stearyl-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate
Wheat germamidopropyl ethyldimonium ethosulfate

Astringent

Aluminum citrate, A. lactate
Astragalus sinicus extract
Astrocaryum murumuru, A. tucuma extract
Azadirachta indica extract
Azelaamide MEA
Bearberry (*Arctostaphylos uva-ursi*) extract
Birch (*Betula alba*) leaf extract
Catalpa kaempferia extract
Celastrus paniculata extract
Coccinea indica extract
Coffee (*Coffea arabica*) bean extract
Euphrasia officinalis extract
Euterpe precatoria extract
Evening primrose (*Oenothera biennis*) extract
Gentian (*Gentiana lutea*) extract
Geranium maculatum extract
Grape (*Vitis vinifera*) leaf extract
Henna (*Lawsonia inermis*) extract
Hierochloa odorata extract
Honeysuckle (*Lonicera caprifolium*) extract
Hops (*Humulus lupulus*) extract
Horsetail extract
Hypericum perforatum extract
Ivy extract
Juniperus communis extract
Kadsura heteroloca extract
Kola (*Cola acuminata*) extract
Lady's mantle (*Alchemilla vulgaris*) extract
Lemon (*Citrus medica limonum*) extract, peel extract
Lemon bioflavonoids extract
Lysimachia foenum-graecum extract
Magnolia spp. extract
Mauritia flexosa extract
Maximiliana regia extract
Melaleuca uncinata, M. wilsonii extract
Melia australasica extract
Nettle (*Urtica dioica*) extract
Oak (*Quercus*) bark extract
Ocimum basilicum, O. sanctum extract
Palmetto extract
Passion flower (*Passiflora laurifolia*) fruit extract
Plantain (*Plantago major*) extract
Polygonum multiflorum extract
Pterocarpus marsipianus extract
Raspberry (*Rubus*) extract

Sambucus nigra oil
Sanguisorba root extract
Selinum spp. extract
Shorea robusta extract
Tannic acid
Walnut (*Juglans regia*) leaf extract, oil
Wheat (*Triticum vulgare*) protein
White nettle (*Lamium album*) extract
Witch hazel (*Hamamelis virginiana*) extract
Xanthozylum bungeanum extract
Zinc lactate
Ziziphus jujuba extract

Binder

Aluminum starch octenylsuccinate
Boron nitride
C20-40, C30-50, C40-60 alcohols
Calcium stearate
Cellulose gum
Dihydroabietyl behenate
Diisostearyl malate
Dioctyl sebacate
Distarch phosphate
Ethylcellulose
Gellan gum
Hydrogenated jojoba oil
Isocetyl alcohol, I. palmitate
Isopropyl isostearate
Isostearyl erucate, I. isostearate
Isostearyl neopentanoate
Maltodextrin
Methylcellulose
Microcrystalline cellulose
Octyl palmitate
Octyldodecyl myristate
bis-Octyldodecyl stearoyl dimer diinoleate
Octyldodecyl stearoyl stearate
Oleyl oleate
PEG-20, -75, -150, -240, -350
Polydipentene
Polyethylene: P., micronized
PTFE
PVP
Sorbitol
Synthetic wax
Tapioca dextrin
Tridecyl behenate, T. neopentanoate
Tridecyl stearoyl stearate
Trisodium HEDTA

Biol. polymer

Distarch phosphate
Dog rose (*Rosa canina*) seed extract
Hydrogen peroxide
Kojic acid
Mulberry (*Morus nigra*) extract
Sanguisorba root extract

Botanical

Acacia
Acacia farnesiana extract
Agrimony (*Agrimonia eupatoria*) extract
Alder (*Alnus firma*) extract
Alfalfa (*Medicago sativa*) extract
Algae (*Ascomyllum nodosum*) extract
Algae (*Lithothamnium calcareum*) extract
Aloe barbadensis, A.b. extract
Aloe capensis extract
Alpine Veronica extract
Althea officinalis extract
Angelica archangelica extract
Anise (*Pimpinella anisum*) extract
Apple (*Pyrus malus*) extract
Apricot (*Prunus armeniaca*) extract
Arnica montana extract
Artemisia capillaris extract
Arichoke (*Cynara scolymus*) extract
Asafetida (*Ferula assa foetida*) extract
Asiasarum sieboldii extract

Functions

Asparagus officinalis extract	Cucumber (Cucumis sativus) extract	Jasmine (Jasminum officinale) extract
Astragalus sinensis extract	Cypress (Cupressus sempervirens) extract	Job's tears (Coix lacryma-jobi) extract
Avena (Cicium rivale) extract	Dandelion (Taraxacum officinale) extract	Jojoba (Buxus chinensis) seed powder
Avocado (Persea gratissima) extract	Date (Phoenix dactylifera) extract	Juniperus communis extract
Balm mint (Melissa officinalis) extract, oil extract	Dead Sea Mud, Salts	Kelp (Macrocystus pyrifera) extract
Banana (Musa sapientum) extract	Dog rose (Rosa canina) hips extract	Kiwi (Actinidia chinensis) fruit extract, seed oil
Barley (Hordeum vulgare) extract	Dyer's broom extract	Kola (Cola acuminata) extract
Basil (Ocimum basilicum) extract	Eleuthero ginseng (Acanthopanax senticosus) extract	Kramena mandra extract
Bearberry (Arctostaphylos uva-ursi) extract	Elm (Ulmus campestris) extract	Lady's mantle (Alchemilla vulgaris) extract
Bee pollen extract	Eucalyptus (Eucalyptus globulus) extract	Lady's Thistle (Silybum marianum) extract
Beet (Beta vulgaris) extract	Eucalyptus globulus oil	Laurel (Laurus nobilis) extract
Betaglucon	Eucommia ulmoides extract	Lavender (Lavandula angustifolia) extract, water
Bilberry (Vaccinium myrtillus) extract	Euphrasia officinalis extract	Lemon (Citrus medica limonum) extract, juice
Biotinonoids	Evening primrose (Oenothera biennis) extract, oil	extract, peel extract
Birch (Betula alba) bark extract, leaf extract	Everlasting (Helichrysum arvenarium) extract	Lemon bioflavonoids extract
Birch (Betula platyphylla japonica) extract	Fennel (Foeniculum vulgare) extract	Lemongrass (Cymbopogon schoenanthus) extract
Bitter orange (Citrus aurantium amara) extract, flower extract, peel extract	Fenugreek extract	Leopard flower (Belamcanda chinensis) root extract
Black cohosh (Cimicifuga racemosa) extract	Fermented rice (Oryza sativa) extract	Lettuce (Lactuca scariola sativa) extract
Black currant (Ribes nigrum) extract	Fern (Dryopteris filix-Mas) extract	Licorice (Glycyrrhiza glabra) extract
Black henna extract	Fig (Ficus carica) extract	Lilac (Syringa vulgaris) extract
Black poplar (Populus nigra) extract	Fir needle extract	Linden (Tilia argentea) extract
Black walnut (Juglans nigra) extract	Fumitory (Fumaria officinalis) extract	Linden (Tilia cordata) extract, water
Bladderwrack (Fucus vesiculosus) extract	Gardenia florida extract	Loquat (Eriobotrya japonica) leaf extract
Borage (Borago officinalis) extract	Garlic (Allium sativum) extract	Maidenhair fern extract
Buckthorn (Frangula alnus) extract	Gelidium cartilagineum	Magnolia kobus extract
Burdock (Arctium lappae) extract	Gentian (Gentiana lutea) extract	Mallow extract
Burdock (Arctium minus) root extract	Geranium maculatum extract	Mandragora officinarum extract
Burnet extract	Ginger root extract	Mannan
Butcherbroom (Ruscus aculeatus) extract	Ginkgo biloba extract	Mangold
Cabbage rose (Rosa centifolia) extract	Ginseng (Panax ginseng) extract	Marine salts
Calamus (Aconis calamus) extract	Glycyrrhetic acid	Mauritania (Chamomilla recutita) extract
Calendula officinalis extract	Glycyrrhizic acid	Meadowsweet (Spiraea ulmaria) extract
Caper (Capparis spinosa) extract	Glycyrrhizin, ammoniated	Melon (Cucumis melo) extract
Capsicum frutescens extract, C.F. oleoresin	Golden seal (Hydrastis canadensis) root extract	MEA iodine
Caraway (Carum carvi) extract	Goldthread (Coptis japonica) extract	Mistletoe (Viscum album) extract
Cartagenan (Chondrus crispus)	Gotu kola extract	Mugwort (Artemisia princeps) extract, water
Carrot (Daucus carota) extract	Grape (Vitis vinifera) distillate, extract	Mulberry (Morus alba) root extract
Carrot (Daucus carota sativa) oil	Grape (Vitis vinifera) leaf, seed extract	Mulberry (Morus bombycis) root extract
Cassia auriculata extract	Grape skin extract	Mushroom extract
Celandine (Chelidonium majus) extract	Grapefruit (Citrus grandis) peel extract	Myrrh (Commiphora myrrha) extract
Chamomile (Anthemis nobilis) extract, oil	Green bean (Phaseolus lunatus) extract	Nasturtium extract
Chaparral (Larrea mexicana) extract	Ground Ivy (Glechoma hederacea) extract	Neroli extract
Cherry (Prunus speciosa) leaf extract	Guarana (Paullinia cupana) extract	Nettle (Urtica dioica) extract
Cherry bark, C.b. extract	Harpagophytum procumbens extract	Oak (Quercus) bark extract
Chestnut (Castanea sativa) extract	Haviflower extract	Oak root extract
Chinese hibiscus (Hibiscus rosa-sinensis) extract	Hazel (Corylus avellana) nut extract	Oat (Avena sativa) bran, bran extract, flour, protein
Chlorella vulgaris extract	Henna (Lawsonia inermis) extract	Oat flower
Cimicifuga foetida rhizome extract	Hesperidin, H. methyl chalcone	Olive (Olea europae) extract, leaf extract
Cinchona succubina extract	Hibiscus sabdariffa extract	Onion (Allium cepa) extract
Citroflavonoid, water soluble	Hibiscus synacus extract	Orange blossom extract
Citrus bioflavonoid complex	High beta-glucan barley flour	Orange (Citrus aurantium dulcis) flower extract, peel extract
Clary extract	Honeysuckle (Lonicera caprifolium) extract	Pansy (Viola tricolor) extract
Clove (Eugenia caryophyllus) extract	Honeysuckle (Lonicera japonica) leaf extract	Papaya (Carica papaya) extract
Clover (Trifolium pratense) extract	Hops (Humulus lupulus) extract	Parsley (Carum petroselinum) extract
Cnidium officinale rhizome extract, C.O. water	Horse chestnut (Aesculus hippocastanum) extract	Passion flower (Passiflora laurifolia) fruit extract
Colfee (Coffea arabica) bean extract	Horseradish (Cochlearia armoracia) extract	Passionflower (Passiflora incarnata) extract
Colloidal oatmeal	Horsetail extract	Pea (Pisum sativum) extract
Coltsfoot (Tussilago farfara) leaf extract	Houttuynia cordata extract	Peach (Prunus persica) extract, leaf extract
Comfrey (Symphytum officinale) leaf extract	Hyacinth (Hyacinthus orientalis) extract	Pelargonium capitatum extract
Condurango extract	Hydrocotyl (Centella asiatica) extract	Pellitory (Panetaria officinalis) extract
Coneflower (Echinacea angustifolia) extract	Hydrolyzed oat protein, soy flour	Pennyroyal (Mentha pulegium) extract
Corallina officinalis	Hypericum perforatum extract	Peony (Paeonia albiflora) extract
Corchorus olitorius extract	Hyssop (Hyssopus officinalis) extract	Peony (Paeonia obovata) root extract
Coriander (Coriandrum sativum) extract	Indian cress (Tropaeolum majus) extract	Peppermint (Mentha piperita) extract, oil
Corn (Zea mays) cob powder, silk extract	Isodonis Japonicus extract	Perilla ocymoides extract
Corn poppy (Papaver rhoeas) extract	Ivy extract	Periwinkle (Vinca minor) extract
Cornflower (Centaurea cyanus) extract	Japanese angelica (Angelica acutiloba) extract, water	PEG-80 jojoba acid/alcohol
Couch (Agropyron repens) grass	Japanese hawthorn (Crataegus cuneata) extract	PEG-120 jojoba acid/alcohol
Crataegus monogyna extract		
Cribrum maritimum extract		

CAMPO Siddha Herbs Extracts

Jothi-Pul (Glow-grass) Siddha Extract for High content bio-available
Natural Radium for anti Kaposi Sarcoma Skin Treatment.
 Roma-Maram (Hairy Tree) Siddha Extract for **ANTI-SENSE DNA**
 Topical applications for HIV+ Lymph-nodes
 Siddha Extracts for post-Chemotherapy Skin-Damage Treatment



CAMPO RESEARCH



Level 36, Hong Leong Building,
 18 Raffles Quay, Singapore 0104
 Tel: (65) - 7653292 Full Colour Fax: (65) - 7653293
 PC - Video Teleconferencing (65) 7653292 - For Tech. Assistance.

Functions

<p> <i>Pfalfla paniculata</i> extract <i>Phellodendron amurense</i> extract Phospholipids Pimento (<i>Pimenta officinalis</i>) extract Pine (<i>Pinus sylvestris</i>) cone, needle extract Pineapple (<i>Ananas sativus</i>) extract Plantain (<i>Plantago major</i>) extract Pollen extract Pongamul Pona Cocos extract <i>Pueraria lobata</i> extract Queen of the meadow extract Quillaja saponaria extract Quince (<i>Pyrus cydonia</i>) seed extract Quinoa (<i>Chenopodium quinoa</i>) extract Rasperry (<i>Rubus</i>) extract <i>Rauwolfia</i> (<i>Serpentina</i>) extract Red clover <i>Rehmannia chinensis</i> extract Resharrow (<i>Ononis spinosa</i>) extract <i>Rhododendron chrysanthum</i> extract <i>Rhodophyceae</i> extract Rhubarb (<i>Rheum palmatum</i>) extract Rice (<i>Oryza sativa</i>) bran extract Rice fatty acid Rose (<i>Rosa multiflora</i>) extract Rosemary (<i>Rosmannus officinalis</i>) extract <i>Ruta tinctorum</i> extract Safflower (<i>Carthamus tinctorius</i>) extract Sage (<i>Salvia officinalis</i>) extract, water <i>Sambucus nigra</i> berry extract, extract Sandalwood (<i>Santalum album</i>) extract <i>Sanguinaria canadensis</i> extract <i>Saponaria officinalis</i> extract <i>Sasa veitchii</i> extract <i>Saxifraga sarmentosa</i> extract <i>Scabiosa arvensis</i> extract <i>Scutellaria baicalensis</i> root extract Silk extract Silver fir (<i>Abies pectinata</i>) extract Sisal (<i>Agave rigida</i>) extract Slippery elm extract Soapberry (<i>Sapindus mukurossi</i>) extract <i>Sophora angustifolia</i> extract <i>Sophora flavescens</i> root extract <i>Sophora japonica</i> extract Soybean (<i>Glycine soja</i>) extract Soy (<i>Glycine soja</i>) germ extract, protein, sterol Spearmint (<i>Mentha viridis</i>) extract, oil Spinach (<i>Spinacia oleracea</i>) extract <i>Spiraea ulmaria</i> extract Sunflower (<i>Helianthus annuus</i>) seed extract Sweet almond (<i>Prunus amygdalus dulcis</i>) extract Sweet cherry (<i>Prunus avium</i>) extract Sweet cicely (<i>Anthriscus cerefolium</i>) extract Sweet clover (<i>Melilotus officinalis</i>) extract Sweet violet (<i>Viola odorata</i>) extract <i>Swertia chirata</i> extract Tea (<i>Camellia sinensis</i>) extract Thistle (<i>Chicus benedictus</i>) extract Thyme (<i>Thymus vulgaris</i>) extract Tomato (<i>Solanum lycopersicum</i>) extract Tormentil (<i>Potentilla erecta</i>) extract Tuberose (<i>Polianthes tuberosa</i>) extract Turmeric (<i>Curcuma longa</i>) extract Valerian (<i>Valeriana officinalis</i>) extract Walnut (<i>Juglans regia</i>) extract, leaf extract Water Lily (<i>Nymphaea alba</i>) root extract Watercress (<i>Nasturtium officinale</i>) extract Wheat (<i>Triticum vulgare</i>) extract, protein Wheat (<i>Triticum vulgare</i>) germ extract Wheat bran lipids White ginger (<i>Hedychium coronatum</i>) extract White nettle (<i>Lamium album</i>) extract Wild agnemony (<i>Potentilla anserina</i>) extract Wild cherry (<i>Prunus serotina</i>) bark extract Wild indigo (<i>Baptisia tinctoria</i>) Wild marjoram (<i>Origanum vulgare</i>) extract Willow (<i>Salix alba</i>) bark extract, extract Willow (<i>Salix alba</i>) leaf extract Witch hazel (<i>Hamamelis virginiana</i>) extract Yarrow (<i>Achillea millefolium</i>) extract Yeast (<i>Saccharomyces cerevisiae</i>) extract (Faex) <i>Yucca vera</i> extract <i>Zanthoxylum piperitum</i> extract Zedoary (<i>Curcuma zedoaria</i>) oil </p>	<p> Wheat (<i>Triticum vulgare</i>) extract, protein Wheat (<i>Triticum vulgare</i>) germ extract Wheat bran lipids White ginger (<i>Hedychium coronatum</i>) extract White nettle (<i>Lamium album</i>) extract Wild agnemony (<i>Potentilla anserina</i>) extract Wild cherry (<i>Prunus serotina</i>) bark extract Wild indigo (<i>Baptisia tinctoria</i>) Wild marjoram (<i>Origanum vulgare</i>) extract Willow (<i>Salix alba</i>) bark extract, extract Willow (<i>Salix alba</i>) leaf extract Witch hazel (<i>Hamamelis virginiana</i>) extract Yarrow (<i>Achillea millefolium</i>) extract Yeast (<i>Saccharomyces cerevisiae</i>) extract (Faex) <i>Yucca vera</i> extract <i>Zanthoxylum piperitum</i> extract Zedoary (<i>Curcuma zedoaria</i>) oil </p> <p> Buffer Ammonium carbonate, A. phosphate Calcium hydroxide, C. phosphate Citric acid Ethanolamine HCl Glycine Phosphoric acid Potassium phosphate Potassium sodium tartrate Sodium acetate, S. citrate Sodium lactate, S. phosphate Succinic acid Tromethamine </p> <p> Carrier Acrylates copolymer, spherical powder Arginine Caprylic/capric triglyceride Caprylic/capric/lauric triglyceride Caprylic/capric/linoleic triglyceride Caprylic/capric/oleic triglycerides Ceteareth-20 Coconut (<i>Cocos nucifera</i>) oil Cyclodextrin Dipropylene glycol Glyceryl caprylate, G. caprylate/caprate Hydrated silica Liposomes Magnesium silicate Methyl propanediol PEG-8/SMDI copolymer Potassium chloride PPG-12/SMDI Copolymer PPG-51/SMDI Copolymer Propylene carbonate, P. glycol Serum albumin Sodium carboxymethyl beta-glucan Sodium chloride Sodium magnesium silicate Tapioca dextrin </p> <p> Chelators beta-Alanine diacetic acid Calcium disodium EDTA Disodium EDTA, -copper EDTA HEDTA Malic acid Monostearyl citrate Pentasodium pentetate Pentetic acid </p>	<p> Phytic acid Potassium aspartate Sodium aspartate Sodium dihydroxyethylglycinate Sodium hexametaphosphate Tetrahydroxypropyl ethylenediamine Tetrasodium EDTA Tripotassium EDTA Trisodium EDTA, HEDTA </p> <p> Cell stimulant <i>Aesculus chinensis</i> extract <i>Artemisia apiacea</i> extract <i>Astrocaryum muru</i>, A. tucuma extract <i>Bacris gasipaes</i> extract <i>Borjooa sorbilis</i> extract <i>Calendula amurensis</i> extract <i>Chrysanthemum monifolium</i> extract <i>Cocinea indica</i> extract Comfrey (<i>Symphytum officinale</i>) leaf extract Condurango extract Dandelion (<i>Taraxacum officinale</i>) extract <i>Echitea glauca</i> extract <i>Equisetum arvense</i> extract Eucalyptus (<i>Eucalyptus globulus</i>) extract <i>Eupatorium fortunei</i> extract <i>Euterpe precatoria</i> extract <i>Ficus racemosa</i> extract Glycoproteins <i>Hierochloa odorata</i> extract Horse chestnut (<i>Aesculia hippocastanum</i>) extract Inga edulis extract <i>Kadsura heteroloca</i> extract <i>Ligustrum lucidum</i> extract <i>Lysimachia foenum-graecum</i> extract <i>Mauritia flexosa</i> extract <i>Maximiliana regia</i> extract <i>Melaleuca bracteata</i>, M. symphyocarp extract <i>Nelumbium speciosum</i> extract <i>Ocimum basilicum</i> extract, O. santum extract <i>Paulownia imperialis</i> extract <i>Pfaffia</i> spp. extract <i>Pterocarpus marsupianus</i> extract <i>Rubus thunbergii</i> extract <i>Selinum</i> spp. extract <i>Shorea robusta</i> extract <i>Xanthoxylum bungeanum</i> extract </p> <p> Cleansing Birch (<i>Betula alba</i>) leaf extract Lemongrass (<i>Cymbopogon schoenanthus</i>) extract Oat (<i>Avena sativa</i>) bran extract Passion flower (<i>Passiflora laurifolia</i>) fruit extract Witch hazel (<i>Hamamelis virginiana</i>) extract Yarrow (<i>Achillea millefolium</i>) extract </p> <p> Conditioner Acetamide MEA 6-(N-Acetylamino)-4-oxyhexyltrimonium chloride Acrylamidopropyltrimonium chloride/acrylamide copolymer Adipic acid/dimethylaminohydroxypropyl diethylene triamine copolymer AMP-isostearoyl hydrolyzed wheat protein Apricot (<i>Prunus armeniaca</i>) kernel oil Behenalkonium chloride Behenamidopropyl dihydroxypropyl dimonium chloride Behenamidopropyl ethyldimonium ethosulfate Behenamidopropyl PG-dimonium chloride </p>
--	--	--

CAMPO Siddha Herb Extracts
CAMPO Rainforest Herb Extracts & Oils
CAMPO Australasian Herbs & Tea Tree Extracts
CAMPO Chinese & Japanese Herb Extracts


CAMPO RESEARCH

Level 38, Hong Leong Building,
16 Raffles Quay, Singapore 0104
Tel: (65) - 7653292 Full Colour Fax: (65) - 7653293
PC - Video Teleconferencing (65) 7653292 - For Tech. Assistance.

Functions

Behenamidopropyltrimethylamine betaine	Hydrolyzed sweet almond protein	Polymethacrylamidopropyltrimonium chloride
Behenamine oxide	Hydrolyzed wheat protein/PVP copolymer	Polyoxyethylene dihydroxypropyl linoleaminium chloride
Behenyl PG-trimonium chloride	Hydrolyzed wheat protein polysiloxane polymer	Polyquaternium-2, -5, -6, -11, -16
Behenyl betaine	Hydroxycetyl hydroxyethyl dimonium chloride	Polyquaternium-17, -18, -24, -29, -44
Benzyltrimonium hydrolyzed collagen	Hydroxyproline	Potassium dimethicone copolyol panthenyl phosphate
Canolamidopropyl betaine	Hydroxypropyl chitosan	Potassium lauroyl collagen amino acids
Capramide DEA	Hydroxypropyl guar hydroxypropyltrimonium chloride	Potassium lauroyl hydrolyzed soy protein
Caprylic/capric/lauric triglyceride	Hydroxypropyl-bis-isostearylamidopropyltrimonium chloride	Potassium lauroyl wheat amino acids
Caprylyl pyrrolidone	Hydroxypropyl bis-stearyldimonium chloride	Potassium lauroyl hydrolyzed collagen
Cassia auriculata extract	Hydroxypropyltrimonium gelatin	PPG-3 lanolin alcohol ether
Cetamine oxide	Hydroxypropyltrimonium hydrolyzed keratin	PPG-9 diethyltrimonium chloride
Cetearalkonium chloride	Il.h. silk	PPG-20 lanolin alcohol ether
Chitosan PCA	Hydroxypropyltrimonium hydrolyzed wheat protein	Proline
Citric acid	Isopropyl hydroxybutylamide dimethicone copolyol	Propylene glycol stearate
Cocamidopropyl dimethylamine, C.D. lactate, C.D. propionate	Isopropyl lanolate	PVP/dimethiconylacrylate/polycarbonyl/polyglycol ester
Cocamidopropyl dimethylaminohydroxypropyl hydrolyzed collagen	Isostearamidopropyl betaine, I. dimethylamine	PVP/dimethylaminoethylmethacrylate copolymer
Cocamidopropyl dimonium	Isostearamidopropyl dimethylamine gluconate	PVP/dimethylaminoethylmethacrylate/polycarbonyl/polyglycol ester
Cocamidopropyl dimethylamine gluconate	Isostearamidopropyl dimethylamine glycolate	PVP/hydrolyzed wheat protein copolymer
Cocamidopropyl PG-dimonium chloride, C.P.C. phosphate	Isostearamidopropyl dimethylamine lactate	Quaternium-22, -26, -33, -61, -62, -70, -80
Coco-morpholine oxide	Isostearamidopropyl ethyldimonium ethosulfate	Quaternium-76 hydrolyzed collagen
Coco/oleamidopropyl betaine	Isostearamidopropyl laurylaceto dimonium chloride	Rapeseedamidopropyl benzyltrimonium chloride
Cocodimonium hydroxypropyl hydrolyzed hair keratin	Isostearamidopropyl morpholine, I.M. lactate	Rapeseedamidopropyl epoxypropyl dimonium chloride
Cocodimonium hydroxypropyl hydrolyzed rice protein	Isostearamidopropyl morpholine oxide	Rapeseedamidopropyl ethyldimonium ethosulfate
Cocodimonium hydroxypropyl hydrolyzed silk	Isostearamidopropyl PG-dimonium chloride	Rice peptide
Cocodimonium hydroxypropyl hydrolyzed soy protein	Isostearaminopropylaluminum chloride	Ricinoleamidopropyl dimonium ethosulfate
Cocoon alcohol	Isostearyl hydrolyzed animal protein	Ricinoleamidopropyl betaine
N-Cocoyl-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate	Isostearylamidopropyl dihydroxypropyl dimonium chloride	Ricinoleamidopropyl dimethylamine lactate
Collagen phthalate	Lactoglobulin	Ricinoleamidopropyl ethyldimonium ethosulfate
Dibehenyl/diarachidyl dimonium chloride	Lauramidopropyl dimethylamine	Ricinoleamidopropyltrimonium chloride
Dibehenyldimonium chloride	Lauramidopropyl PG-dimonium chloride, I.P.C. phosphate	Ricinoleamidopropyltrimonium ethosulfate
Dicetyldimonium chloride	Lauramine oxide	Silicone quaternium-3, -4
Didecyldimonium chloride	Laurampho PG-glycinate phosphate	Silk amino acids
Dihydroxyethyl cocamine oxide	Lauroyl hydrolyzed collagen, L.h. elastin	Sodium/TEA-lauroyl collagen amino acids
Dihydroxyethyl dihydroxypropyl stearamonium chloride	Lauroyl silk amino acids	Sodium/TEA-lauroyl hydrolyzed keratin
Dihydroxyethyl dihydroxypropyl stearamonium chloride	Lauryl methyl gluceth-10 hydroxypropyl-dimonium chloride	Sodium/TEA-lauroyl keratin amino acids
Dihydroxyethyl tallow glycinate	Lauryl phosphate, L. pyrrolidone	Sodium citrate
Dihydroxyethyl tallowamine oxide	Laurylidimonium hydroxypropyl hydrolyzed collagen, keratin, soy protein	Sodium cocoyl hydrolyzed soy protein
Dilauryl acetyl dimonium chloride	Linoleamidopropyl dimethylamine	Sodium hydrogenated tallow dimethyl glycinate
Dilinoamidopropyl dimethylamine	Milk amino acids	Sodium lauroyl collagen, keratin amino acids
Dimethyl hydrogenated tallowamine	Milk protein (Lactis proteinum)	Sodium lauroyl wheat amino acids
Dimethyl lauramine, D.I. isostearate	Myristalkonium chloride	Sodium stearamphoacetate
Dimethyl myristamine, soyamine, stearamine	Myristamidopropyl betaine, M. dimethylamine	Soluble keratin, wheat protein
Dimethylamidopropylamine dimerate	Myristonium bromide	Soyamide DEA
Disodium hydrogenated cottonseed glyceride sulfosuccinate	Oat (Avena sativa) protein	Soyamidopropyl benzyltrimonium chloride
Disodium laureth sulfosuccinate	Oleamide	Soyamidopropyl betaine, S. dimethylamine
Disodium lauroamphodiacetate	Oleamidopropyl betaine, O. dimethylamine	Soyamidopropyl ethyldimonium ethosulfate
Distearyldimonium chloride	Oleamidopropyl dimethylamine hydrolyzed collagen	Soyethyl morpholinium ethosulfate
Ethyl ester of hydrolyzed keratin	Oleamidopropylamine oxide	Soyethyltrimonium ethosulfate
N-Ethylether-bis-1,4-(N-isostearylamidopropyl)-N,N-dimethyl ammonium chlo	Oleamine	Stearamide MEA
Glutamic acid	Oleamine oxide	Stearamidocetyl diethylamine, ethanolamine
Glycerol collagenate	Oleoyl sarcosine	Stearamidopropyl benzyl dimonium chloride
Glycine	Oleyl betaine	Stearamidopropyl cetyl dimonium tosylate
Guar hydroxypropyltrimonium chloride	Oleyl dimethylamidopropyl ethonium ethosulfate	Stearamidopropyl dimethylamine stearate
Henna (Lawsonia inermis) extract	Palmitamidopropyl betaine	Stearamidopropyl ethyldimonium ethosulfate
Hydrogenated tallowamine oxide	Palmitamidopropyl dimethylamine	Stearamidopropyl morpholine lactate
Hydrogenated tallowtrimonium chloride	Palmitamine, P. oxide	Stearamidopropyl PG-dimonium chloride
Hydrolyzed concanin protein	Panthenyl hydroxypropyl steardimonium chloride	phosphate
Hydrolyzed egg protein	PEG-2 milk solids	Stearamine oxide
Hydrolyzed extensin	PEG-3 lauramine oxide	Steardimonium hydroxypropyl hydrolyzed collagen, keratin
Hydrolyzed fibronectin	PEG-5 stearyl ammonium lactate	Steardimonium panthenol
Hydrolyzed fish protein	PEG-15 cocomonium chloride	Stearyl amidoethyl diethylamine
Hydrolyzed keratin	PEG-15 cocopolamine	Steartimonium bromide
Hydrolyzed lactalbumin	PEG-15 tallowmonium chloride	Stearyl dimethicone
Hydrolyzed milk protein	PEG-27	Tallowamidopropyl dimethylamine
Hydrolyzed oats	PEG-40	Tetramethyl trihydroxy hexadecane
Hydrolyzed reuculin	PEG-85 lanolin	TEA-cocoyl hydrolyzed collagen
Hydrolyzed soy protein	PEG-7000	Trachea hydrolysate
	Polydimethicone copolyol	Tricetylmonium chloride
		Tridecyl salicylate
		Triethonium hydrolyzed collagen ethosulfate
		Wheat germamidopropylaluminum chloride
		Wheat germamidopropyl dimethylamine lactate

Functions

Wheat germamidopropyl ethyldimonium
ethosulfate
Wheat peptide
Yeast powder, deproteinized

Coupling agent

Acetyl monoethanolamine
Butyloctan-1-ol
Myreth-J
Oleyl alcohol
PPG-10 butanediol
PPG-10 cetyl ether
PPG-10 oleyl ether
PPG-15 stearyl ether
PPG-22 butyl ether
PPG-23 oleyl ether
PPG-50 oleyl ether
Trideceth-7 carboxylic acid

Denaturant

Brucine sulfate
Denatonium benzoate, saccharide
Nicotine sulfate
Sucrose octaacetate
Thymol

Dental powder

Dicalcium phosphate
Silica
Sodium monofluorophosphate
Stannous fluoride

Deodorant

Abietic acid
Azadirachta indica extract
Chlorophyllin-copper complex
Eugenia jambolana extract
Farnesol
Fermented vegetable
Mauntia flexosa extract
Salvia miltiorrhiza extract
Sodium aluminum chlorohydroxy lactate
Spondias amara extract
Triethyl citrate
Zinc phenol sulfonate, Z. ncinoleate

Depilatory

Barium sulfide
Beeswax, oxidized
Calcium thioglycolate
L-cysteine HCL
Potassium thioglycolate
Sodium thioglycolate
Thioglycerin

Detergent

Ammonium laureth sulfate
Ammonium lauryl sulfate
Capramide DEA
Cocamidopropyl dimethylamine lactate
Decyl glucoside
Decyltetradeceth-25
DEA lauryl sulfate
Diamyl sodium sulfosuccinate
Dicyclohexyl sodium sulfosuccinate
Diisobutyl sodium sulfosuccinate
Disodium caproamphodipropionate
Disodium caproamphodipropionate
Disodium capryloamphodipropionate
Disodium capryloamphodipropionate
Disodium cetaryl sulfosuccinate
Disodium cocamid MEA-sulfosuccinate
Disodium cocamid MIPA-sulfosuccinate
Disodium cocoamphodipropionate
Disodium deceth-6 sulfosuccinate
Disodium isodecyl sulfosuccinate
Disodium lauramido MEA-sulfosuccinate
Disodium lauramido PEG-2 sulfosuccinate
Disodium laureth sulfosuccinate

Disodium lauroamphodipropionate
Disodium lauroamphodipropionate
Disodium lauryl sulfosuccinate
Disodium myristamido MEA-sulfosuccinate
Disodium nonoxynol-10 sulfosuccinate
Disodium oleamid PEG-2 sulfosuccinate
Disodium PEG-4 cocamid MIPA-sulfosuccinate
Disodium ricinoleamid MEA-sulfosuccinate
Disodium tallowiminodipropionate
Dodecylbenzene sulfonic acid
Dodecylbenzene sulfonic acid
Dodecylbenzene sulfonic acid
Isopropylamine dodecylbenzenesulfonate
Isostearamidopropyl betaine
Isosteareth-6 carboxylic acid
Isostearamphodipropionate
Isostearyl hydroxyethyl imidazoline
Lauramidopropylamine oxide
Laureth-11
Lauroampho PG-glycinate phosphate
Lauryl glucoside, L. phosphate
Magnesium laureth sulfate, M. lauryl sulfate
Magnesium PEG-3 cocamide sulfate
MEA-dodecylbenzenesulfonate
MEA-laureth sulfate
MEA-lauryl sulfate
MIPA-lauryl sulfate
Myristamine oxide
Myristic acid
Nonoxynol-10
Oleoamphohydroxypropylsulfonate
Oleth-12, -15
Oleyl betaine
Palmitamidopropyl betaine
PEG-10 glyceryl stearate
PEG-15 glyceryl stearate
PEG-25 glyceryl isostearate
Potassium cocoyl hydrolyzed collagen
Sodium caproamphoacetate
Sodium cocoamphoacetate
Sodium cocoamphodipropionate
Sodium cocomonoglyceride sulfate
Sodium cocoyl hydrolyzed soy protein
Sodium cocoyl isethionate
Sodium C12-15 pareth-25 sulfate
Sodium C14-16 olefin sulfonate
Sodium C14-17 alkyl seculfonate
Sodium deceth sulfate
Sodium decyl diphenyl ether sulfonate
Sodium dodecylbenzenesulfonate
Sodium dodecylidiphenyl ether sulfonate
Sodium iodate
Sodium laureth-2 sulfate
Sodium laureth-3 sulfate
Sodium laureth-7 sulfate
Sodium laureth-12 sulfate
Sodium laureth-13-carboxylate
Sodium laureth sulfate
Sodium lauriminodipropionate
Sodium lauroamphodipropionate
Sodium lauryl methyl alaninate
Sodium lauryl phosphate, S.L. sulfate
Sodium lauryl sulfoacetate
Sodium methyl oleoyl taurate
Sodium methyl cocoyl taurate
Sodium methyl lauroyl taurate
Sodium methyl naphthalenesulfonate
Sodium myreth sulfate
Sodium myristyl sulfate
Sodium oetyl sulfate, oleyl sulfate
Sodium POE alkyl ether acetate
Sodium trideceth-7 carboxylate
Sodium trideceth sulfate
Sodium tridecyl sulfate
Stearth-11, -30
TEA-dodecylbenzenesulfonate
TEA-laureth sulfate
TEA-lauryl sulfate
TEA-palm kernel sarcosinate

TEA-PEG-3 cocamide sulfate
Undecylenamidopropyl betaine

Disinfectant

Benzalkonium chloride
Chlorophene
Didecylmonium chloride
Myristalkonium saccharinate
Shikonin
Sodium capryloamphoacetate
Tea tree (Melaleuca alternifolia) oil
p-Tertiaryphenol

Dispersant

Alkylated polyvinylpyrrolidone
C20-40, C30-50, C40-60 alcohols
Castor (Ricinus communis) oil
Ceteareth-20
Cetyl PPG-2 isodeceth-7 carboxylate
Cholesteryl/beheryl/octyldodecyl lauryl glutamate
Decaglycerol monodiolate
Diisocetyl dodecanedioate
Diisostearyl adipate
Dimethicone copolyol methyl ether
Diocetyldodecyl dimer diisoleate
Diocetyldodecyl dodecanedioate
Ethyl hydroxymethyl oleyl oxazoline
Glyceryl caprylate, G. caprylate/caprate
Glyceryl diisostearate
Hydrogenated castor oil, H. lecithin
Hydrogenated tallow glycerides
Isobutylene/MA copolymer
Isocetyl alcohol
Isopropyl C12-15-pareth-9-carboxylate
Isostearyl neopenanoate
Lanolin acid
Laureth-4, -6, -16
Melania
Nonoxynol-2, -18, -20, -30, -40
Octoxynol-5, -10
Octoxynol 16, 30, 40, 70
Octyldodeceth-5
Octyldodecyl/dimethicone copolyol citrate
Oleth-40
Oleyl alcohol
PEG-5 castor oil, glyceryl sesquiolate
PEG-6 beeswax
PEG-8/SMDI copolymer
PEG-9 castor oil, oleate, stearate
PEG-10 dioleate, stearamine
PEG-12 beeswax
PEG-12 glyceryl dioleate, laurate
PEG-15 castor oil
PEG-20 almond glycerides
PEG-20 glyceryl isostearate
PEG-20 sorbitan trisostearate
PEG-25 castor oil
PEG-30 dipolyhydroxystearate
PEG-40 hydrogenated castor oil PCA isostearate
PEG-60 shea butter glycerides
Poloxamer 101, 122, 181, 182, 184
Polyglyceryl-2 sesquiosostearate
Polyglyceryl-3 diisostearate, oleate
Polyglyceryl-5 diisostearate
Polyglyceryl-6 mixed fatty acids
Polyglyceryl-10 diisostearate, distearate
Polyglyceryl-10 decaoleate
Polyhydroxystearic acid
Polysorbate 40, 80
Potassium polyacrylate
PPG-3 PEG-6 oleyl ether
PPG-9 diethylmonium phosphate
PPG-12/SMDI Copolymer
PPG-15 stearyl ether
PPG-25, PPG-40 diethylmonium chloride
PPG-51/SMDI Copolymer
PVP/icosene copolymer
PVP/hexadecene copolymer

Functions

Rapeseed oil, ethoxylated high erucic acid
Ricinolevl alcohol
Sodium ceteth-1 1-carboxylate
Sodium laurylsulfonate, S. polymethacrylate
Sodium polyphthalenesulfonate
Sorbitan oleate
Steareth-10
Tricortanvl PVP
Trisostearin PEG-6 esters
Truxtyldodecyl citrate

Emollient

Acetylated elvent stearate
Acetylated hydrogenated lanolin
Acetylated hydrogenated lard glycende
Acetylated hydrogenated vegetable glycende
Acetylated lanolin, A.I. alcohol
Acetylated lard glycende
Acetylated monomylcendes
Acetylated palm kernel glycendes
Aleurites molluccana ethyl ester
Allantoin
Aluminum/magnesium hydroxide stearate
AMP-isoctearyl hydrolyzed soy protein
Apricot (Prunus armeniaca) kernel oil
Arachidyl behenate
Argania spinosa oil
Avocado (Persea gratissima) oil, unsaponifiables
Avocado oil ethyl ester
Babassu (Orbignya oleifera) oil
Butyl isostearate, B. stearate
Behenamidoethyl dihydroxypropyl dimonium chloride
Behenoxy dimethicone
Behenyl alcohol, B. behenate
Behenyl erucate, B. isostearate
Benzyl laurate
Bladderwrack (Fucus vesiculosus) extract
Borage (Borago officinalis) seed oil
Borageamidoethyl phosphatidyl PG-dimonium chloride
Brain extract
Brazil nut (Benthoetia excelsa) oil
Butyl myristate, oleate, stearate
Butyloctanol
Butyloctyl oleate
C12-13, C12-16, C14-15 alcohols
C12-15 alcohols octanoate
C12-15 alkyl benzoate
dl-C12-15 alkyl fumarate
C12-15 alkyl lactate
Camellia kissi oil
Tea (Camellia sinensis) oil
C10-30 cholesterol/lanosterol esters
Canola oil
Caprylic/capric triglycende
Caprylic/capric triglycende PEG-4 esters
Caprylic/capric/laure triglycende
Caprylic/capric/linoleic triglycende
Caprylic/capric/oleic triglycendes
Caprylic/capric/steane triglycende
Caprylic/capric/succinic triglycende
Capsicum frutescens oleoresin
Carrot (Daucus carota sativa) oil
Cashew (Anacardium occidentale) nut oil
Castor (Ricinus communis) oil
Cetearyl behenate, C. candelillate
Cetearyl isononanoate, C. octanoate
Cetearyl palmitate, C. stearate
Ceteth-10
Cetostearyl stearate
Cetyl C12-15 pareth-9 carboxylate
Cetyl acetate, C. alcohol
Cetyl esters, C. lactate
Cetyl myristate, C. octanoate
Cetyl oleate, C. palmitate
Cetyl PPG-2 isodeceth-7 carboxylate
Cetyl ricinoleate, C. stearate

Cetyl stearyl octanoate
Chia (Salvia hispanica) oil
Cholesteric esters
Cholesterol
Cholesteryl/behenyl/oxtyldodecyl lauroyl glutamate
Cholesteryl hydroxystearate
Cholesteryl stearate
Choleth-24
C 18-70 Isoparaffin
C10-18, C12-18 triglycendes
C12-15 linear alcohols 2-ethylhexanoate
Cicumidopropyl PG-dimonium chloride
Cocoa (Theobroma cacao) butter
Coco-caprylate/caprate
Coco-rapeseedate
Coconut (Cocos nucifera) oil
Cocoyl hydrolyzed soy protein
Collagen phthalate
Colloidal oatmeal
Comirex (Symphytum officinale) leaf extract
Corn (Zea mays) oil
Corn poppy (Papaver rhoeas) extract
Cottonseed (Gossypium) oil
Cuttlefish extract
Cyclomethicone
Deceth-4 phosphate
Decyl oleate
Decyltetradecanol
Dialkyldimethylpolysiloxane
Dibutyl sebacate
Dicapryl adipate
Dicaprylyl ether, D. malate
Diethylene glycol diisononanoate
Diethylene glycol dioctanoate
bis-Diglyceryl/caprylate/caprate/isostearate/
hydroxystearate/adipate
bis-Diglyceryl/caprylate/caprate/isostearate/
stearate/hydroxystearate/adipate

Dihydroabietyl behenate
Dihydroxyethyl tallowamine oleate
Diisobutyl adipate
Diisocetyl adipate, dodecanedioate
Disodecyl adipate
Diisopropyl adipate, dimer dilinoleate
Diisopropyl sebacate
Diisostearyl trimethylolpropane siloxy silicate
Diisostearyl adipate
Diisostearyl dimer dilinoleate
Diisostearyl fumarate, D. malate
Dilinoleic acid
Dimethicone
Dimethicone copolyol
Dimethicone copolyol acetate, D.C. almondate
Dimethicone copolyol isostearate, D.C. lactate
Dimethicone copolyol methyl ether
Dimethicone copolyol phthalate
Dimethicone propylethylendiamine behenate
Dimethiconol stearate
Dimethyl lauramine oleate
Dioctyl adipate
Dioctyl dimer dilinoleate
Dioctylcyclohexane
Dioctyldodecyl dimer dilinoleate
Dioctyldodecyl dodecanedioate
Dioctyl malate, D. sebacate, succinate
Dipentaerythritol fatty acid ester
Dipentaerythritol hexacaprylate/hexacaprate
Dipentaerythritol hexahydroxystearate/isostearate
Distearyl dimethylamine dilinoleate
Ditridecyl adipate
Dog rose (Rosa canina) hips oil
Egg (Ovum) yolk extract
Emu (Dromiceus) oil
Erucyl erucate
Ethyl avocado
Ethylhexyl isopalmitate

COSMETIC AND PHARMACEUTICAL INGREDIENTS

CAMPHOR USP
CARBOXYMETHYLCELLULOSE USP
CETINA (CETYL ESTERS & STEARAMIDE DEA)
SPERMWAX® (CETYL ESTERS WAX)
CHOLESTEROL NF
DENATONIUM BENZOATE NF
GLYCINE USP
IPG (ISOPENTYLDIOL)
MENTHOL USP
ROBANE (SQUALANE NF)
SUPRAENE® (SQUALENE)
UREA PEROXIDE USP

ROBEKO INC.
99 PARK AVENUE • NEW YORK, NY 10016
212-986-6410
FAX, 212-986-6419

OUR 78TH YEAR



Functions

2-Ethylhexyl isostearate	Isononyl isononanoate	Octyldodecanoil
Ethyl linolenate, E. myristate	Isopentylidol	Octyldodecyl behenate, O. benzoate
Ethyl myristate, E. myristate	Isopropyl avocadate	Octyldodecyl erucate, O. myristate
Ethyl oleate, E. olive	Isopropyl C12-15-pareth-9-carboxylate	Octyldodecyl oleate, O. ricinoleate
Evening primrose (<i>Oenothera biennis</i>) extract, oil	Isopropyl isostearate	Octyldodecyl stearate
Glycereth-4.5-lactate	Isopropyl lanolate, I. linoleate	bis-Octyldodecyl stearoyl dimer dilinoleate
Glycereth-5 lactate	Isopropyl myristate, I. palmitate	Octyldodecyl stearoyl stearate
Glycereth-7 benzoate	Isopropyl PPG-2-isodeceth-1 carboxylate	Oleamine oxide
Glycereth-7 diisononanoate	Isopropyl stearate	Oleic/palmitoleic/linoleic glycerides
Glycereth-7 triacetate	Isoribide laurate	Oleic alcohol
Glycereth-7 trioctanoate	Isostearic acid	Oleostearine
Glycereth-12, -26	Isostearyl alcohol	Oleyl alcohol, O. erucate, O. oleate
Glycerol tricaprylate/caprate	Isostearyl behenate, I. benzoate	Olive (<i>Olea europae</i>) oil
Glyceryl adipate, G. dioleate	Isostearyl diglycerol succinate	Orange (<i>Citrus aurantium dulcis</i>) peel wax
Glyceryl isostearate, G. lanolate	Isostearyl erucate, I. erucyl erucate	Orange roughy (<i>Hoplostethus atlanticus</i>) oil
Glyceryl linoleate, G. monopyrrolutamate	Isostearyl isostearate, I. lactate	Palm (<i>Elaeis guineensis</i>) oil
Glyceryl myristate, G. oleate	Isostearyl malate, I. myristate	Palm kernel glycerides
Glyceryl ricinoleate	Isostearyl neopentanoate, palmitate	Palmitic acid
Glyceryl triacetyl hydroxystearate	Isostearyl stearoyl stearate	Panthenyl triacetate
Glyceryl triacetyl ricinoleate	Isostearylmidopropyl dihydroxypropyl dimonium chloride	Partially hydrogenated canola oil
Glycosaminoglycans	Isotridecyl isononanoate	Partially hydrogenated soybean oil
Glycosphingolipids	Isotridecyl myristate	Peach (<i>Prunus persica</i>) extract
Gold of Pleasure oil	Jojoba (<i>Buxus chinensis</i>) oil	Peanut (<i>Arachis hypogaea</i>) oil
Grape (<i>Vitis vinifera</i>) seed oil	Jojoba butter, J. esters	Pecan (<i>Carya illinoensis</i>) oil
Hazel (<i>Corylus avellana</i>) nut oil	Jojoba oil, synthetic	PEG-2 diisononanoate, P. dioctanoate
Helianthus annuus ethyl ester	Kukui (<i>Aleurites moluccana</i>) nut oil	PEG-2 milk solids
Hexadecyl isopalmitate	Lactamide DGA	PEG-4
Hexamethyldisiloxane	Laureth-10 acetate	PEG-4 diheptanoate, P. dilaurate
Hexyl laurate	Laolin, L. acid	PEG-5 C8-12 alcohols citrate
Hexyldecanol	Laolin alcohol, L. oil	PEG-5 C14-18 alcohols citrate
Hexyldecyl stearate	Laolin, ultra anhydrous	PEG-5 hydrogenated castor oil
Honey extract	Laolin wax	PEG-5 hydrogenated castor oil triisostearate
Hybrid safflower (<i>Carthamus tinctorius</i>) oil	Lanosterol	PEG-6
Hybrid sunflower (<i>Helianthus annuus</i>) oil	Lard glyceride	PEG-6 capric/caprylic glycerides
Hydrogenated C6-14 olefin polymers	Laureth-2, -3	PEG-7 glyceryl cocoate
Hydrogenated castor oil	Laureth-2 acetate, L. benzoate	PEG-8
Hydrogenated castor oil laurate	Laureth-2-octanoate	PEG-8 dilaurate, P. dioleate
Hydrogenated coconut oil	Laurel/palmitic/oleic triglyceride	PEG-8/SMD1 copolymer
Hydrogenated cottonseed oil	Lauryl behenate, L. lactate	PEG-9 stearyl stearate
Hydrogenated C12-18 triglycerides	Lauryl phosphate	PEG-10 stearyl stearate
Hydrogenated lanolin	Lauryldimethylamine isostearate	PEG-12
Hydrogenated lanolin, distilled	Lesquerella fendleri oil	PEG-12 dioleate, P. palm kernel glycerides
Hydrogenated lecithin	Linoleic acid	PEG-15 cocamine oleate/phosphate
Hydrogenated milk lipids	Macadamia ternstroemia nut oil	PEG-18
Hydrogenated mink oil	Maleated soybean oil	PEG-20
Hydrogenated palm kernel glycerides	Mango (<i>Mangifera indica</i>) oil, seed oil	PEG-20 hydrogenated castor oil isostearate
Hydrogenated palm oil	Mango kernel oil	PEG-20 hydrogenated castor oil triisostearate
Hydrogenated polyisobutene	Meadowfoam (<i>Limnanthes alba</i>) seed oil	PEG-20 hydrogenated lanolin
Hydrogenated soybean oil	Menhaden (<i>Brevoortia tyrannus</i>) oil	PEG-24 hydrogenated lanolin
Hydrogenated starch hydrolysate	Methyl acetyl ricinoleate	PEG-25 PABA, P. propylene glycol stearate
Hydrogenated tallow glycende	Methyl gluceth-20	PEG-40 glyceryl laurate
Hydrogenated tallow glycende lactate	Methyl gluceth-20 benzoate, M. g. distearate	PEG-40 hydrogenated castor oil isostearate
Hydrogenated turtle oil	Methyl hydroxystearate, M. ricinoleate	PEG-40 hydrogenated castor oil laurate
Hydrogenated vegetable glycerides	Microcrystalline wax	PEG-40 hydrogenated castor oil triisostearate
Hydrogenated vegetable oil	Mineral oil (<i>Paraffinum liquidum</i>)	PEG-40 jojoba oil
Hydrolyzed collagen	Mink oil	PEG-50 hydrogenated castor oil laurate
Hydrolyzed conchiorin protein	Musk rose (<i>Rosa moschata</i>) oil	PEG-50 hydrogenated castor oil triisostearate
Hydrolyzed keratin	Myreth-3	PEG-60 shea butter glycerides
Hydrolyzed mushroom (<i>Tricholoma matsutake</i>) extract	Myreth-3 caprate, M. laurate	PEG-70 mango glycerides
Hydrolyzed oat protein	Myreth-3 myristate, M. octanoate	PEG-75
Hydroxylated lanolin	Myristyl alcohol, M. lactate	PEG-75 lanolin, P. shea butter glycerides
Hydroxylated milk glycerides	Myristyl myristate, M. octanoate	PEG-75 shea butter glycerides
Hydroxystearic acid	Myristyl propionate, M. stearate	PEG-150
Ilupe butter	Neatsfoot oil	PEG/PPG-17/6 copolymer
Isobutyl palmitate, I. stearate	Neem (<i>Melia azadirachta</i>) seed oil	Pentaerythrityl dioleate
Isocetyl behenate, I. octanoate	Neopentyl glycol dicaprate	Pentaerythrityl isostearate/caprate/caprylate/adipate
Isocetyl palmitate, I. salicylate	Neopentyl glycol dicaprate/dicaprylate	Pentaerythrityl stearate
Isocetyl stearate	Neopentyl glycol diisooctanoate	Pentaerythrityl stearate/caprate/caprylate/adipate
Isodeceth-2 cocoate	Neopentyl glycol dioctanoate	Pentaerythrityl tetracaprylate/tetracaprate
Isodecyl citrate, I. cocoate	Oat (<i>Avena sativa</i>) bran extract, extract, flour	Pentaerythrityl tetraisononanoate, P. tetraisostearate
Isodecyl isononanoate, I. laurate	Octacosanyl stearate	Pentaerythrityl tetralaureate, P. tetraoctanoate
Isodecyl neopentanoate	Octyl cocoate	Pentaerythrityl tetraoleate, P. tetrapelargonate
Isodecyl octanoate, I. oleate	Octyl hydroxystearate, O. isononanoate	Pentaerythrityl tetrastearate
Isodecyl stearate	Octyl neopentanoate, O. octanoate	Perfluorodecalin
Isododecane	Octyl oleate, O. palmitate	Perfluoropolydimethylisopropyl ether
Isocicosane	Octyl pelargonate, O. stearate	Petrolatum
Isohexadecane	Octyldecanol	Phenethyl dimethicone
		Phenyl dimethicone, P. methicone, P. trimethicone

Functions

Phytantriol	PPG-8/SMDI copolymer	Propylene glycol myristyl ether acetate
Pistachio (Pistacia vera) nut oil	PPG-9	Propylene glycol stearate, SE
Placental enzymes	PPG-9-buteth-12	Pumpkin (Cucurbita pepo) seed oil
Pollen extract	PPG-9-butyl ether	Quinoa (Chenopodium quinoa) oil
Poloxamer 105 benzoate	PPG-10 butanediol, P. cetyl ether	Rapeseed (Brassica campestris) oil
Poloxamer 182 dibenzoate	PPG-10 methyl glucose ether	Rice (Oryza sativa) bran oil, bran wax
Polybutene	PPG-10 oleyl ether	Rice fatty acid
Polydecene	PPG-11 stearyl ether	Safflower (Carthamus tinctorius) oil
Polydimethicone copolyol	PPG-12 buteth-16	Salmon (Salmo) egg extract
Polyethylene glycol	PPG-12 PEG-50 lanolin	Sesame (Sesamum indicum) oil
Polyglyceryl-2 diisostearate, P. tetrastearate	PPG-12-PEG-65 lanolin oil	Shark liver oil
Polyglyceryl-2 trisostearate	PPG-12/SMDI Copolymer	Shea butter (Butyrospermum parkii)
Polyglyceryl-3 diisostearate, P. oleate	PPG-14 butyl ether	Shea butter (Butyrospermum parkii) extract
Polyglyceryl-3 stearate	PPG-15 butyl ether, P. stearyl ether	Shea butter, ethoxylated
Polyglyceryl-6 dioleate	PPG-15 stearyl ether benzoate	Shorea stenoptera butter
Polyglyceryl-10 decanoate, P. decastearate	PPG-16 butyl ether	Silybum marianum ethyl ester
Polyglyceryl-10 tetraoleate	PPG-18 butyl ether	Sitostearyl acetate
Polyisobutene	PPG-20	Skin lipids
Polyisobutene:isohexapentacontahexane	PPG-20-buteth-30	Slippery elm extract
Polyisobutene:isooctahexacontane	PPG-20 cetyl ether	Sodium C8-16 isoalkylsuccinyl lactoglobulin sulfonate
Polyisobutene:isopentacontaoctane	PPG-24-glycereth-24	Sodium carboxymethyl beta-glucan
Polyisoprene	PPG-26	Sodium ceteth-13-carboxylate
Polyoxyethylene polyoxypropylene glycol	PPG-27 glyceryl ether	Sodium dimethicone copolyol acetyl methylaurate
Polyquaternium-2	PPG-28-buteth-35	Sodium glyceryl oleate phosphate
Polysiloxane polyalkylene copolymer	PPG-30	Sodium hyaluronate, S. polymethacrylate
Poly sorbate 40	PPG-30 cetyl ether	Sorbeth-20
Potassium dimethicone copolyol phosphate	PPG-40 butyl ether	Sorbitan isostearate, S. palmitate
PPG-2-buteth-3	PPG-50 cetyl ether, P. oleyl ether	Sorbitan sesquioleate, S. sesquisteate
PPG-2 lanolin alcohol ether	PPG-51/SMDI Copolymer	Sorbitan trioleate
PPG-2 myristyl ether propionate	PPG-53 butyl ether	Soybean (Glycine soja) oil
PPG-3 hydrogenated castor oil	Propylene glycol ceteth-3 acetate	Spermaceil
PPG-3 myristyl ether	Propylene glycol dicaprylate	Sphingolipids
PPG-5-buteth-7	Propylene glycol dipalmylate:dicaprate	Squalene
PPG-5-laureth-5	Propylene glycol diisostearate, P.g. dioctanoate	Stearamidopropyl cetecaryl dimonium tosylate
PPG-5-butyl ether	Propylene glycol dipelargonate	Stearth-4 stearate
PPG-5 lanolin wax	Propylene glycol isoceteth-3 acetate	Stearic acid, S. hydrazide
PPG-5 pentaerythrityl ether	Propylene glycol isostearate, P.g. laurate	Stearoxy dimethicone
PPG-7-buteth-10	Propylene glycol myristate	

ANIMAL OR VEGETABLE?

New V-Series Cerasynt® emulsifiers give you the choice

ISP Van Dyk has added vegetable-based Cerasynt® derivatives to their outstanding emulsifier line. Cerasynt SD-V and IP-V provide the same excellent performance as the original animal-derived products. They are ideal for use as secondary emulsifiers, stabilizers and opacifiers in a wide variety of cosmetic creams and lotions. For information, call **201-450-7724**.

ISP

VAN DYK

a subsidiary of International Specialty Products

For samples, call the ISP Sample Center at 1-800-243-6788. To place an order, call ISP Customer & Sales Service at 1-800-622-4423.

Functions

Stearoxymethiconedimethicone copolymer
 Stearyl behenate, S. benzoate
 Stearyl dimethicone, S. erucate
 Stearyl heptanoate, S. propionate
 Stearyl stearate
 Stearyl stearoyl stearate
 Sucrose cocate
 Sunflower (Helianthus annuus) seed oil
 Sweet almond (Prunus amygdalus dulcis) oil
 Sweet cherry (Prunus avium) pit oil
 Synthetic jojoba oil
 Synthetic wax
 Tallow
 Tetradecylcyclostearyl stearate
 Tocopheryl acetate
 Tricaprin
 Tricaprylin
 Tricaprylyl citrate
 Tricholoma matsutake extract
 Tridecyl behenate, T. cocate
 Tridecyl erucate, T. neopentanoate
 Tridecyl octanoate, T. stearate
 Tridecyl stearoyl stearate
 Tridecyl trimellitate
 Trihexyldecyl citrate
 Trisoceryl citrate
 Trisosteann
 Trisostearyl citrate
 Trisostearyl linoleate
 Trilaunin
 Trilinolein
 Trimethylolpropane triacrylate/tricaprate
 Trimethylolpropane triacetate
 Trimethylolpropane trioleate
 Trimyrstin
 Trioctanoic
 Trioctyldodecyl citrate
 Triolein
 Tripalmitin
 Tripropylene glycol citrate
 Tristearin
 Triundecanoic
 Vegetable oil
 Walnut (Juglans regia) oil
 Wheat (Triticum vulgare) germ oil

Emulsifier
 Acetylated hydrogenated lard glyceride
 Acetylated hydrogenated vegetable glyceride
 Acetylated monoglycerides
 Acrylates/C10-C30 alkyl acrylate crosspolymer
 Acrylates/vinyl isodecanoate crosspolymer
 Acrylic acid/acrylonitrile copolymer
 2-Aminobutanol
 Ammonium acrylates/acrylonitrile copolymer
 Arachidyl alcohol
 Beeswax
 Behenamidopropyl dihydroxypropyl dimonium chloride
 Beheneth-5 -10 -20 -30
 Behenic acid
 Behenyl betaine
 Borageamidopropyl phosphatidyl PG-dimonium chloride
 Butyloctanol
 C12-20 acid PEG-8 ester
 C18-36 acid
 Calcium dodecylbenzene sulfonate
 Calcium protein complex

Calcium stearate
 Calcium stearyl lactylate
 Capramide DEA
 Caprylic/capric acid
 Caprylic/capric glycerides
 Castor oil, ethoxylated
 Cetalkonium chloride
 Ceteareth-2 -4 -5 -6
 Ceteareth-2 phosphate
 Ceteareth-3 phosphate
 Ceteareth-8 -10 -11 -12
 Ceteareth-10 phosphate
 Ceteareth-15 -17 -20 -25
 Ceteareth-27 -29 -30 -34
 Cetearyl alcohol
 Cetearyl glucoside
 Ceteth-2 -4 -6 -10 -12 -13
 Ceteth-16 -20 -25 -30 -33
 Cetethyldimonium bromide
 Cetrimonium chloride
 Cetyl dimethicone copolyol
 Cetyl phosphate
 Cholesterol
 Choleth-10 -15 -24
 Cocamide DEA, C. MEA
 Cocamidopropyl dimethylamine
 Cocamidopropyl PG-dimonium chloride phosphate
 Cocamine
 Coceth-7 carboxylic acid
 Coconut acid
 Copper protein complex
 Cottonseed glyceride
 C12-13 pareth-3 -4 -9 -23
 C16-18 pareth-3 -5.5 -13 -19
 Cyclodextrin
 Decaglycerol monodiolate
 DEA-ceteareth-2-phosphate
 DEA-cetyl phosphate
 DEA-cyclocarboxypropylolate
 DEA-oleth-3 phosphate
 DEA-oleth-5-phosphate
 DEA oleth-10 phosphate
 DEA-oleth-20-phosphate
 Dicitateareth-10 phosphonic acid
 Diethanolamine
 Diethylaminooethyl stearate
 Diglyceryl stearate malate
 Dihydrocholeth-15 -20 -30
 Dihydrogenated tallow phthalic acid amide
 Dilauryl acetyl dimonium chloride
 Dilinoleamidopropyl dimethylamine dimethicone copolyol phosphate
 Dilinoleic acid
 Dimethicone copolyol almondate
 Dimethicone copolyol isostearate
 Dimethicone copolyol laurate
 Dimethicone copolyol methyl ether
 Dimethicone copolyol olivate
 Dimethicone copolyol phthalate
 Dipalmitolethyl hydroxyethylmonium methosulfate
 Dipropylene glycol
 Disodium hydrogenated cottonseed glyceride sulfosuccinate
 Disodium ricinoleamide MEA-sulfosuccinate
 Disodium stearyl sulfosuccinate
 Disodium sulfosuccinamide
 Distearyl phthalic acid amide

N-Dodecyl-N,N-dimethyl-N-(dodecyl acetate) ammonium chloride
 Dodecylphenol-ethylene oxide condensate
 Egg (Ovum) yolk extract
 Emulsifying wax NF
 Ethoxylated fatty alcohol
 N-Ethylether-bis-1,4-(N-iso-stearylaminopropyl)-N,N-dimethyl ammonium chlo
 Ethyl hexanediol
 Euglena gracilis polysaccharide
 Glyceth-26 phosphate
 Glyceryl caprylate, G. caprylate/caprate
 Glyceryl citrate/lactate/linoleate/oleate
 Glyceryl cocate, G. dilaurate
 Glyceryl dilaurate, G. dioleate
 Glyceryl distearate, G. hydroxystearate
 Glyceryl isostearate, G. lanolate
 Glyceryl laurate, G. linoleate
 Glyceryl mono-di-tri-caprylate
 Glyceryl myristate, G. oleate
 Glyceryl palmitate, G. ricinoleate
 Glyceryl ricinoleate SE
 Glyceryl stearate, G. stearate citrate
 Glyceryl stearate lactate
 Glyceryl stearate SE
 Glyceryl undecylenate
 Glycol distearate, G. oleate
 Glycol palmitate, G. stearate
 Glycol stearate SE
 Glycolamide stearate
 Glycosphingolipids
 Hydrogenated coco-glycerides
 Hydrogenated cottonseed glyceride
 Hydrogenated lanolin
 Hydrogenated lecithin
 Hydrogenated palm oil
 Hydrogenated soy glyceride
 Hydrogenated tallow glycerides
 Hydrogenated tallow glycerides citrate
 Hydroxycetyl phosphate
 Hydroxylated lanolin
 Hydroxylated lecithin
 Hydroxyoctacosanyl hydroxystearate
 Hydroxypropyl-bis-isostearylamidopropylidimonium chloride
 Isoceteareth-8 stearate
 Isoceteareth-10 stearate
 Isoceteareth-20
 Isocetyl alcohol
 Isolaureth-6
 Isostearamidopropyl dimethylamine gluconate
 Isostearamidopropyl dimethylamine glycolate
 Isostearamidopropyl laurylacetylmonium chloride
 Isosteareth-2 -3 -10 -12 -20 -22 -50
 Isosteareth-2-octanoate
 Isosteareth-10 stearate
 Isostearic acid
 Isostearyl diglyceryl succinate
 Isostearylaminopropyl dihydroxypropyl dimonium chloride
 Karaya (Sterculia urens) gum
 Laneth-5 -10 -15 -16 -20 -40
 Laneth-10 acetate
 Lanolin
 Lanolin alcohol
 Lanolin, ultra anhydrous
 Lanolin wax
 Lauramide DEA, L. MEA

3 BETTER IDEAS.

1 BETTER SOURCE.

CARDOPOL
Ultra
 New, easiest to
 disperse carboxomer

CARDOPOL
ETD 2020
 POLYMER
 For surfactant-based
 products

PEMULEN
 POLYMERIC EMULSIFIERS
 Eliminates surfactant-based
 emulsifiers

BF Goodrich
 Talk to the global leader.

Functions

Laureamidopropyl dimethylamine	PEG-5 lanolate, P. oleamine	PEG-20 lanolin, P. laurate
Laureamidopropyl PG-dimonium chloride	PEG-5 soy sterol, P. soyamine	PEG-20 oleate
Laureth-1 -2 -3 -4 -5	PEG-5 stearamine, P. stearate	PEG-20 methyl glucose sesquistearate
Laureth-2-octanoate	PEG-5 tallow amine	PEG-20 sorbitan beeswax
Laureth-3 phosphate	PEG-6 capric/caprylic glycerides	PEG-20 sorbitan isostearate
Laureth-4 carboxylic acid	PEG-6 cocamide	PEG-20 sorbitan trisostearate
Laureth-5 carboxylic acid	PEG-6 C12-14 ether	PEG-20 sorbitan trioleate
Laureth-6 -7 -9 -11 -12	PEG-6 dilaurate, P. dioleate	PEG-20 stearate, P. tallow amine
Laureth-11 carboxylic acid	PEG-6 distearate, P. isostearate	PEG-23 oleate, P. stearate
Laureth-16 -20 -23 -25 -30	PEG-6 lauramide, P. laurate	PEG-24 hydrogenated lanolin
Lauryl PCA	PEG-6 oleate, P. palmitate	PEG-25 castor oil
Lauryl methicone copolyol	PEG-6 sorbitan beeswax	PEG-25 phytosterol
Lecithin	PEG-6 sorbitan laurate	PEG-25 propylene glycol stearate
Linoleamidopropyl PG-dimonium chloride	PEG-6 sorbitan oleate	PEG-25 soy sterol, P. stearate
phosphate	PEG-6 sorbitan stearate	PEG-29 castor oil
Lithium stearate	PEG-6 stearate	PEG-30 castor oil
Magnesium sulfate hepta-hydrate	PEG-6-32	PEG-30 dipolyhydroxystearate
Maleated soybean oil	PEG-6-32 stearate	PEG-30 glyceryl cocoate
Methoxy PEG-17/dodecyl glycol copolymer	PEG-7 glyceryl cocoate	PEG-30 glyceryl isostearate
Methyl gluceth-20 distearate	PEG-7 hydrogenated castor oil	PEG-30 glyceryl laurate
Methyl glucose dioleate, M. g. sesquistearate	PEG-7 oleate	PEG-30 glyceryl oleate
Methyl glucose sesquistearate	PEG-7.5 tallowamine	PEG-30 glyceryl stearate
MEA-laureth sulfate	PEG-8	PEG-30 hydrogenated castor oil
Myreth-3 -4 -7	PEG-8 beeswax, P. castor oil	PEG-30 lanolin
Myreth-3 myristate	PEG-8 C12-14 ether	PEG-30 sorbitan tetraoleate
Myristamidopropyl dimethylamine	PEG-8 dilaurate, P. dioleate	PEG-32 dilaurate, P. dioleate
Nonoxynol-1 -2 -4 -5 -6 -7	PEG-8 distearate	PEG-32 distearate, P. laurate
Nonoxynol-8 -9 -10 -11 -12 -13	PEG-8 glyceryl laurate	PEG-32 oleate, P. stearate
Nonoxynol-14 -15 -18 -20 -30 -40 -50	PEG-8 laurate, P. oleate	PEG-33 castor oil
Nonyl nonoxynol-5 -10	PEG-8, P. tallate	PEG-35 castor oil, P. stearate
Oat (Avena sativa) flour	PEG-9 castor oil	PEG-40 castor oil
Octoxynol-1 -3 -5 -8 -10	PEG-9 diisostearate	PEG-40 glyceryl isostearate
Octoxynol 16, 30, 40	PEG-9 dioleate, P. distearate	PEG-40 glyceryl laurate
2-Octyl dodecyl alcohol	PEG-9 laurate, P. oleate	PEG-40 glyceryl triisostearate
Octyldodecanol	PEG-9 stearate	PEG-40 hydrogenated castor oil
Octyldodeceth-20 -25	PEG-10 castor oil, P. cocamine	PEG-40 hydrogenated castor oil PCA isostearate
Oleamide DEA	PEG-10 coconut oil esters	PEG-40 sorbitan diisostearate
Oleamidopropyl dimethylamine	PEG-10 C12-18 alcohols	PEG-40 sorbitan lanolate
Oleamine oxide	PEG-10 dioleate	PEG-40 sorbitan tetraoleate
Oleic acid	PEG-10 glyceryl isostearate	PEG-40 stearate
Oleth-2 -3 -4 -5 -6 -7 -8 -9	PEG-10 hydrogenated castor oil	PEG-40/dodecyl glycol copolymer
Oleth-10 -12 -15 -20 -23	PEG-10 hydrogenated castor oil triisostearate	PEG-42 babassu glycerides
Oleth-25 -30 -40 -50	PEG-10 lanolate	PEG-44 sorbitan laurate
Oleth 13	PEG-10 polyglyceryl-2 laurate	PEG-45 palm kernel glycerides
Oleth-2 phosphate	PEG-10 sorbitan laurate	PEG-45 safflower glycerides
Oleth-3 phosphate	PEG-10 soy sterol, P. stearamine	PEG-50 lanolin, P. stearamine
Oleth-5 phosphate	PEG-10 stearate	PEG-50 stearate
Oleth-10 phosphate	PEG-11 babassu glycerides	PEG-60 almond glycerides
Oleth-20 phosphate	PEG-11 castor oil	PEG-60 castor oil
Palm acid	PEG-12 dilaurate, P. dioleate	PEG-60 corn glycerides
Palmitamidopropyl dimethylamine	PEG-12 distearate	PEG-60 glyceryl triisostearate
Palmitic acid	PEG-12 glyceryl dioleate	PEG-60 hydrogenated castor oil
PEG-2 cocamine, P. distearate	PEG-12 laurate, P. oleate	PEG-60 hydrogenated castor oil isostearate
PEG-2 hydrogenated tallow amine	PEG-12 stearate, P. tallate	PEG-60 hydrogenated castor oil triisostearate
PEG-2 laurate, P. laurate SE	PEG-14 avocado glycerides	PEG-60 shea butter glycerides
PEG-2 oleamine, P. oleate	PEG-15 castor oil	PEG-60 sorbitan tetraoleate
PEG-2 soyamine, P. stearamine	PEG-15 cocamine	PEG-70 mango glycerides
PEG-2 stearate, P. stearate SE	PEG-15 glyceryl isostearate	PEG-75
PEG-3 cocamide	PEG-15 glyceryl laurate	PEG-75 castor oil, P. dilaurate
PEG-3 C12-C18 alcohols	PEG-15 glyceryl ricinoleate	PEG-75 dioleate, P. distearate
PEG-3 glyceryl isostearate	PEG-15 oleamine, P. oleate	PEG-75 lanolin, P. laurate
PEG-3 glyceryl triisostearate	PEG-15, P. stearamine	PEG-75 oleate
PEG-3 glyceryl tristearate	PEG-15 tallow amine	PEG-75 shea butter glycerides
PEG-3 lanolate, P. sorbitan oleate	PEG-15 tallow polyamine	PEG-75 shorea butter glycerides
PEG-3 stearate	PEG-16	PEG-75 stearate
PEG-4 dioleate, P. diisostearate	PEG-16 hydrogenated castor oil	PEG-80 sorbitan laurate
PEG-4 dilaurate, P. distearate	PEG-16 soy sterol	PEG-90 stearate
PEG-4 glyceryl distearate	PEG-18 stearate	PEG-100 castor oil
PEG-4 laurate, P. oleate	PEG-20 almond glycerides	PEG-100 hydrogenated castor oil
PEG-4 stearate	PEG-20 castor oil, P. dilaurate	PEG-100 lanolin, P. stearate
PEG-4 stearyl stearate	PEG-20 dioleate, P. distearate	PEG-120 distearate
PEG-4 tallate	PEG-20 glyceryl laurate	PEG-150 dilaurate, P. dioleate
PEG-5 castor oil, P. cocamine	PEG-20 glyceryl oleate	PEG-150 distearate, P. lanolin
PEG-5 C12-C18 alcohols	PEG-20 glyceryl stearate	PEG-150 laurate, P. oleate
PEG-5 glyceryl isostearate	PEG-20 glyceryl triisostearate	PEG-150 stearate
PEG-5 glyceryl sesquioleate	PEG-20 glyceryl tristearate	PEG-200 castor oil
PEG-5 glyceryl stearate	PEG-20 hydrogenated castor oil	PEG-200 glyceryl stearate
PEG-5 glyceryl triisostearate	PEG-20 hydrogenated lanolin	PEG-200 hydrogenated castor oil

Functions

PEG-200 laurate, P. oleate
 PEG-400 laurate
 Phosphate esters
 Phosphated amine oxides
 Phospholipids
 Poloxamer 101, 145, 122, 123, 124
 Poloxamer 181, 142, 184, 185, 235, 237
 Poloxamer 238, 334, 338, 407
 Polyglyceryl-2 oleate
 Polyglyceryl-2 polyhydroxystearate
 Polyglyceryl-2 sesquiossearate
 Polyglyceryl-2 stearate
 Polyglyceryl-2 PE/1-4-distearate
 Polyglyceryl-2 PE/1-4 stearate
 Polyglyceryl-3 distearate, P. dioleate
 Polyglyceryl-3 distearate
 Polyglyceryl-3 methylglucose distearate
 Polyglyceryl-3 oleate, P. polyricinoleate
 Polyglyceryl-3 stearate
 Polyglyceryl-4 oleate, P. stearate
 Polyglyceryl-6 dioleate, P. distearate
 Polyglyceryl-6 laurate, P. myristate
 Polyglyceryl-6 oleate, P. polyricinoleate
 Polyglyceryl-6 stearate
 Polyglyceryl-8 oleate
 Polyglyceryl-10 decanoate
 Polyglyceryl-10 distearate
 Polyglyceryl-10 dioleate, P. dipalmitate
 Polyglyceryl-10 distearate, P. isostearate
 Polyglyceryl-10 laurate, P. linoleate
 Polyglyceryl-10 mixed fatty acids
 Polyglyceryl-10 myristate
 Polyglyceryl-10 oleate
 Polyglyceryl-10 pentastearate
 Polyglyceryl-10 stearate
 Polyglyceryl-10 tetrastearate
 Polyglyceryl-10 trioleate
 Polyoxyethylene polyoxypropylene glycol
 Polyquaternium-5, -11
 Polysorbate 20, 21, 40, 60, 61
 Polysorbate 65, 80, 81, 85
 Potassium alginate, P. cetyl phosphate
 Potassium laurate, P. myristate
 Potassium tallowate
 PPG-1-PEG-9 lauryl glycol ether
 PPG-2-ceteareth-9
 PPG-3 isosteareth-9
 PPG-3 PEG-6 oleyl ether
 PPG-5-buteth-7
 PPG-5-ceteth-20
 PPG-5-ceteth-10 phosphate
 PPG-8 oleate
 PPG-10 cetyl ether phosphate
 PPG-12-PEG-50 lanolin
 PPG-15 stearyl ether
 PPG-24-buteth-27
 PPG-25 laurath-25
 PPG-26-buteth-26
 PPG-26 oleate
 PPG-36 oleate
 Propylene glycol alginate, P.g. dioleate
 Propylene glycol hydroxystearate
 Propylene glycol laurate, P.g. ricinoleate
 Propylene glycol ricinoleate SE
 Propylene glycol stearate
 Propylene glycol stearate, SE
 Quaternium-33
 Rapeseedamidopropyl ethyldimonium ethosulfate
 Rice (*Oryza sativa*) bran wax
 Ricinoleamide DEA
 Ricinoleic acid
 Saponins
 Selenium protein complex
 Silicone quaternium-5, -6
 Sodium acrylates/acrylonitrile isodecanoate crosspolymer
 Sodium capryl lactylate
 Sodium carboxomer
 Sodium cetyl sulfate

Sodium C12-15 pareth-15 sulfonate
 Sodium isostearoyl lactylate
 Sodium laureth-17 carboxylate
 Sodium lauroyl lactylate
 Sodium lauryl sulfate
 Sodium nonoxynol-6 phosphate
 Sodium octyl sulfate
 Sodium oleate
 Sodium oleyl sulfate
 Sodium phosphate
 Sodium stearoyl lactylate
 Sorbeth-20
 Sorbitan isostearate, S. laurate
 Sorbitan oleate, S. palmitate
 Sorbitan sesquiossearate
 Sorbitan sesquioleate, S. sesquisearate
 Sorbitan stearate, S. trisostearate
 Sorbitan trioleate, S. tristearate
 Soyamidopropyl dimethylamine
 Soyamine
 Stearamide DEA
 Stearamide DIBA-stearate
 Stearamidoethyl diethylamine
 Stearamidopropyl dimethylamine lactate
 Stearamidopropyl PG-dimonium chloride phosphate
 Stearamine
 Stearamine oxide
 Steareth-2, -4, -6, -7, -10, -11, -13
 Steareth-2 phosphate
 Steareth-15, -20, -21, -30, -100
 Stearic acid
 Sucrose cocoate, S. distearate
 Sucrose stearate
 Synthetic beeswax
 Tallow glyceride, acetylated hydrogenated
 Tallowamide DEA

Tallowamidopropyl dimethylamine
 Talloweth-6
 Tetrasodium dicarboxyethyl stearyl sulfosuccinamide
 TEA-acrylates/acrylonitrile copolymer
 Tissue extract
 Tricetareth-4 phosphate
 Trideceth-3, -5, -6, -7, -8
 Trideceth-9, -10, -12, -15
 Tridecyl ethoxylate
 Triethanolamine
 Trilaureth-4 phosphate
 Triolein
 Trisodium HEDTA
 Tristearin

Enzyme

Fermented vegetable
 Ganoderma lucidum oil
 Lipase
 Papain
 Soy (Glycine soja) protein
 Superoxide dismutase

Essential oil

Aesculus chinensis extract
 Artemisia apiacea extract
 Brassica rapa-depressa extract
 Caraway (*Carum carvi*) oil
 Cardamon (*Elettaria cardamomum*) oil
 Clove (*Eugenia caryophyllus*) oil
 Eclipta alba extract
 Eucalyptus globulus oil
 Euphorbia fortunei extract
 Euterpe precatoria extract
 Hierochloa odorata extract
 Kadsura heteroloca extract



Trivent Chemical Company, Inc.

4266 US Rt. 1, P.O. Box 194 • Monmouth Junction, NJ 08852 • 908-438-0995 • Fax 908-438-1020

Functions

Ligustrum lucidum extract	PVM/MA decadiene crosspolymer	Lauramidopropyl betaine
Lysimachia torenum-graecum extract	PVP/Dimethiconylacrylate/polycarbamyl/polyglycol ester	Lauryl betaine
Melaleuca bracteata extract	PVP/dimethylaminoethylmethacrylate copolymer	Myristamidopropyl dimethylamine dimethicone copolyol phosphate
Melaleuca hypericifolia extract	PVP/dimethylaminoethylmethacrylate/polycarbamyl/polyglycol ester	Myristamine oxide
Melaleuca symphocarp extract	PVP/eicosene copolymer	Octyldodecyl benzoate
Melaleuca uncinata extract	PVP/hexadecene copolymer	Oleamide DEA, C. MIPA
Melaleuca wilsonii extract	PVP/hydrolyzed wheat protein copolymer	Oleyl betaine
Nasturtium sinensis extract	Rice peptide	Palm kernelamide DEA
Nelumbium speciosum extract	Sericin	PEG-3 lauramine oxide
Paulownia imperialis extract	Shea butter (Butyrospermum parkii)	PPG-15 stearyl ether benzoate
Rosemary (Rosmarinus officinalis) oil	Shellac	PEG-7000
Sclerum spp. extract	Sodium C12-15 pareth-7 sulfonate	Sodium cocoamphacetate
Trichomonas japonica extract	Sodium hyaluronate	Sodium cocoyl isethionate
Withania somniferum extract	Soluble collagen	Sodium laureth sulfate
Yuzu oil	Soluble keratin	Sodium lauroyl wheat amino acids
Ziziphys jujuba extract	Soluble wheat protein	Sodium octoxynol-2 ethane sulfonate
	TEA-acrylates/acrylonitril copolymer	Soyamidopropyl betaine
	Tosylamide/epoxy resin	Tallowamide MEA
	Tricontanyl PVP	
	Trithonium hydrolyzed collagen ethosulfate	
	Wheat peptide	
Exfoliant	Fixative	Foam stabilizer
Apricot (Prunus armeniaca) kernel powder	Acrylates copolymer	Babassuamidopropylamine oxide
Glycolic acid	Adipic acid/dimethylaminohydroxypropyl diethylene triamine copolymer	Behenamine oxide
Jojoba (Buxus chinensis) seed powder	AMP-acrylates copolymer	Caprylyl pyrrolidone
Lactic acid	Hydrolyzed zein	Cetamine oxide
Papain	Methacryloyl ethyl betaine/acrylates copolymer	Cocamide DEA, C. MEA, C. MIPA
PEG 11-Avocado Glycerdies	Methyl rosinat	Cocamidopropyl betaine
Willow (Salix alba) bark extract	Polyquaternium-4, -10, -29	Cocamidopropyl hydroxysultaine
	PPG-20 methyl glucose ether	Cocamidopropyl lauryl ether
	Sodium polystyrene sulfonate	Cocamidopropylamine oxide
Fiber	Flavor (aroma)	Cocamine oxide
Corn (Zea mays) cob powder	Benzaldehyde	Dihydroxyethyl C12-15 alkoxypropylamine oxide
Nylon-66	Caraway (Carum carvi) oil	Dihydroxyethyl cocamine oxide
Oat (Avena sativa) bran, meal	Cardamon (Elettaria cardamomum) oil	Dihydroxyethyl tallowamine oxide
Rayon	Cinnamon (Cinnamomum cassia) oil	Erucamidopropyl hydroxysultaine
	Clove (Eugenia caryophyllus) oil	Hydroxypropyl methylcellulose
	Ethyl vanillin	Isostearamide DEA
	Eucalyptus globulus oil	Lauramide DEA, L. MEA
	Flavor (aroma)	Lauramidopropylamine oxide
	Glutamic acid	Lauramine oxide
	Glycyrhethinic acid	Laureth-10
	Glycyrhizic acid	Lauric-linoleic DEA
	Glycyrhizin, ammoniated	Lauroyl-linoleyl diethanolamide
	Methyl salicylate	Lauroyl-myristoyl diethanolamide
	Orange (Citrus aurantium dulcis) oil	Lauryl pyrrolidone
	Peppermint (Mentha piperita) oil	Linoleamide MEA
	Rosemary (Rosmarinus officinalis) oil	Myristamide DEA, M. MEA
	Sodium glycyrhizinate	Oleamide MEA
	Thymol	Palmitamide MEA
	Vanillin	PEG-3 lauramide
		PEG-4 oleamide
		Ricinoleamide MEA
		Sesamide DEA
		Wheat germamide DEA
	Foam booster	Foamer
	Alkyldimethylamine oxide	Ammonium laureth sulfate
	Babassuamidopropyl betaine	Ammonium laureth-5 sulfate
	Babassuamidopropylamine oxide	Ammonium laureth-12 sulfate
	Caprylyl pyrrolidone	Ammonium lauryl sulfate, A. 1. sulfosuccinate
	Carageenan (Chondrus crispus)	Ammonium myreth sulfate
	Cocamide DEA, C. MIPA	Ammonium nonoxynol 4 sulfate
	Cocamidopropyl betaine	Capryl caprylylglucoside
	Cocamidopropyl dimethylamine lactate	Cetyl betaine
	Cocamidopropyl hydroxysultaine	Cocamide
	Coco-betaine	Cocamidopropyl dimethylamine
	Coco/oleamidopropyl betaine	Cocamidopropyl dimethylamine lactate
	Cocoyl amido hydroxy sulfo betaine	DEA-laureth sulfate
	Cocoyl monoethanolamide ethoxylate	DEA lauryl sulfate
	DEA-hydrolyzed lecithin	Decyl glucoside
	Dimethyl lauramine	Disodium caproamphodiacetate
	Disodium cocamide MEA-sulfosuccinate	Disodium caproamphodipropionate
	Disodium cocoamphodiacetate	Disodium capryloamphodiacetate
	Disodium lauramide MEA-sulfosuccinate	Disodium cocoamphodipropionate
	Disodium laureth sulfosuccinate	Disodium lauroamphodiacetate
	Lauramide MIPA	Disodium lauroamphodipropionate
		Disodium lauryl sulfosuccinate
		Disodium oleamide MEA-sulfosuccinate

Functions

Disodium oleate
Disodium PEG-4 cocamidopropylamine sulfosuccinate
Isostearamidopropylamine oxide
Lauryl glucoside
Methyl gluceth-20
MEA-laureth sulfate
Mixed isopropanolamines myristate
MIPA-lauryl sulfate
PEG-40 sorbitan laurate
PEG lauryl ether sulfate
Potassium cocate, P. lauryl sulfate
Quillaja saponaria extract
Sodium caproamphacetate
Sodium caprylamphoacetate
Sodium caprylamphohydroxypropylsulfonate
Sodium cocamphoacetate
Sodium cocamphopropionate
Sodium C12-15 pareth-25 sulfate
Sodium C12-15 pareth-3 sulfonate
Sodium C12-15 pareth-15 sulfonate
Sodium C14-16 olefin sulfonate
Sodium deceth sulfate
Sodium laureth-2 sulfate
Sodium laureth-3 sulfate
Sodium laureth-7 sulfate
Sodium lauriminodipropionate
Sodium laurylether sulfosuccinate
Sodium lauryl sulfate, S. I. sulfoacetate
Sodium lauryl sulfosuccinate
Sodium magnesium laureth sulfate
Sodium myreth sulfate, S. myristyl sulfate
Sodium trideceth sulfate
Sodium tridecyl sulfate
TEA-dodecylbenzenesulfonate
TEA-laureth sulfate
TEA-lauryl collagen amino acids
TEA-lauryl keratin amino acids
TEA-lauryl sulfate
TEA-palm kernel sarcosinate
Wheat germamidopropyl betaine
Yucca vera extract

Fragrance
Chamaecyparis obtusa oil
Orange (Citrus aurantium dulcis) oil
Peppermint (Mentha piperita) oil
Phenethyl alcohol

Fragrance solvent
Benzyl benzoate
Diethyl phthalate
Triacetin
Triethyl citrate

Fungicide
Astrocaryum murumuru extract
Azadirachta indica extract
Capitan
Diiodomethyltolylsulfone
Ficus racemosa extract
Hexetidine
Ligusticum jeholense extract
Maunitha flexuosa extract
Melaleuca symphyocarp extract
Melia australasica extract
Melia azadirachta extract
Mushroom (Cordyceps sabolifera) extract
Mushroom (Coriolus versicolor) extract
Sodium undecylenate
Tea tree (Melaleuca alternifolia) oil
Thiabendazole
Undecylenamide MEA
Zinc undecylenate
Ziziphus jujuba extract

Gellant
Acrylic acid/acrylonitril copolymer
Agar
Alginate

Aluminum distearate, A. tristearate
Ammonium acrylates/acrylonitril copolymer
Behenic acid
Calcium alginate
Carbomer
Carboxymethylchitosan
N,O-Carboxymethylchitosonium
Cartagenan (Chondrus crispus)
Ceresin
Cetearyl candelillate
Dibenzylidene sorbitol
Ethylene/acrylic acid copolymer
Ethylene/VA copolymer
Gellan gum
Hexanediol behenyl beeswax
Hydrogenated jojoba oil
Hydrogenated jojoba wax
Hydroxysebacic acid
Jojoba wax
Laneth-5, -15
Monimorillonite
Myreth-3-octanoate
Octacosanyl stearate
Oleth-3 phosphate
Oleth-10 phosphate
Poloxamer 105, 123, 124, 185, 235
Poloxamer 237, 238, 338, 407
Polyethylene
Polyethylene, oxidized
Polyquaternium-31
Potassium alginate, P. chloride
Sodium nonoxynol-6 phosphate
Sodium tallowate
Synthetic beeswax
TEA-acrylates/acrylonitril copolymer
Tribehenin

Glosser
C18-36 acid glycol ester
Diphenyl dimethicone
Methyl gluceth-10
Octyldodecyl lactate
Phenyl methicone, P. trimethicone
Polyglyceryl-2 dioleate
Polyisobutene
Polyisobutene/isohexapentacontahexactane
Polyisobutene/isooctahexacontane
Polymethacrylamidopropyltrimonium chloride
PPG-10 methyl glucose ether
PPG-36 oleate
Tea (Camellia sinensis) oil
Tribehenin

Hair care
Gentiana scabra extract
Maidenhair fern extract
Nicotinamide
Nicotinic acid
Paeonia lactiflora extract
Watercress (Nasturtium officinale) extract

Hair conditioner
Amino bispropyl dimethicone
Amodimethicone
AMPD-isostearyl hydrolyzed collagen
Aqua Ichthammol
Babassu (Orbignya oleifera) oil
Babassuamidopropalkonium chloride
Behenamidopropyl dimethylamine
Behenamidopropyl hydroxyethyl dimonium chloride
Behenitrimonium chloride
Biotin
Bishydroxyethyl bisceryl malonamide
Borageamidopropyl phosphatidyl PG-dimonium chloride
Brazil nut (Bertholletia excelsa) oil

Cetearyl trimonium methosulphate
Cetrimonium bromide, C. chloride
Cetyl pyridinium chloride
Chia (Salvia hispanica) oil
Chrysanthemum morifolium extract
Cinchona succirubra extract
Cocamidopropyl dimethylamine propionate
Coccinea indica extract
Cocodimonium hydroxypropyl hydrolyzed collagen
Cocodimonium hydroxypropyl hydrolyzed keratin
Cocodimonium hydroxypropyl silk amino acids
Cocodimonium hydroxypropyl hydrolyzed wheat protein
Cocodimonium hydroxypropyloxyethyl cellulose
Cocotrimonium chloride
Collagen amino acids
Cyclomethicone
L-cysteine HCL
Dibenzylidimonium methosulfate
Dicetyltrimonium chloride
Dicocodimonium chloride
Dihydroxyethyl tallowamine oleate
Dimethicone
Dimethicone copolyol acetate, D. c. almondate
Dimethicone copolyol amine
Dimethicone copolyol bishydroxyethylamine
Dimethicone copolyol isosteate, D. c. laurate
Dimethicone copolyol oliveate
Dimethicone hydroxypropyl trimonium chloride
Dimethyl lauramine dimer diisoleate
Dioleamidoethyl hydroxyethylmonium methosulfate
Dipalmityloxyethyl hydroxyethylmonium methosulfate
Diphenyl dimethicone
Dilalldimonium chloride
N-Dodecyl-N,N-dimethyl-N-(dodecyl acetate) ammonium chloride
Eniada phaseoloides extract
Ethyl ester of hydrolyzed animal protein
Gelatin
Ginseng hydroxypropyltrimonium chloride butylene glycol
Hematin
Honey (Mel)
Hydrolyzed collagen
Hydrolyzed hair keratin
Hydrolyzed vegetable protein
Hydrolyzed wheat protein/dimethicone copolyol acetyl copolymer
Hydrolyzed wheat protein hydroxypropyl polysiloxane
Hydroxyethyl cetyltrimonium phosphate
Hydroxypropyltrimonium hydrolyzed collagen
Hydroxypropyl trimonium hydrolyzed wheat protein polysiloxane copolymer
Hysop (Hysopus officinalis) extract
Inga edulis extract
Isostearamidopropylamine oxide
Isostearyl hydrolyzed collagen
Keratin amino acids
Kiwi (Actinidia chinensis) fruit extract
Kola (Cola acuminata) extract
Laminaria japonica extract
Laurtrimonium chloride
Lauryl hydroxypropyl trimonium polysiloxane copolymer
Lauryldimethylamine isosteate
Lauryldimonium hydroxypropyl hydrolyzed collagen
Lauryldimonium hydroxypropyl hydrolyzed wheat protein
Linoleamidopropyl dimethylamine dimer diisoleate
Linoleamidopropyl dimethylamine
Lysimachia foenum-graecum extract
Melaleuca hypericifolia extract
Ocimum sanctum extract
Olealkonium chloride

Functions

Oleyl dimethylamidopropyl ethonium ethosulfate
 Palmitamidodecanediol
 Panthenyl ethyl ether
 Paulownia imperialis extract
 Peach (*Prunus persica*) leaf extract
 PEG-2 cocomonium chloride
 PEG-120 jojoba acid/alcohol
 PG-hydroxycellulose lauryldimonium chloride
 PG-hydroxyethylcellulose cocodimonium chloride
 PG-hydroxyethylcellulose lauryldimonium chloride
 PG-hydroxyethylcellulose stearyldimonium chloride
 Phenyl trimethicone
 Phospholipids
 Phytantriol
 Polyoxethylene polyoxypropylene glycol
 Polypropylene glycol
 Polyquaternium-4, -6, -7, -10
 Polyquaternium-22, -28, -39
 PPG-5-ceteih-10 phosphate
 Propyltrimonium hydrolyzed collagen
 Propyltrimonium hydrolyzed soy protein
 Propyltrimonium hydrolyzed wheat protein
 Quaternium-18, -75, -81, -82
 Quaternium-79 hydrolyzed keratin
 Quaternium-79 hydrolyzed silk
 Sambucus nigra extract, oil
 Sesamidopropalkonium chloride
 Silicone quaternium-1, -8
 Sodium cocamphoacetate
 Sodium cocoyl hydrolyzed collagen
 Sodium polyvinylene sulfonate
 N-Soya-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate
 Steapyrium chloride
 Stearalkonium chloride
 Stearamidopropyl dimethylamine
 Steardimonium hydroxypropyl hydrolyzed wheat protein
 Steartrimonium chloride
 Steartrimonium hydroxyethyl hydrolyzed collagen
 N-Stearyl-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate
 Stenocalyx micalli extract
 Sulfur
 Tallowbenzylidimethylammonium chloride, hydrogenated
 Tallowtrimonium chloride
 Tea (*Camellia sinensis*) oil
 TEA-cocoyl hydrolyzed soy protein
 Thenoyl methionate
 Trimethylsilylamodimethicone
 Wheat amino acids

Hair set resin polymer

Acrylates/acrylamide copolymer
 Acrylates/PVP copolymer
 Acrylates/hydroxyesters acrylates copolymer
 Acrylates/octylarylamide copolymer
 AMP-acrylates copolymer
 Butylester of PVM-MA copolymer
 Carboxylated vinylacetate terpolymer
 Diglycol/CHDM/isophthalates/SIP copolymer
 Eclipta alba extract
 Ethyl ester of PVM/MA copolymer
 Hydroxypropyl chitosan
 Isopropyl ester of PVM/MA copolymer
 Octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer
 Polymethacrylamidopropyltrimonium chloride
 Polypropylene glycol oligosuccinate
 PVP
 PVP/dimethylaminoethylmethacrylate copolymer
 PVP/Polycarbonyl polyglycol ester
 PVP/VA copolymer
 PVP/VA-vinyl propionate copolymer
 Sodium polyacrylate

VA/butyl maleate/isobornyl acrylate copolymer
 VA/crotonates/vinyl neodecanoate copolymer
 VA/crotonates/vinyl propionate copolymer
 VA/crotonates copolymer
 Vinyl caprolactam/PVP/
 dimethylaminoethylmethacrylate copolymer

Hair sheen

Maidenhair fern extract
 Tetrahydroxypropyl methicone

Hair waving

Ammonium thioglycolate, A. thiolactate
 Argania spinosa oil
 L-cysteine HCL
 Cystine
 Diammonium dithiodiglycolate
 Dilauryl thioldipropionate
 Ethanolamine sulfate, E. thioglycolate
 Ethanolamine thiolactate
 Glyceryl thioglycolate
 Hydroxymethyl dioxazabicyclooctane
 Jojoba esters
 Monoethanolamine thiolactate
 Shea butter, ethoxylated
 Sodium thioglycolate
 Thiolglycerol
 Thioglycolic acid
 Thiolactic acid

Humectant

Acetamide MEA
 Acetyl monoethanolamine
 6-(N-Acetyl amino)-4-oxahexyltrimonium chloride
 Adenosine phosphate
 Ammonium lactate
 Atelocollagen
 Calcium pantothenate
 Calcium stearoyl lactylate
 Carboxymethyl chitin
 Carboxymethyl chitosan succinamide
 Chitosan PCA
 Cholesteryl hydroxystearate
 Collagen amino-polysiloxane hydrolyzate
 Colloidal oatmeal
 Copper PCA methylsilanol
 Dimethicone copolyol laurate
 Dipotassium glycyrrhizinate
 Ethyl ester of hydrolyzed silk
 Fatty quaternary amine chloride complex
 Glucose glutamate
 Glycereth-4,5-lactate
 Glycereth-7, -12, -26
 Glycenn
 Honey extract
 Hydrogenated passion fruit oil
 Hydrolyzed casein
 Hydrolyzed fibronectin
 Hydrolyzed glycosaminoglycans
 Hydrolyzed oat protein
 Hydrolyzed silk
 Hydrolyzed soy protein
 Hydroxypropyl chitosan
 Hydroxypropyltrimonium hydrolyzed casein
 Hydroxypropyltrimonium hydrolyzed silk
 Hydroxypropyltrimonium hydrolyzed soy protein
 Hydroxypropyltrimonium hydrolyzed wheat protein
 Keratin amino acids
 Lactamide DGA, MEA
 Lactamidopropyl trimonium chloride
 Lactic acid
 Lactose
 Lauroyl lysine
 Maltitol
 Mannitol
 Methyl gluceth-10, -20
 Natto gum
 Oat (*Avena sativa*) extract, protein
 Panthenol

Panthenyl ethyl ether
 PCA
 PEG-4
 Polyamino sugar condensate
 Potassium lactate
 Propylene glycol
 Propyltrimonium hydrolyzed collagen
 Propyltrimonium hydrolyzed soy protein
 Propyltrimonium hydrolyzed wheat protein
 Quaternium-22
 Rice (*Oryza sativa*) germ oil
 Sea Salts (Mans salt)
 Shea butter (*Butyrospermum parkii*)
 Silk powder
 Sodium behenoyl lactylate
 Sodium caproyl lactylate
 Sodium cocoyl lactylate
 Sodium hyaluronate
 Sodium isostearoyl lactylate
 Sodium lactate, S. lauroyl lactylate, S. PCA
 Sodium polyglutamate
 Sodium stearoyl lactylate
 Sorbitan laurate
 Sorbitan sesquiossearate
 Sorbitol
 Sphingolipids
 TEA-PCA
 Urea

Hydrotrope

Ammonium cumenesulfonate
 Ammonium xylenesulfonate
 Cetamine oxide
 Cocamidopropylamine oxide
 Lauramine oxide
 Potassium toluenesulfonate
 PPG-2-isodeceth-4, -6, -9, -12
 Sodium cumene sulfonate
 Sodium laureth-13-carboxylate
 Sodium toluene sulfonate
 Sodium xylene sulfonate
 Trideceth-19-carboxylic acid

Intermediate

Caprylic acid
 Deceth-3
 Diethyl succinate
 Dimethylaminopropylamine
 DM hydantoin
 Dodecylbenzene sulfonic acid
 Ethylene dichloride
 4-Fluoro 3-nitro aniline
 Lauramine
 Methyl benzoate, M. cocoate
 Methyl isostearate, M. laurate
 Methyl myristate, M. palmitate
 Oleic acid
 Ricinoleic acid
 Tall oil acid
 Tallow acid

Lathering agent

Ammonium cocoyl sarcosinate
 Ammonium C12-15 alkyl sulfate
 Ammonium lauroyl sarcosinate
 Cocamide MEA ethoxylate
 Cocamidopropyl dimethylaminohydroxypropyl hydrolyzed collagen
 Lauroyl sarcosine
 Myristoyl sarcosine
 Sodium cocoyl sarcosinate
 Sodium lauroyl sarcosinate
 Sodium methyl cocoyl laurate
 Sodium myristoyl sarcosinate
 TEA-cocoyl sarcosinate
 TEA-lauroyl sarcosinate

Lubricant

Aluminum salt octenyl succinate
 Amodimethicone

Functions

Boron nitride
Calcium aluminum borosilicate
Calcium stearate
Caprylic/capric triglyceride
Coceth-7 carboxylic acid
Coconut (Cocos nucifera) oil
Cyclomethicone
Diisodecyl adipate
Diisostearyl fumarate
Dimethicone copolyol
Glyceryl isostearate, G. oleate
Glyceryl polymethacrylate
Gold of Pleasure oil
Hyaluronic acid
Hydrogenated coconut oil
Hydrogenated cottonseed oil
Hydrogenated palm oil
Hydrogenated soybean/cottonseed oil
Hydrogenated soybean oil
Hydrogenated vegetable oil
Hydrolyzed oat flour
Hydroxypropyl guar
Isodecyl stearate
Isopropyl lanolate
Isostearyl diglyceryl succinate
Jojoba esters
Lanolin oil
Laureth-3 phosphate
Magnesium myristate, M. stearate
Mango (Mangifera indica) oil
Mineral oil (Paraffinum liquidum)
Mink oil
Monostearyl citrate
Neatsfoot oil
Oleostearine
Partially hydrogenated soybean oil
PEG-2 stearate
PEG-4 dilaureate
PEG-5M
PEG-9M
PEG-23M
PEG-27 lanolin
PEG-30 lanolin
PEG-40 lanolin, P. stearate
PEG-45M
PEG-90M
PEG-160M
PEG/PPG-17/6 copolymer
Pentaerythrityl tetrapalargonate
Petrolatum
Phenethyl dimethicone
Phenyl methicone
Polyacrylamidomethylpropane sulfonic acid
Polybutene
Polydimethicone copolyol
Polyglycerol ester of mixed vegetable fatty acids
Polymethylsilsesquioxane
Potassium laurate, P. myristate
Potassium palmitate
PPG-2 myristyl ether propionate
PPG-3 myristyl ether
PPG-9-buteth-12
PPG-11 stearyl ether
PPG-12-buteth-16
PPG-12-PEG-50 lanolin
PPG-14 butyl ether
PPG-20 cetyl ether
PPG-20-buteth-30
PPG-24-buteth-27
PPG-28-buteth-35
PPG-36 oleate
PPG-40 butyl ether
Quaternium-79 hydrolyzed keratin
Quaternium-79 hydrolyzed silk
Rice (Oryza sativa) starch
Shea butter (Butyrospermum parkii) extract
Shorea stenoptera butter
Silica
Stearamide MEA, S. MEA-stearate
Stearoxytrimethylsilane

Stearyl dimethicone
Trisostearyl citrate
Triolein
Trisodium HEDTA
Triundecanoin
Zinc laurate, Z. stearate

Miscellaneous

Adhesion promoter—Glyceryl/diethylene glycol/adipate crosspolymer
Analgesic—Glycol salicylate
Anesthetic—Benzocaine
Anti-elastic—Hydrolyzed Ulva lactuca extract
Anti-itching—Sodium shale oil sulfonate
Antiacid—Magnesium hydroxide, Magnesium silicate, Simethicone
Antifoam—Dimethicone silylate, Simethicone
Antilipase—Laminaria saccharina extract
Antipruritic—Coal tar
Antispasmodic—Garlic (Allium sativum) extract
Antiwrinkle—Chinese hibiscus (Hibiscus rosa-sinensis) extract
Barrier—Glyceryl/diethylene glycol/adipate crosspolymer
Cell regeneration—Glycoproteins, Hydrolyzed Ulva lactuca extract
Co-emulsifier—Cholesteryl/heptenyl/octyldodecyl lauroyl glutamate, Isododecane
Colloid—Gelatin
Cooling agent—Menthyl PCA, Menthone glycerin acetal
Detoxifier—Clover (Trifolium pratense) extract
Dye stabilizer—Uric acid
Filler—Mica
Fragrance stabilizer—2,2',4,4'-Tetrahydroxybenzophenone
Free radical scavenger—Melanin
IR filter—Corallina officinalis

Lanolin substitute—PEG-80 jojoba acid/alcohol
Lipolytic—Gelidium cartilagineum
Oxidant—Barium peroxide, Hydrogen peroxide, Urea peroxide
Oxygen carrier—Perfluorodecalin
Peroxide stabilizer—Phenacetin, Sodium stannate
Scalp stimulant—Birch (Betula alba) leaf extract
Sebostatic—Laminaria saccharina extract
Shine enhancer—Hydrolyzed wheat protein hydroxypropyl polysiloxane
Skin barrier lipid—Ceramide 3, N(27-Stearoyloxy-heptacosanoyl) phytosphingosine
Skin clarifier—Oat (Avena sativa) bran extract
Skin purifier—Birch (Betula alba) leaf extract
Substantivity—Dimethicone copolyol bishydroxyethylamine, Dimethicone hydroxypropyl trimonium chloride, Trimethylsilylamodimethicone
Sunless tanning—Acetyl tyrosine, Eclipta alba extract in white emulsion
Tonic—Kiwi (Actinidia chinensis) fruit extract, Matricaria (Chamomilla recutita) extract, Orange (Citrus aurantium dulcis) peel extract
Viscosity stabilizer—Diisodecyl adipate
Spreading agent—Stearyl heptanoate
Wound healing—Comfrey (Symphytum officinale) leaf extract
Waterproofing agent—PVP/eicosene copolymer, PVP/hexadecene copolymer, Tricontanyl PVP

Moisture barrier

Acrylates/octylarylamide copolymer
Beta-glucan
C16-18 alkyl methicone
Cholesterol
Glycolipids
Isoeicosane

BERNEL

CHEMICAL COMPANY

Up to date, innovative technology for the cosmetic industry has been the driving force behind Bernel Chemical Company since its founding in 1982. Combining over 60 years of cosmetic expertise and marketing knowledge, we have introduced more than 20 raw materials for use by the cosmetic chemist.

Our product is innovation. Finding unique materials, such as MARRIX SF and CUPL[®] PIC, that contribute to the growth of our customers has established Bernel products worldwide.

BERNEL

CHEMICAL COMPANY

174 Grand Ave., Englewood, NJ 07631
Phone: 201-569-8934 • Fax: 201-569-1741

Functions

Isohexadecane	Embolica officinalis extract	Methylsilanol elastinate, M. mannuronate
Lanosterol	Ethyl minkate	Milk amino acids
Octyl pelargonate, O. stearate	Eugenia jambolana extract	Mineral oil (Paraffinum liquidum)
Polyisobutene	Evening primrose (Oenothera biennis) extract, oil	Molybdenum aspartate
Polyisobutene/isohexapentacontahexane	Galla sinensis extract	Mouriri apiranga extract
Polyisobutene/isooctahexacontane	Ganoderma lucidum oil	Natto gum
Silica silvate	Ginseng (Panax ginseng) extract	Nelumbium speciosum extract
Trihydroxypalmitamidohydroxy propyl myristyl ether	Gleditsia sinensis extract	Neopentyl glycol dicaprate
Trimethylsiloxysilicate	Glycereth-12	Oat (Avena sativa) protein
	Glycerol alginate, G. collagenate	Octyl hydroxystearate
	Glycerol polymethacrylate	Ophiopogon japonicus extract
	Glycolic acid	Orange (Citrus aurantium dulcis) peel wax
	Glycolipids	Palmetto extract
	Glycosaminoglycans	Pantethine
	Glycosphingolipids	Panthenyl ethyl ether
	Gnetum amazonicum extract	Paraffin
	Grape (Vitis vinifera) seed oil	Partially hydrogenated soybean oil
	Hazel (Corylus avellana) nut oil	Peanut (Arachis hypogaea) oil
	Honey extract	Pecan (Carya illinoensis) oil
	Hyaluronic acid	PEG-4, -6, -8, -12
	Hybrid safflower (Carthamus tinctorius) oil	PEG-70 mango glycerides
	Hydrogenated castor oil	PEG-75 shea butter glycerides
	Hydrogenated coconut oil	PEG-75 shorea butter glycerides
	Hydrogenated cottonseed oil	PEG-100 stearate
	Hydrogenated lecithin	Pentaerythritol isostearate/caprate/caprylate/adipate
	Hydrogenated palm oil	Pentaerythritol stearate/caprate/caprylate/adipate
	Hydrogenated polyisobutene	Pentylene glycol
	Hydrogenated soybean oil	Perfluoropolyethylisopropyl ether
	Hydrogenated soybean/cottonseed oil	Petrolatum
	Hydrogenated vegetable oil	Petroleum wax
	Hydrolyzed carbolipoprotein	Pfaffia spp. extract
	Hydrolyzed collagen	Pistachio (Pistacia vera) nut oil
	Hydrolyzed elastin	Placental protein
	Hydrolyzed fibronectin	Plankton extract
	Hydrolyzed glycosaminoglycans	Polyamino sugar condensate
	Hydrolyzed keratin	Polybutene
	Hydrolyzed milk protein	Polyunsaturated fatty acids
	Hydrolyzed oils	Potassium DNA, P. lactate, P. PCA
	Hydrolyzed pea protein	PPG-8/SMDI copolymer
	Hydrolyzed placental protein	PPG-20 methyl glucose ether distearate
	Hydrolyzed rice protein	Propylene glycol dicaprylate/dicaprate
	Hydrolyzed transgenic collagen	Propylene glycol dioctanoate
	Hydrolyzed serum protein	Pumpkin (Cucurbita pepo) seed oil
	Hydrolyzed silk	Quinoa (Chenopodium quinoa) extract
	Hydrolyzed sweet almond protein	Rapeseed (Brassica campestris) oil
	Hydrolyzed wheat protein	Rehmannia chinensis extract
	Hydroxyethyl chitosan	Rice (Oryza sativa) bran oil
	Inositol	Rose Water
	Isodecyl salicylate	Royal jelly extract
	Isostearyl hydrolyzed animal protein	Saccharide isomerase
	Jajoba (Buxus chinensis) oil	Saccharomyces lysate extract
	Jajoba esters	Saccharomyces/soy protein ferment
	Keratin amino acids	Safflower (Carthamus tinctorius) oil
	Kiwi (Actinidia chinensis) fruit extract	Selenium aspartate, S. protein complex
	Kola (Cola acuminata) extract	Sericin
	Kukui (Aleurites moluccana) nut oil	Serum albumin
	Lactamide DGA, L. MEA	Sesame (Sesamum indicum) oil
	Lactic acid	Shea butter (Butyrospermum parkii)
	Lactobacillus/whey ferment	Shea butter (Butyrospermum parkii) extract
	Lactococcus hydrolysate	Shorea stenoptera butter
	Lactoyl methylsilanol elastinate	Silk amino acids
	Lanolin alcohol	Sodium carboxymethyl beta-glucan
	Lauryl PCA	Sodium chondroitin sulfate
	Lecithin	Sodium DNA, S. hyaluronate
	Lesquerella fendleri oil	Sodium lactate, S. PCA
	Liposomes	Soluble collagen
	Lysine PCA	Soluble transgenic elastin
	Macadamia ternifolia nut oil	Soybean (Glycine soja) oil
	Magnesium aspartate	Spherical cellulose acetate
	Maltitol	Spondias amara extract
	Manganese aspartate	Squalene
	Mango (Mangifera indica) oil	Stomach extract
	Mannan	Sunflower (Helianthus annuus) seed oil
	Marine polyaminosaccharide	Superoxide dismutase
	Mauritella armata extract	Tissue extract
	Maximiliana regia extract	Tocopheryl acetate, T. linoleate
	Meadowfoam (Limnanthes alba) seed oil	Tomato (Solanum lycopersicum) extract
	Melaleuca hypericifolia extract	

Formenil (Potentilla erecta) extract
Trehalose
Tridecanoin
Vegetable oil
Walnut (Juglans regia) oil
Watercress (Nasturtium officinale) extract
Wheat (Triticum vulgare) germ extract, germ oil
Yarrow (Achillea millefolium) extract
Wheat amino acids
Yeast (Saccharomyces cerevisiae) extract (Faex)
Yogurt filtrate
Zinc aspartate
Ziziphus jujuba extract

Naturilizer
2-Aminobutanol
Aminoethyl propanediol
Aminomethyl propanediol
Aminomethyl propanol
Ammonium carbonate
Calcium hydroxide
Diethanolamine
Ethanolamine
Glucamine
Isopropanolamine
Isopropylamine
2-Methyl-4-hydroxypyridine
Morpholine
Sodium bromate
Succinic acid
Tetrahydroxypropyl ethylenediamine
Triethanolamine
Tromethamine

Oil absorbent
Hydrated silica
Polymethyl methacrylate
Silicon dioxide hydrate
Walnut (Juglans regia) shell powder

Ointment base
Borage (Borago officinalis) seed oil
Caprylic/capric/stearic triglyceride
Glyceryl cocoate
Hydrogenated coco-glycerides
Lanolin
Mink oil
Oleostearine
Tallow

Opacifier
Barium sulfate
C12-16 alcohols
Cetearyl octanoate
Cetyl myristate, C. palmitate
Cocamidopropyl lauryl ether
Glyceryl distearate
Glyceryl hydroxystearate
Glyceryl myristate, G. stearate
Glycol distearate, G. stearate
Magnesium myristate
PEG-2 distearate, P. stearate
PEG-2 stearate SE
PEG-3 distearate
Propylene glycol myristate, P. g. stearate
Stearamide
Stearamide DIBA-stearate
Stearamide MEA
Stearamide MEA-stearate
Stearamidopropyl dimethylamine lactate

Stearyl stearate
Styrene homopolymer
Styrene/acrylates copolymer
Styrene/PVP copolymer
Triisostearin PEG-6 esters

Plasticizer
Acetyl tributyl citrate
Acetyl triethyl citrate
AMP-isostearyl hydrolyzed wheat protein
AMPD-isostearyl hydrolyzed collagen
Cyclohexane dimethanol dibenzoate
Dibutyl phthalate
Diethyl phthalate
Diethylene glycol dibenzoate
Diisopropyl sebacate
Dimethicone copolyol
Dimethyl phthalate
Dipropylene glycol dibenzoate
Ethyl ester of hydrolyzed keratin
Glycerol tnbenzoate
Glycol
Hydrolyzed serum protein
Isocetyl salicylate
Isodecyl benzoate
Isocicosane
Isopropyl lanolate
Isostearyl hydrolyzed collagen
Lauroyl hydrolyzed collagen
Marine collagen
Monostearyl citrate
Neopentyl glycol dibenzoate
Octyl benzoate, O. laurate
PEG-60 Shea butter glycerides
Pentaerythrityl tetrabenzoate
Polyoxyethylene glycol dibenzoate
Polypropylene glycol dibenzoate
PPG-12-PEG-50 lanolin
PPG-20 cetyl ether
PPG-20 lanolin alcohol ether
Propylene glycol dibenzoate
Propylene glycol myristyl ether acetate
Rice (Oryza sativa) bran wax
Serum protein
Tosylamide/epoxy resin
Triacetin
Tributyl citrate
Triethyl citrate
Trimethyl pentanediol dibenzoate
Trimethylethancmbenzoate

Polish
Acrylates copolymer
Aluminum silicate
Neasfoot oil
Tallow

Polymer
Acrylamide sodium acrylate copolymer
Acrylates-VA crosspolymer
Acrylates/acrylamide copolymer
Acrylates/hydroxyesters acrylates copolymer
Acrylates/octylacrylamide copolymer
Acrylates/steareth-20 methacrylate copolymer
Adipic acid-epoxypropyl diethylenetriamine copolymer
Adipic acid/dimethylaminohydroxypropyl diethylene triamine copolymer
Ammonium acrylates copolymer

Ammonium acrylates copolymer
AMP-acrylates copolymer
AMP-isostearyl hydrolyzed collagen
Butylester of PVM-MA copolymer
Calcium carrageenan
Carboxylated vinylacetate terpolymer
Ceteareth-2 phosphate
Ceteareth-5 phosphate
Ceteareth-10 phosphate
Ceteareth-29, -34
Coco-glucoside
Cocodimonium hydroxypropyloxyethyl cellulose
C12-13 pareth-4, -9, -23
DEA-ceteareth-2-phosphate
DEA-oleth-5-phosphate
DEA-oleth-20-phosphate
Diglycol/CHDM/isophthalate/SIP copolymer
Diisopropyl dimer dilinoleate
Diisostearyl trimethylolpropane siloxy silicate
Diisostearyl dimer dilinoleate
Dilinoleic acid
Dodecanedioic acid/cetearyl alcohol/glycol copolymer
Eclipta alba extract
Ethyl ester of PVM/MA copolymer
Ethylene/acrylic acid copolymer
Ethylene/VA copolymer
Glycereth-26 phosphate
Hyaluronic acid
Hydrolyzed RNA
Hydrolyzed wheat protein polysiloxane polymer
Hydroxypropyltrimonium hydrolyzed collagen
Hydroxypropyltrimonium hydrolyzed wheat protein
Laneth-40
Lauryldimonium hydroxypropyl hydrolyzed soy protein
Methacryloyl ethyl betaine/acrylates copolymer
Octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer
Oleth-2 phosphate
Oleth-5 phosphate
PEG-3 lanolate
PEG-4 stearate
PEG-5M
PEG-7 glyceryl cocoate
PEG-8 glyceryl laurate
PEG-8/SMDI copolymer
PEG-9 castor oil
PEG-9M
PEG-11 babassu glycerides
PEG-12 palm kernel glycerides
PEG-12 stearate
PEG-14 avocado glycerides
PEG-15 glyceryl laurate
PEG-20 com glycerides
PEG-20 evening primrose glycerides
PEG-20 glyceryl oleate
PEG-23 oleate
PEG-23M
PEG-29 castor oil
PEG-42 babassu glycerides
PEG-45 safflower glycerides
PEG-45M
PEG-60 evening primrose glycerides
PEG-60 hydrogenated castor oil
PEG-75 castor oil
PEG-90M
PEG-120 distearate

Functions

PEG-150 lanolin
 PEG-160M
 PG-hydroxycellulose lauryldimonium chloride
 PG-hydroxyethylcellulose cocodimonium chloride
 PG-hydroxyethylcellulose stearyldimonium chloride
 Polyethylene, ionomer
 Polyethylene, micronized
 Polyethylene, oxidized
 Polyglyceryl-2 polyhydroxystearate
 Polymethacrylamidopropyltrimonium chloride
 Polyquaternium-6, -7, -10, -11, -22, -39
 Polysilicone-8
 Potassium alginate
 Potassium lauroyl collagen amino acids
 Potassium lauroyl hydrolyzed soy protein
 Potassium lauroyl wheat amino acids
 PPG-8/SMDI copolymer
 PPG-12/SMDI copolymer
 PPG-51/SMDI copolymer
 PVM/MA decadiene crosspolymer
 PVP/dimethylaminomethylmethacrylate copolymer
 PVP/VA copolymer
 Sodium cocoyl hydrolyzed wheat protein
 Steardimonium hydroxypropyl hydrolyzed wheat protein
 Steareth-2 phosphate
 TEA-acrylates/acrylonitril copolymer
 Tosylamide/epoxy resin
 Tosylamide/formaldehyde resin
 Trideceth-5, -6, -7, -8
 VA/butyl maleate/isobornyl acrylate copolymer
 VA/crotonates/vinyl neodecanoate copolymer
 Vinyl caprolactam/PVP/
 dimethylaminoethylmethacrylate copolymer
 Wheat (Triticum vulgare) protein
 Xanthan gum

Powder

Acrylates copolymer, spherical powder
 Attapulgit
 Boron nitride
 Calcium aluminum borosilicate
 Calcium carbonate
 Cellulose triacetate
 Corn (Zea mays) cob powder, starch
 Hydrogenated jojoba wax
 Magnesium carbonate, M. myristate
 Magnesium stearate
 Mica
 Microcrystalline cellulose
 Nylon-6
 Nylon powder
 Oat (Avena sativa) starch
 Polyamide 12
 Polyethylene
 Polymethyl methacrylate
 Polymethylsilsesquioxane
 PTFE
 Silica
 Silk powder
 Spherical cellulose acetate
 Talc
 Tapioca dextrin
 Zinc laurate

Powder, absorbent

Aluminum starch octenylsuccinate
 Clays (white, yellow, red, green, pink)
 Sorbitol
 Tapioca

Preservative

Alcohol
 Ascorbic acid
 Ascorbyl palmitate

Benzalkonium chloride
 Benzethonium chloride
 Benzoic acid
 Benzyl alcohol
 Benzylparaben
 5-Bromo-5-nitro-1,3-dioxane
 2-Bromo-2-nitropropane-1,3-diol
 Butylparaben
 Calcium propionate
 Cetrimonium bromide
 Cetyl pyridinium chloride
 Chloroxylenol
 Chlorophenesin
 o-Cymen-3-ol
 Diazolidinyl urea
 Dichlorobenzyl alcohol
 Dichlorophene
 Diiodomethyltolylsulfone
 Dimethyl hydroxymethyl pyrazole
 Dimethyl oxazolidine
 Disodium EDTA
 DMDM hydantoin
 EDTA
 Erythorbic acid
 7-Ethylbicyclooxazolidine
 Ethylparaben
 Fomistopsis officinalis oil
 Formaldehyde
 Glutaral
 Glyceryl laurate
 HEDTA
 Hexamidine diisethionate
 Hexetidine
 Imidazolidinyl urea
 Isobutylparaben
 Isopropyl sorbate
 Isopropylparaben
 MDM hydantoin
 Methenammmonium chloride
 Methyl paraben sodium
 Methylchloroisothiazolinone
 Methylidibromo glutaronitrile
 Methylisothiazolinone
 Methylparaben
 Mushroom (Cordyceps sabolifera) extract
 Myrrimonium bromide
 Pentasodium pentetate
 Pentenoic acid
 Phenethyl alcohol
 Phenol
 Phenyl mercuric acetate
 o-Phenylphenol
 Polyaminopropyl biguanide
 Polymethoxy bicyclic oxazolidine
 Potassium sorbate
 Propylparaben
 Quaternium-15
 Salicylic acid
 Sodium benzoate, S. bisulfate
 Sodium butylparaben, S. dehydroacetate
 Sodium erythorbate, S. ethyl paraben
 Sodium hydroxymethylglycinate
 Sodium metabisulfite, S. methylparaben
 Sodium o-phenylphenate
 Sodium propionate, S. propylparaben
 Sodium pyrrithione, S. salicylate
 Sodium sulfite
 Sorbic acid
 Tetrasodium EDTA
 Thimerosal
 Thymol
 Tris (hydroxymethyl) nitromethane
 Trisodium EDTA, T. HEDTA
 Usnic acid
 Zinc PCA

Propellant

Butane
 Dimethyl ether
 Hydrofluorocarbon 152a

**In the World of
Natural Waxes**

Carnauba Wax
Beeswax
Candelilla Wax



STRAHL & PITSCHE INC.

Ceresine And Ozokerite
Paraffin And Microcrystalline



There is no one else!
 Dedicated to Natural Waxes and
 Specialty Blends Since 1904



STRAHL & PITSCHE INC., P.O. BOX 1000, 210 EAST EAST NICE RD., WEST BABYLON, NY 11794 U.S.A.
 (516) 587-9000 / FAX: (516) 587-9120

Functions

Isobutane
Propane

Protein

Albumen
Atelocollagen
Bleeta hvacinthina extract
Chrysanthemum montolium extract
Cocodimonium hydroxypropyl hydrolyzed collagen
Cocodimonium hydroxypropyl hydrolyzed keratin
Cocodimonium hydroxypropyl hydrolyzed soy protein
Cocodimonium hydroxypropyl hydrolyzed wheat protein
Cocoyl hydrolyzed collagen
Collagen, C. phthalate
Collagen amino-polyvinylane hydrolyzate
Deoxyribonucleic acid
Desamido collagen
Elastin amino acids
Embryo extract
Ethyl ester of hydrolyzed animal protein
Fibronectin
Gelatin
Human placental protein
Hydrolyzed collagen
Hydrolyzed extensin
Hydrolyzed fish protein
Hydrolyzed hemoglobin
Hydrolyzed keratin
Hydrolyzed lactalbumin
Hydrolyzed milk protein
Hydrolyzed soy flour
Hydrolyzed sweet almond protein
Hydroxypropyltrimonium hydrolyzed collagen
Isostearyl hydrolyzed collagen
Keratin
Lactoterrin
Lactoglobulin
Lauryldimonium hydroxypropyl hydrolyzed collagen
Marine collagen
Methylsilanol elastinate
Potassium abietoyl hydrolyzed collagen
Potassium cocoyl hydrolyzed collagen
Potassium myristoyl hydrolyzed collagen
Potassium oleoyl hydrolyzed collagen
Potassium undecylenoyl hydrolyzed collagen
Propyltrimonium hydrolyzed collagen
Propyltrimonium hydrolyzed soy protein
Propyltrimonium hydrolyzed wheat protein
Protein hydrolysates
Quaternium-79 hydrolyzed keratin
Quaternium-79 hydrolyzed silk
Rice peptide
RNA
Serum albumin, S. protein
Silk powder

Sodium caseinate

Sodium cocoyl hydrolyzed collagen
Sodium cocoyl hydrolyzed soy protein
Sodium myristoyl hydrolyzed collagen
Sodium oleoyl hydrolyzed collagen
Sodium stearyl hydrolyzed collagen
Sodium undecylenoyl hydrolyzed collagen
Sodium/TEA-lauryl hydrolyzed collagen
Sodium/TEA-lauryl hydrolyzed keratin
Soluble collagen
Soluble keratin
Soluble wheat protein
Soy (Glycine soja) protein
Steardimonium hydroxypropyl hydrolyzed collagen
Steardimonium hydroxyethyl hydrolyzed collagen
TEA-cocoyl hydrolyzed collagen
TEA-cocoyl hydrolyzed soy protein
TEA-lauryl collagen amino acids
TEA-lauryl keratin amino acids
Trachea hydrolysate
Triethonium hydrolyzed collagen ethosulfate
Wheat (Triticum vulgare) germ extract, protein
Wheat amino acids
Wheat peptide
Wheat protein

Protein, hydrolyzed

Ethyl ester of hydrolyzed silk
Hydrolyzed casein
Hydrolyzed elastin
Hydrolyzed mushroom (Tricholoma matsutake) extract
Hydrolyzed pea protein
Hydrolyzed rice protein
Hydrolyzed serum protein
Hydrolyzed silk
Hydrolyzed soy protein
Hydrolyzed vegetable protein
Hydrolyzed wheat protein
Hydroxypropyltrimonium hydrolyzed casein
Hydroxypropyltrimonium hydrolyzed silk
Hydroxypropyltrimonium hydrolyzed soy protein
Hydroxypropyltrimonium hydrolyzed wheat protein

Reducing agent

Dimyristyl thiodipropionate
Hydrolyzed zein, iodized
Hydrolyzed zein, sulfonized
Zinc formaldehyde sulfoxylate

Refatting agent

Caprylic/capric triglyceride PEG-4 esters
Cocamide MIPA
Diisostearyl dimer dilinoleate
Hydrogenated palm kernel glycerides
Isostearyl erucate, I. isostearate
Lecithin

Liposomes

Magnesium sulfate hepta-hydrate
Octyldodecyl behenate, (O). myristate
bis-Octyldodecyl stearyl dimer dilinoleate
Octyldodecyl stearyl stearate
Octyl hydroxystearate
PEG-3 stearate
PEG-4 oleamide
PEG-6 capric/caprylic glycerides
PEG-7 glyceryl cocoate
PEG-16
Propylene glycol dipalargonate

Resin

Acrylates/hydroxyesters acrylates copolymer
Ethylene vinyl acetate
Glyceryl abietate
Methacryloyl ethyl betaine/acrylates copolymer
4-Methyl benzenesulfonamide
Polypropylene
Polyquaternium-16, -44
Sucrose benzoate

Sequestrant

Calcium acetate, C. phosphate, C. sulfate
Encapsulation and entrapment systems
Pentasodium triphosphate
Phosphoric acid
Potassium phosphate, P. sodium tartrate
Silicon dioxide hydrate
Sodium citrate, S. gluconate
Sorbitol
Tartaric acid
Tripotassium EDTA
Trisodium NTA

Silicone

Amino bispropyl dimethicone
Ammonium dimethicone copolyol sulfate
Amodimethicone
Behenoxy dimethicone
C16-18 alkyl methicone
Cetyl dimethicone copolyol
Cyclomethicone Diisostearyl trimethylolpropane siloxy silicate
Diisodecyl adipate
Diisostearyl trimethylolpropane siloxy silicate
Dimethicone
Dimethicone copolyol
Dimethicone copolyol almondate
Dimethicone copolyol isostearate
Dimethicone copolyol olivate, D. c. phthalate
Dimethicone copolyolamine
Dimethiconol fluoroalcohol dilinoleic acid
Dimethiconol hydroxystearate, D. stearate
Diphenyl dimethicone
Disodium-PG-propyldimethicone thiosulfate
Isopropyl hydroxybutylamide dimethicone copolyol
Methicone

Smoother, Shinier Hair?
Try our cross-linking conditioner: **CRODASONE W**

Croda
7 Century Dr Parsippany NJ 07054 201-644-4900

BROOKS INDUSTRIES inc.

Cosmetic Ingredients & Ideas®

70 Tyler Place, South Plainfield, NJ 07080 USA
Tel 908-561-5200 Fax 908-561-9174

Proteins

Hydrocoll, Solu-Soy, Wheat-Pro

Functions

Octamethyl cyclotetrasiloxane
Phenyl methicone, P. trimethicone
Polyether trisiloxane
Polymethylsilsequioxane
Polysilicone-A
Quaternium-80
Silicone quaternium-1, -8
Sodium-PG-propyl thiosulfate dimethicone
Stearoxymethiconedimethicone copolymer
Trimethylsilylamindimethicone

Skin calming agent

Cornflower (*Centaurea cyanus*) extract
Fennel (*Foeniculum vulgare*) extract
Fenugreek extract
Linden (*Tilia cordata*) extract
Valerian (*Valeriana officinalis*) extract

Skin cleanser

Dog rose (*Rosa canina*) hips extract
Papaya (*Carica papaya*) extract
Peach (*Prunus persica*) extract
Rose (*Rosa multiflora*) extract
Willow (*Salix alba*) extract

Skin conditioner

Artemisia apiacea extract
Astrocaryum tucuma extract
Bactris gasipaes extract
Biotin
Bishydroxyethyl biscetyl malonamide
Bletia hyacinthina extract
Borage (*Borago officinalis*) seed oil
Borageamidopropyl phosphatidyl PG-dimonium chloride
Carboxysteine
Catalpa kaempferia extract
Coco phosphatidyl PG-dimonium chloride
Cocodimonium hydroxypropyl hydrolyzed keratin
Collagen amino acids
Cyclomethicone
Dimethicone, D. copolyl acetate
Emblica officinalis extract
Equisetum arvense extract
Ethyl ester of hydrolyzed animal protein
Evening primrose (*Oenothera biennis*) oil
Fomes fomentarius extract
Fomistopsis officinalis oil
Gelatin
Ginseng hydroxypropyltrimonium chloride
butylene glycol
Glycolipids
Glycosphingolipids
Gnetum amazonicum extract
Honey (Mel)
Hydrolyzed carbolipoprotein
Hydrolyzed elastin
Hydrolyzed pea protein
Hydrolyzed rice protein
Hydrolyzed serum protein
Hydrolyzed silk
Hydrolyzed soy protein
Hydrolyzed vegetable protein
Hydrolyzed wheat protein
Inga edulis extract
Kiwi (*Actinidia chinensis*) fruit extract
Laminaria japonica extract
Lecithin
Marsilea minuta extract
Nettle (*Urtica dioica*) extract
Palmitamidodecanediol
Pearls (*Margarita margarita*)
PEG-42 Ebirko ceramides extract
Phenyl trimethicone
Phytantriol
Polygonum multiflorum extract
Polyquaternium-7, -22, -30
Polysiloxane, dimethylsiloxane

Potassium cocoyl hydrolyzed collagen
Retinyl palmitate polypeptide
Salvia miltiorrhiza extract
Silk
Sodium cocoyl hydrolyzed collagen
Soluble transgenic elastin
Stearammonium hydroxyethyl hydrolyzed collagen
Sicaryl methicone

Skin healing

Calendula officinalis extract
Glycoproteins
Hydrocortyl (*Centella asiatica*) extract
Oat (*Avena sativa*) extract
Sandalwood (*Santalum album*) extract
Spearmint (*Mentha viridis*) extract

Skin lightening/whitening agent

Ascorbic acid polypeptide
Bearberry (*Arctostaphylos uva-ursi*) extract
Hydroquinone-beta-D-glucopyranoside
Lemon (*Citrus medica limonum*) peel extract
Pearls (*Margarita margarita*)

Skin protectant

Acetyl methionyl methylsilanol elastinate
Allantoin, A. aluminum hydroxide
Aloe barbadensis, A. b. extract
Aluminum starch octenylsuccinate
Anise (*Pimpinella anisum*) extract
Arnica montana extract
Artemisia apiacea extract
Ascorbyl methylsilanol pectinate
Astrocaryum tucuma extract
Bactris gasipaes extract
Betaglucon
Bishydroxyethyl biscetyl malonamide
Bletia hyacinthina extract
C 18-70 Isoparaffin
Calendula amurensis extract
Carboxymethyl chitin
Carcinia cambogia extract
Carrot (*Daucus carota*) extract
Carrot (*Daucus carota sativa*) oil
Catalpa kaempferia extract
Chenopodium album extract
Chitosan
Chrysanthemum morifolium extract
Collagen
Corn poppy (*Papaver rhoeas*) extract
Crataegus cuneata extract
Crataegus monogina extract
Cypress (*Cupressus sempervirens*) extract
Dimethicone
Dimethiconol fluoroalcohol dilinoleic acid
Dimethiconol hydroxystearate, D. stearate
Dimethylsilanol hyaluronate
Echitea glauca extract
Embryo extract
Entada phaseoloides extract
Equisetum arvense extract
Eupatorium fortunei extract
Euterpe precatoria extract
Fenugreek extract
Fomistopsis officinalis oil, F. pinicola extract
Galla sinensis extract
Gentian (*Gentiana lutea*) extract
Gleditsia sinensis extract
Glyceryl ricinoleate
Glycolipids
Hierochloa odorata extract
Hyaluronic acid
Hydrogenated lecithin
Hydrolyzed lupine protein
Hydrolyzed milk protein
Hydrolyzed mushroom (*Tricholoma matsutake*) extract
Indian cress (*Tropaeolum minus*) extract

Isodecyl salicylate
Jojoba (*Buxus chinensis*) oil
Lady's Thistle (*Silybum marianum*) extract
Laminaria japonica extract
Ligusticum jeholense extract
Liposomes
Magnolia spp. extract
Mango kernel oil
Marsilea minuta extract
Melaleuca hypericifolia extract
Melaleuca uncinata extract
Melaleuca wilsonii extract
Methylsilanol tri PEG-8 glyceryl cocoate
Oat (*Avena sativa*) meal
Oyster (*OSTREA*) shell extract
Palmitamidodecanediol
Pearls (*Margarita margarita*)
Pentahydrosqualene
Perfluorodecalin
Perfluoropolymethylisopropyl ether
Petrolatum
PEG-8/SMDI copolymer
PEG-42 Ebirko ceramides extract
Pfaffia spp. extract
Phospholipids
Plankton extract
Polygonum multiflorum extract
Pongamol
PPG-12/SMDI Copolymer
PPG-51/SMDI Copolymer
Propyltrimonium hydrolyzed collagen
Quinoa (*Chenopodium quinoa*) extract, oil
Salvia miltiorrhiza extract
Sambucus nigra extract
Shark liver oil
Shorea robusta extract
Sodium chondroitin sulfate
Soluble transgenic elastin
Stearammonium hydroxyethyl hydrolyzed collagen
Sterculia plataniifolia extract
Superoxide dismutase
Trachea hydrolysate
Wheat (*Triticum vulgare*) germ extract, protein
White nettle (*Lamium album*) extract
Withania somniferum extract
Xanthoxylum bungeanum extract
Zinc oxide

Skin smoothing agent

Althea officinalis extract
Coltsfoot (*Tussilago farfara*) leaf extract
Comfrey (*Symphytum officinale*) leaf extract
Plantain (*Plantago major*) extract
Sericin

Skin softening

Clays (white, yellow, red, green, pink)
Cucumber (*Cucumis sativus*) extract
Kelp (*Macrocystis pyrifera*) extract
Peach (*Prunus persica*) extract
Phenethyl dimethicone

Skin soothing

Calendula officinalis extract
Cherry bark extract
Cucumber (*Cucumis sativus*) extract
Garlic (*Allium sativum*) extract
Hyssop (*Hyssopus officinalis*) extract
Jasmine (*Jasminum officinale*) extract
Kelp (*Macrocystis pyrifera*) extract
Mango kernel oil
Meadowsweet (*Spiraea ulmaria*) extract
Quince (*Pyrus cydonia*) seed extract
Slippery elm extract
Valerian (*Valeriana officinalis*) extract
Willow (*Salix alba*) extract
Witch hazel (*Hamamelis virginiana*) extract
Yarrow (*Achillea millefolium*) extract

Functions

Solubilizer

Acetyl monoethanolamine
Almond oil PEG-6 esters
2-Aminobutanol
Aminoethyl propanediol
Aminoethyl propanediol, A. propanol
Apricot kernel oil PEG-6 esters
Benzalkonium chloride
Butoxydiglycol
Butyl glucoside
Butylene glycol
Butyloctanol
Capric-caprylic mono-diglyceride
Capryl caprylylglycoside
Caprylic/capric triglyceride
Caprylic/capric/linoleic triglyceride
Caprylic/capric/oleic triglycerides
Caprylyl/capryl glucoside
Ceteareth-20
Ceteth-10
Cetyl PPG-2 isodeceth-7 carboxylate
Cholesterol
Cotton oil PEG-6 esters
Decaglycerol monodistearate
Diethanolamine
Dilaureth-10 phosphate
Dimethyl octylnediol
Dioleth-8 phosphate
Glycereth-7 -26
Glyceryl caprylate, G. dilaurate
Glyceryl caprylate/caprate
Isocicosane
Isopropanolamine
Isosteareth-20
Laneth-5, -15
Laureth-23
Methylated cyclodextrin
Myreth-3
Myreth-3-octanoate
Nonoxynol-10, -12, -14, -40, -50
Octoxynol-11, -40
Oleamphosphorylpropylsulfonate
Oleth-3, -5, -10, -15, -20, -25, -50
Oleth-20 phosphate
PEG-4, -6, -8, -12, -16, -20, -32, -40,
PEG-4 dilaurate
PEG-6 capric/caprylic glycerides
PEG-6 methyl ether
PEG-8 distearate
PEG-12 laurate

PEG-15 castor oil
PEG-18 stearate
PEG-20 glyceryl isostearate, P. g. laurate
PEG-20 glyceryl oleate, P. g. stearate
PEG-20 methyl glucose sesquisteate
PEG-20 sorbitan isostearate
PEG-20 sorbitan triisostearate
PEG-24 hydrogenated lanolin
PEG-25 castor oil
PEG-25 hydrogenated castor oil
PEG-30 castor oil
PEG-30 glyceryl cocoate
PEG-30 glyceryl isostearate
PEG-30 glyceryl laurate
PEG-30 glyceryl oleate
PEG-30 glyceryl stearate
PEG-33 castor oil
PEG-35 castor oil
PEG-36 castor oil
PEG-40 castor oil
PEG-40 glyceryl laurate, P. g. stearate
PEG-40 hydrogenated castor oil
PEG-40 hydrogenated castor oil PCA isostearate
PEG-40 sorbitan diisostearate
PEG-45 palm kernel glycerides
PEG-48 hydrogenated castor oil
PEG-50 castor oil
PEG-50 hydrogenated castor oil
PEG-60 almond glycerides
PEG-60 castor oil
PEG-60 com glycerides
PEG-60 glyceryl isostearate, P. g. stearate
PEG-60 hydrogenated castor oil
PEG-60 lanolin
PEG-70 mango glycerides
PEG-75 lanolin
PEG-75 shea butter glycerides
PEG-75 shea butter glycerides
PEG-80 hydrogenated castor oil
PEG-80 jojoba acid/alcohol
PEG-80 sorbitan laurate
PEG-100 castor oil
PEG-100 hydrogenated castor oil
PEG-120 jojoba acid/alcohol
PEG-200 trihydroxystearin
Poloxamer 407
Polyglyceryl-3 oleate
Polyglyceryl-6 dioleate
Polyglyceryl-10 decaoleate, P. tetraoleate
Polysorbate 20, 60, 80
PPG-2-isodeceth-4, -6, -9, -12

PPG-3 isosteareth-9
PPG-3 isoceteth-20 acetate
PPG-3-ceteth-10 phosphate
PPG-5-ceteth-20
PPG-6-decyltetradeceth-12, -20, -30
PPG-12-PEG-65 lanolin oil
PPG-15 stearyl ether
PPG-18 butyl ether
PPG-24 butyl ether
PPG-26-buteth-26
PPG-33 butyl ether
PPG-33-buteth-45
PPG-40-PEG-60 lanolin oil
PPG-50 cetyl ether
Propylene glycol dicaprylate, dicaprylate/
dicaprate
Ricinoleamide DEA
Ricinoleth-40
Sodium alpha olefin sulfonate
Sodium lauryl sulfate
Sodium methylphenylsulfonate
Triethanolamine
Trioxanoin
Tromethamine

Solvent

Acetic acid
Acetone
Alcohol, A. denat.
Benzophenone
Butoxydiglycol
Butyl acetate
n-Butyl alcohol
Butyl myristate, B. stearate
Butylene glycol
C9-11 isoparaffin
C10-11 isoparaffin
C10-13 isoparaffin
Caprylic alcohol
Castor (Ricinus communis) oil
Cetearyl octanoate
Cetyl stearyl octanoate
Chlorobutanol
Decyl alcohol
Diethylene glycol
Diethylene glycol dibenzoate
Diethyl sebacate
Diisocetyl adipate
Diisopropyl adipate, D. sebacate
Dimethyl phthalate
Dipropylene glycol

Pigments • Preservatives • Surfactants • Emulsifiers • Emollients • Esters • Proteins • Lanolin



Costec

We Specialize in Personal Care.

Performance Chemicals for Specialty Requirements.

Costec, Inc. • 655 First Bank Drive • Palatine, IL 60067 USA
Telephone: 847/359-5713 • Fax: 800/5 COSTEC

Aloe Vera • Jojoba Oil • Botanicals • Pigments • Preservatives • Surfactants • Emulsifiers

Functions

Dipropylene glycol dibenzoate	Glyceryl diisostearate, G. stearate SE	Superfating agent
Ethoxydiglycol	Glyceryl mono-di-in-caprylate	Linoleamide DEA
Ethyl acetate, E. lactate	Hydrogenated coco-glycerides	PEG-20 almond glycerides
Ethyl myristate, E. oleate	Hydrogenated C12-18 triglycerides	PEG-60 lanolin
2-Ethylhexyl isostearate	Hydrogenated tallow glycerides	PEG-75 lanolin
Glycerin	Hydrolyzed oat flour	
Glycolfurfol	Hydroxyoctacosanyl hydroxystearate	
Heptane	Karaya (<i>Stemculia urens</i>) gum	Surfactant
Hexyl alcohol	Laureth-3	Alkyl dimethyl betaine
Hexylene glycol	Maltitol	Alkyldimethylamine oxide
Isobutyl stearate	Methylated cyclodextrin	Ammonium cocoyl sarcosinate
Isocetyl salicylate	Oleamide	Ammonium C12-15 alkyl sulfate
Isodecyl benzoate, I. isononanoate	PEG-40 stearate	Ammonium dimethicone copolyol sulfate
Isodecyl octanoate, I. oleate	PEG-40/dodecyl glycol copolymer	Ammonium laureth-5 sulfate
Isododecane	Perfluoropolyethylisopropyl ether	Ammonium laureth-12 sulfate
Isocicosane	Polyethylene paste	Ammonium laureth sulfate
Isohexadecane	PPG-5 lanolin wax	Ammonium lauroyl sarcosinate
Isopropyl alcohol, I. myristate	PPG-7-buteth-10	Ammonium lauryl sulfate, A. I. sulfosuccinate
Isostearyl stearoyl stearate	PPG-10 cetyl ether phosphate	Ammonium myreth sulfate
Laureth-2 acetate	Propylene carbonate, P. glycol alginate	Ammonium nonoxynol 4 sulfate
Methoxydiglycol	PVM/MA decadiene crosspolymer	Azelamide MEA
Methoxyisopropanol	Sodium acrylates/vinyl isodecanoate crosspolymer	C20-40 alcohol ethoxylate
Methyl alcohol	Sodium carbomer	C30-50 alcohol ethoxylate
Methyl propanediol	Sorbitan laurate	C40-60 alcohol ethoxylate
Methylene chloride	Stearic hydrazide	Calcium dodecylbenzene sulfonate
MEK	2,2',4,4'-Tetrahydroxybenzophenone	Calcium laurate
MIBK	Tricaprin	Ceteareth-2 phosphate
Morpholine	Tricaprylin	Ceteareth-5 phosphate
Octyl benzoate, O. isononanoate	Triaurin	Ceteareth-10 phosphate
Octyl laurate, O. palmitate	Trimyristin	Cetoleth-25
Octyldodecyl lactate	Tripalmitin	Cetyl betaine, C. phosphate
Olive oil PEG-6 esters	Tristearin	Cocamide MEA ethoxylate
Peanut oil PEG-6 esters		Cocamidopropyl betaine, potassium salt
Pentane	Stimulant	Cocamidopropyl betaine ammonium salt
Petroleum distillates	Capiscum frutescens extract	Cocamidopropyl hydroxy sulfate
PEG-6 methyl ether	Eleuthero ginseng (<i>Acanthopanax senticosus</i>) extract	Cocamidopropyl hydroxy sulfate, ammonium salt
PEG-12	Guarana (<i>Paullinia cupana</i>) extract	Cocamidopropyl hydroxy sulfate, potassium salt
PEG-20 hydrogenated castor oil	Lactococcus hydrolysate	Cocamidopropylamine oxide
PEG-33 castor oil	Methylsilanol elastinate	Coceth-7 carboxylic acid
PEG-50 glyceryl cocoate	Methylsilanol hydroxyproline aspartate	Coco-glucoside
Polyglyceryl-2 dioleate	TEA-hydroiodide	Cocamphodiacetate lauryl-laureth sulfate
Polyglyceryl-3 diisostearate	Tocopheryl nicotinate	Cocamphodiacetate lauryl sulfate
Polyoxyethylene glycol dibenzoate	Urocanic acid	Cocamphodiacetate trideceth sulfate
Polypropylene glycol dibenzoate	Yeast (<i>Saccharomyces cerevisiae</i>) extract (Faex)	Coco phosphatidyl PG-dimonium chloride
PPG-2 myristyl ether propionate	Zedoary (<i>Curcuma zedoaria</i>) oil	N-Cocoyl-(3-amidopropyl)-N,N-dimethyl-N-ethyl ammonium ethyl sulfate
PPG-3	Zinc DNA	Cocoyl glutamic acid
PPG-20 lanolin alcohol ether		Cocoyl hydrolyzed soy protein
Propyl alcohol	Sunscreen	Cocoyl hydroxyethyl imidazoline
Propylene carbonate	Basil (<i>Basilicum santum</i>) oil extract	C11-15 pareth-9, -12, -20, -30, -40
Propylene glycol	Basil (<i>Ocimum basilicum</i>) extract	C12-13 pareth sulfate
Propylene glycol dibenzoate	Benzophenone-3 -4	C12-13 pareth-5 carboxylic acid
Propylene glycol methyl ether	3-Benzylidene camphor	C12-15 pareth-12
Propylene glycol myristate	Borjoia sorbilis extract	C14-15 pareth-8 carboxylic acid
Pyridine	C12-15 alkyl benzoate	DEA-oleth-5-phosphate
Sesame (<i>Sesamum indicum</i>) oil	Coffee (<i>Coffea arabica</i>) bean extract	DEA-oleth-20-phosphate
Stearyl heptanoate	Ethyl salicylate	Deceth-3, -6, -8
Toluene	Glyceryl PABA	Decyltetradeceth-25
Xylene	Homosalate	Diceteareth-10 phosphoric acid
SPF booster	Hydroquinone-beta-D-glucopyranoside	Dimethicone copolyol
Borjoia sorbilis extract	Isoamyl p-methoxycinnamate	Dimethicone copolyol almondate, D. c. isostearate
Isohexadecyl salicylate	Isopropylbenzyl salicylate	Dimethicone copolyol laurate, D. c. olivate
Styrene/acrylates copolymer	Job's tears (<i>Coix lacryma-jobi</i>) extract	Dimethicone copolyol phthalate
Titanium dioxide	Menthyl anthranilate	Dimethicone copolyolamine
Yeast (<i>Saccharomyces cerevisiae</i>) extract (Faex)	Octyl dimethyl PABA, O. methoxycinnamate	Dimethicone propyl PG-betaine
Stabilizer	Octyl salicylate, O. triazone	Diocryldodeceth-2 lauroyl glutamate
Acrylates-VA crosspolymer	Oryzanol	Diocryldodeceth-5 lauroyl glutamate
Acrylates/ceteeth-20 methacrylates copolymer	Pansy (<i>Viola tricolor</i>) extract	Diocryldodecyl lauroyl glutamate
Acrylates/steareth-20 methacrylate copolymer	PEG-25 PABA	Disodium capryloamphodiacetate
Acrylates/vinyl isodecanoate crosspolymer	Phenylbenzimidazole sulfonic acid	Disodium cocoamphodiacetate
Alkyldimethylamine oxide	Rice (<i>Oryza sativa</i>) bran oil	Disodium hydrogenated tallow glutamate
C10 polycarbonyl polyglycol ester	TEA-salicylate	Disodium laneth-5 sulfosuccinate
Calcium alginate	Titanium dioxide	Disodium lauramido MEA-sulfosuccinate
Cocamidopropyl dimethylamine lactate	Sunscreen UVB	Disodium laureth sulfosuccinate
Cocamine oxide	Benzophenone-5	Disodium oleamido MIPA-sulfosuccinate
Colloidal silica sols	Eclipta alba extract	Disodium oleamido PEG-2 sulfosuccinate
Cyclodextrin	PEG-25 PABA	Disodium oleth-3 sulfosuccinate
Disodium EDTA	Steareth-100	Disodium ricinoleamido MEA-sulfosuccinate
Gellan gum	Tridecyl salicylate	Disodium tallamido MEA-sulfosuccinate
		Disteareth-2 lauroyl glutamate

Functions

Disteareth-5 lauryl glutamate
 Ethoxylated fatty alcohol
 Ethoxylated glycerol sorbitan saturated fatty acid ester
 Ethoxylated glycerol sorbitan unsaturated fatty acid ester
 Glycereth-25 PCA isostearate
 Glycereth-26 phosphate
 Glyceryl hydroxystearate
 Hydrogenated tallowyl glutamic acid
 Isopropyl hydroxybutylamide dimethicone copolyol
 Lauramidopropyl betaine
 Laureth-1, -2, -3, -4, -7, -12, -16
 Laureth-3 carboxylic acid, L. phosphate
 Laureth-5 carboxylic acid
 Laureth-11 carboxylic acid
 Lauroyl sarcosine
 Lauryl dimethylamine cyclocarboxypropyloleate
 Lauryl hydroxyethyl imidazoline
 Linoleamide DEA
 Magnesium laureth-8 sulfate
 Meroxapol 105, 171, 172
 MEA-lauryl sulfate
 Mixed isopropanolamines myristate
 Myreth-7
 Myristoyl sarcosine
 Myristyl alcohol
 Nonoxynol-7, -9, -13, -15
 Nonoxynol-10 carboxylic acid
 Octoxynol-10, -12
 Octyldodeceth-10, -16
 Oleoyl sarcosine
 Oleth-2 phosphate
 Oleth-5 phosphate
 Oleyl betaine
 Oleyl hydroxyethyl imidazoline
 Palmitamine oxide
 Palmityl betaine
 PCA ethyl cocoyl arginate
 PEG-7 hydrogenated castor oil
 PEG-8 caprylic/capric glycerides
 PEG-8 laurate
 PEG-8 stearate
 PEG-15 glyceryl stearate
 PEG-25 glyceryl isostearate
 PEG-27 lanolin
 PEG-30 lanolin
 PEG-40 castor oil
 PEG-40 glyceryl stearate
 PEG-40 jojoba oil, P. lanolin
 PEG-60 glyceryl isostearate, P. g. stearate

PEG-80 jojoba oil, P. sorbitan laurate
 PEG-120 jojoba oil
 Pentasodium triphosphate
 Poloxamer 101, 122
 Polyglyceryl-2 dioleate
 Polysiloxane-polyether copolymer
 Potassium cocoyl glycinate
 Potassium cocoyl hydrolyzed collagen
 Potassium C9-15 phosphate ester
 Potassium lauroyl hydrolyzed collagen
 Potassium lauryl sulfate
 Potassium myristoyl hydrolyzed collagen
 Potassium oleyl hydrolyzed collagen
 Potassium palmitate
 Potassium undecylenoyl hydrolyzed collagen
 PPG-2-isodeceth-4, -6, -9, -12
 PPG-6 C12-18 pareth-11
 Protein hydrolysates
 Quaternium-80
 Quillaja saponaria extract
 Raffinose laurate, R. myristate, R. oleate
 Raffinose palmitate, R. stearate
 Ricinoleamidopropyl betaine
 Silicone quaternium-1, -8, -9
 Sodium alpha olefin sulfonate
 Sodium cocoumphoacetate
 Sodium cocoyl hydrolyzed wheat protein
 Sodium cocoyl isethionate
 Sodium C12-13 sulfate
 Sodium C12-14 pareth-2 sulfate
 Sodium C12-15 pareth-3 sulfonate
 Sodium C12-15 pareth-7 carboxylate
 Sodium C12-15 pareth-7 sulfonate
 Sodium C12-15 pareth-8 carboxylate
 Sodium C12-15 pareth-15 sulfonate
 Sodium C12-18 alkyl sulfate
 Sodium C13-17 alkane sulfonate
 Sodium C14-16 olefin sulfonate
 Sodium cetearyl sulfate
 Sodium cetyl oleyl sulfate
 Sodium coco-tallow sulfate
 Sodium cocoyl glutamate
 Sodium cocoyl hydrolyzed collagen
 Sodium cocoyl hydrolyzed soy protein
 Sodium cocoyl sarcosinate
 Sodium dimethicone copolyol aceryl methyltaurate
 Sodium hydrogenated tallow glutamate
 Sodium isodecyl sulfate
 Sodium laureth-5 carboxylate
 Sodium laureth-11 carboxylate
 Sodium laureth-13 carboxylate
 Sodium laureth sulfate
 Sodium lauroamphoacetate

Sodium lauroyl glutamate
 Sodium lauroyl hydrolyzed collagen
 Sodium lauroyl sarcosinate, S. l. taurate
 Sodium magnesium laureth sulfate
 Sodium methyl cocoyl taurate
 Sodium methyl oleoyl taurate
 Sodium myristoyl glutamate
 Sodium myristoyl hydrolyzed collagen
 Sodium myristoyl sarcosinate
 Sodium myristyl sulfate
 Sodium nonoxynol-6 phosphate
 Sodium octoxynol-2 ethane sulfonate
 Sodium octyl sulfate
 Sodium oleyl hydrolyzed collagen
 Sodium stearoyl hydrolyzed collagen
 Sodium undeceth sulfate
 Sodium undecylenoyl hydrolyzed collagen
 Sodium/TEA-lauroyl hydrolyzed collagen
 Sodium/TEA-lauroyl hydrolyzed keratin
 Sorbitan isostearate
 Stearoyl sarcosine
 Sulfated castor oil
 TEA-cocoyl glutamate
 TEA-cocoyl hydrolyzed collagen
 TEA-cocoyl hydrolyzed soy protein
 TEA-C12-15 alkyl sulfate
 TEA-hydrogenated tallow glutamate
 TEA-lauroyl glutamate
 TEA-lauroyl keratin amino acids
 TEA-lauroyl sarcosinate
 TEA-lauryl sulfate
 TEA-myristoyl hydrolyzed collagen
 Tocophereth-5 -10 -18 -20 -30 -50 -70
 Trideceth-7 carboxylic acid
 Trideceth-9
 Trideceth-19-carboxylic acid
 Tridecyl ethoxylate
 Triethanolamine C10-14 sulfate
 Trilauryl phosphate
 Wheat germamidopropyl betaine
 Yucca vera extract

Suspending agent

Acrylates/ceteeth-20 methacrylates copolymer
 Acrylates/steareth-20 methacrylate copolymer
 Algin
 Bentonite
 C10 polycarbamyl polyglycol ester
 Calcium alginate
 Carbomer, C. 934
 Carrageenan (Chondrus crispus)
 Cellulose gum
 Cetyl hydroxyethylcellulose

Witco

Witco Corporation
 Oleo/Surfactants Group
 Call 800 779-4826
 203 881-6277 (outside the U.S.)
 Fax 203 552-2850


Meeting daily your changing
 needs for surfactants globally



McINTYRE
 GROUP LTD.

OUR NAME IS MCINTYRE
 OUR GAME IS AMPHOTERICS

24601 Governors Highway • University Park, IL 60466
 (708) 534-6200 • FAX: (708) 534-6216 • 1-800-645-6457



Functions

Dihydroxycatenated tallow phthalic acid amide
Distearyl phthalic acid amide
Guar (Cyanopsis tetragonoloba) gum
Hectonite
Hydroxypropylcellulose
Isobutylene/MA copolymer
Magnesium aluminum silicate
Methylcellulose
Pentasodium triphosphate
Polyethylene, P. micronized
Propylene glycol alginate
Quaternium-18 bentonite
Quaternium-18 hectorite
Sodium magnesium silicate
Sodium polynaphthalenesulfonate
Stearalkonium bentonite, S. hectorite
Styrene-10 allyl ether/acrylate copolymer
Tragacanth (Astragalus gummi) gum
Tribehenol
Trihydroxystearin
Trimethamine magnesium aluminum silicate
Xanthan gum

Sweetener

Calcium saccharin
Fructose
Glycyrrhizic acid
Glycyrrhizic acid
Glycyrrhizin, ammoniated
Hydrolyzed corn starch
Lactose
Maltitol
Mannitol
Saccharin
Sodium saccharin
Sorbitol
Sucrose

Tanning accelerator

Acetyl tyrosine
Carrot (Daucus carota) extract
Copper acetyl tyrosinate methylsilanol
Dihydroxyacetone
Disodium methyl tyrosinate
Eclipta alba extract in white emulsion
Glucose tyrosinate

Thickener

Acrylates-VA crosspolymer
Acrylates/C10-C30 alkyl acrylate crosspolymer
Acrylates/ceteth-20 itaconate copolymer
Acrylates/ceteth-20 methacrylates copolymer
Acrylates/steareth-20 itaconate copolymer
Acrylates/steareth-20 methacrylate copolymer
Acrylates/steareth-50 acrylate copolymer
Acrylates/vinyl isodecanoate crosspolymer
Acrylic acid/acrylonitrile copolymer
Algin
Aluminum/magnesium hydroxide stearate
Ammonium acrylates/acrylonitrile copolymer
Ammonium alginate
Arachidyl alcohol
Behenic acid
Behenyl alcohol, B. behenate
Bentonite
C10 polyacrylamyl polyglycol ester
C12-15 alcohols
C12-16 alcohols
C18-36 acid

Calcium alginate
Calcium carrageenan
Caprylic alcohol
Carbomer
Carboxymethyl hydroxyethylcellulose
Carrageenan (Chondrus crispus)
Cellulose, C. gum
Cetearyl alcohol, C. behenate
Cetearyl octanoate, C. stearate
Cetostearyl stearate
Cetyl alcohol
Cetyl hydroxyethylcellulose
Cetyl myristate, C. palmitate
Cocamide
Cocamide MEA, C. MIPA
Cocamidopropylamine oxide
Coco-betaine
Coco-rapeseedate
Coco/oleamidopropyl betaine
Cocoyl amido hydroxy sulfo betaine
Cocoyl monoethanolamide ethoxylate
Colloidal silica sols
DEA-hydrolyzed lecithin
DEA-linoleate
DEA-oleth-3 phosphate
DEA-oleth-10 phosphate
Decyl alcohol
Dextran
Dextrin
Dilaureth-10 phosphate
Dioleth-8 phosphate
DNHF
Ethoxylated fatty alcohol
Gellan gum
Glycerol behenate, G. stearate
Glycerol polymethacrylate
Guar (Cyanopsis tetragonoloba) gum
Guar hydroxypropyltrimonium chloride
Hectonite
Hexyl alcohol
Hydrated silica
Hydrogenated rapeseed oil
Hydrogenated starch hydrolysate
Hydrogenated talloweth-60 myristyl glycol
Hydrolyzed oat flour
Hydrolyzed transgenic collagen
Hydroxyethylcellulose
Hydroxypropyl chitosan
Hydroxypropyl guar
Hydroxypropyl methylcellulose
Hydroxypropylcellulose
Isoceteth-10
Isocetamide DEA
Isocetamidopropylamine oxide
Isocetamphopropionate
Jojoba wax
Karaya (Sterculia urens) gum
Lecithide DEA, L. MEA, L. MIPA
Lecithidopropyl betaine
Lecith-10
Lecith-linoleic DEA
Lecith-linoleoyl diethanolamide
Lecith-myristoyl diethanolamide
Lecithyl alcohol, L. betaine
Lecithamide DEA, L. MEA
Lecithic acid
Lecithic acid
Lecithic acid
Lecithic bean (Ceratonia siliqua) gum
Magnesium aluminum silicate

MDM hydantoin
Methylcellulose
Montmorillonite
Myristamide DEA, M. MEA
Myristamine oxide
Myristyl alcohol
Octacosanyl stearate
Oleamide, O. DEA, O. MEA
Palmitamide MEA
Pectin
PEG-2 laurate
PEG-3 distearate, P. lauramide
PEG-3 lauramine oxide
PEG-4 diisostearate, P. oleamide
PEG-5M
PEG-6 beeswax
PEG-7 hydrogenated castor oil
PEG-8
PEG-8 dioleate, P. distearate
PEG-8 stearate
PEG-9M
PEG-12 beeswax
PEG-18 glyceryl oleate/cocaoate
PEG-23M
PEG-28 glyceryl tallowate
PEG-40 jojoba oil
PEG-45M
PEG-50 tallow amide
PEG-55 propylene glycol oleate
PEG-75 stearate
PEG-90M
PEG-100 stearate
PEG-120 methyl glucose dioleate
PEG-150 distearate
PEG-150 pentaerythrityl tetrastearate
PEG-160M
PEG-200 glyceryl stearate
PEG-200 glyceryl tallowate
Pentaerythrityl tetrabehenate
Pentaerythrityl tetrastearate
Poloxamer 105, 124, 185, 237, 238, 338, 407
Polyacrylic acid
Polysorbate 20
Potassium alginate, P. chloride
Potassium oleate, P. stearate
PPG-5-ceteth-10 phosphate
Propylene glycol stearate
PVM/MA decadiene crosspolymer
PVP
Quaternium-18 bentonite
Quaternium-18 hectorite
Rapeseed oil, ethoxylated high erucic acid
Ricinoamide MEA
Sesamide DEA
Sodium acrylates/vinyl isodecanoate crosspolymer
Sodium carbomer, S. carrageenan
Sodium ceteth-13-carboxylate
Sodium chloride
Sodium magnesium silicate, S. stearate
Sorbitan sesquiosostearate, S. tristearate
Soyamide DEA
Soyamidopropyl betaine
Starch polyacrylonitrile copolymer-potassium salt
Starch polyacrylonitrile copolymer-sodium salt
Stearalkonium bentonite, S. hectorite
Stearamide
Stearamide DEA, S. MEA, S. MEA-stearate
Stearamidopropyl dimethylamine lactate
Stearamine oxide

3 BETTER IDEAS.**1 BETTER SOURCE.**

CARBOPOL
Ultraz
New, easiest to
use carbomer

CARBOPOL
HTD
POLYMER
For surfactant-based
emulsifiers

PEMULEN
POLYMERIC EMULSIFIERS
Eliminates surfactant-based
emulsifiers

BF Goodrich
Talk to the global leader.

Functions

Steareth-10 allyl ether/acrylates copolymer
Stearic acid
Stearyl alcohol
Synthetic beeswax
Tallowamide MEA
TEA-acrylates/acrylonitrile copolymer
Tragacanth (Astragalus gummifer) gum
Tribehenin
Trihydroxysteann
Tromethamine mordenium aluminum silicate
Wheat germamide DEA
Wheat germamidopropyl betaine
Xanthan gum

Thixotropes

Bentonite
Hectonite
Sodium magnesium silicate
Stearalkonium bentonite

Toner

Althea officinalis extract
Clover (Trifolium pratense) extract
Dog rose (Rosa canina) hips extract
Ginseng (Panax ginseng) extract
Horsetail extract
Lemon bioflavonoids extract
Meadowsweet (Spiraea ulmaria) extract
Nettle (Urtica dioica) extract
Rose (Rosa multiflora) extract
Rosemary (Rosmarinus officinalis) extract

UVA absorber

Benzophenone-1, -2, -3, -4, -6, -8, -9, -11, -12
Butyl methoxydibenzoylmethane
Corallina officinalis
Isopropyl dibenzoylmethane
Menthyl anthranilate
2,2',4,4'-Tetrahydroxybenzophenone
Titanium dioxide
Zinc oxide

UVB absorber

Argania spinosa oil
Benzophenone-1, -2, -3, -4, -6, -9, -11
Corallina officinalis
DEA-methoxycinnamate
Drometrizole
Ethyl dihydroxypropyl PABA
Eucrylene
Homosalate
Isoamyl p-methoxycinnamate
Isopropyl methoxycinnamate
Isopropylbenzyl salicylate
4-Methylbenzylidene camphor
Octocrylene
Octrizole
Octyl dimethyl PABA
Octyl methoxycinnamate
Octyl salicylate, O, triazone
PABA
PEG-25 PABA
Phenylbenzimidazole sulfonic acid
Shea butter, ethoxylated
TEA-salicylate
Titanium dioxide
TriPABA panthenol
Zinc oxide

Vegetable oil

Apricot (Prunus armeniaca) kernel oil
Avocado (Persea gratissima) oil
Baobab oil
Calendula officinalis oil
Chaulmoogra (Tarakogenos kurzii) oil
Coconut (Cocos nucifera) oil
Corn (Zea mays) oil
Cottonseed (Gossypium) oil

Gold of pleasure oil
Grape (Vitis vinifera) seed oil
Hazel (Corylus avellana) nut oil
Hybrid sunflower (Helianthus annuus) oil
Hydrogenated coconut oil
Hydrogenated cottonseed oil
Hydrogenated vegetable oil
Jojoba (Buxus chinensis) oil
Kukui (Alcurites moluccana) nut oil
Macadamia ternifolia nut oil
Meadowfoam (Limnanthes alba) seed oil
Mexican puppy oil
Palm (Elaeis guineensis) kernel oil
Partially hydrogenated soybean oil
Peach (Prunus persica) kernel oil
Peanut (Arachis hypogaea) oil
Pecan (Carya illinoensis) oil
Pumpkin (Cucurbita pepo) seed oil
Quinoa (Chenopodium quinoa) oil
Rapeseed (Brassica campestris) oil
Rice (Oryza sativa) bran oil
Safflower (Carthamus tinctorius) oil
Seabuckthorn oil
Sesame (Sesamum indicum) oil
Sisymbrium irio oil
Soybean (Glycine soja) oil
Sunflower (Helianthus annuus) seed oil
Walnut (Juglans regia) oil
Wheat (Triticum vulgare) germ oil
Wild borage oil

Vitamin

Aesculus chinensis extract
Ascorbic acid
Ascorbic acid polypeptide
Ascorbyl palmitate
Biotin
Calcium pantothenate
Cholecalciferol
Cyanocobalamin
Eclipta alba extract
Embilca officinalis extract
Equisetum arvense extract
Ergocalciferol
Esculin
Ethyl linoleate
Folic acid
Laminaria japonica extract
Marsilea minuta extract
Melaleuca bracteata extract
Menadione
Nasturtium sinensis extract
Nelumbium speciosum extract
Niacin
Niacinamide, N, ascorbate
Nicotinamide
Nicotinic acid
Ocimum basilicum extract
Panthenyl triacetate
Pantothenic acid
Phytonadione
Pyridoxine HCL
Retinol
Retinyl acetate, R, palmitate
Retinyl palmitate polypeptide
Retinyl propionate
Riboflavin tetraacetate
Sodium ascorbate
Thiamine HCL
Tocopherol
Tocopheryl acetate, T, succinate

Wax

Bayberry (Myrica cerifera) wax
Behenoxy dimethicone
C16-18 alkyl methicone
Candelilla (Euphorbia cerifera) wax
Carnauba (Copernicia cerifera) wax

Ceresin
Cetyl dimethicone, C, isoctanoate
Diaktyldimethylpolysiloxane
Dimethiconol hydroxystearate
Dimethiconol stearate
Hydrogenated castor oil
Hydrogenated cottonseed oil
Hydrogenated jojoba oil, H, J, wax
Hydrogenated palm kernel oil
Hydrogenated rapeseed oil
Hydrogenated rice bran wax
Hydrogenated vegetable oil
Isocetadecyl isononanoate
Japan (Rhus succedanea) wax
Jojoba esters
Montan (Montan cera) wax
Oiticury wax
Ozokerite
Polyglyceryl-3 beeswax
Spermaceti
Stearoxymethicone/dimethicone copolymer
Stearoxymethylsilane
Synthetic candelilla wax
Synthetic carnauba

Wetting agent

Benzalkonium chloride
Benzethonium chloride
Cetalkonium chloride
Ceteareth-20
Ceteth-20
Cetyl pyridinium chloride
Cocoamphodipropionic acid
Decaglycerol monodiolate
Deceth-9
Dihydroabietyl methacrylate
Dimethicone copolyol methyl ether
Dimethicone copolyol phthalate
Diethyl sodium sulfosuccinate
Ethyl hydroxymethyl oleyl oxazoline
Hydroxylated milk glycerides
Isolaureth-6
Lanolin acid
Lauryl pyrrolidone
Lecithin
Methyl hydrogenated rosinate
Methyl rosinate
Nonyl nonoxynol-5
Octoxynol-8, 70
Oleth-15
Oleth-20 phosphate
PEG-9 castor oil
PEG-15 castor oil
PEG-20 glyceryl stearate
PEG-20 sorbitan trisostearate
PEG-45 palm kernel glycerides
PEG-60 almond glycerides, P, corn glycerides
PEG-60 shea butter glycerides
PEG-70 mango glycerides
PEG-75 shorea butter glycerides
PEG-80 sorbitan laurate
Poloxamer 123, 181, 182, 184, 235, 334
Polyether trisiloxane
Polyglyceryl-3 oleate
Polyglyceryl-6 dioleate
Polyglyceryl-10 tetraoleate
Polysorbate 60, 80
PPG-2-isodeceth-4, -6, -9, -12
PPG-10 lanolin alcohol ether
Propylene glycol
Sodium butoxyethoxy acetate
Sodium capryloamphohydroxypropylsulfonate
Sodium decyl diphenyl ether sulfonate
Sodium dodecyl diphenyl ether sulfonate
Sodium lauryl sulfate
Sulfated castor oil
Trisocetyl citrate
Trisostearin PEG-6 esters
Yucca vera extract

Claims:

1. A cosmetic composition, comprising:

a cosmetically acceptable carrier, comprising a reverse thermal viscosifying polymer network comprising at least one poloxamer component randomly bonded to at least one poly(acrylic acid) component said polymer network capable of aggregation
5 in response to a change in temperature; and

a cosmetically active agent which imparts a preselected cosmetic effect, said carrier and said agent disposed within an aqueous-based medium.

2. A cosmetic composition for topical application, comprising:

10 a cosmetically acceptable carrier, comprising a reverse thermal viscosifying polymer network comprising at least one poloxamer component capable of aggregation in response to a change in temperature randomly bonded to at least one poly(acrylic acid) component; and

15 a cosmetically active agent selected to treat imperfections or disorders of the skin, said carrier and said agent disposed within an aqueous-based medium.

3. The cosmetic composition of claim 1, wherein the cosmetic composition is a shampoo and the cosmetically active agent comprises a cleansing surfactant.

20 4. The cosmetic composition of claim 1, wherein the cosmetic composition is a moisturizer and the cosmetically active agent comprises a moisturizer.

25 5. The cosmetic composition of claim 1, wherein the cosmetic composition is a sunscreen and the cosmetically active agent comprises a uv-absorbing agent.

6. The cosmetic composition of claim 1, wherein the cosmetic composition is an acne cream and the cosmetically active agent comprises an antiacne agent.

5 7. The cosmetic composition of claim 1, wherein the cosmetic composition is a hair straightener and the cosmetic agent comprises a base for increasing the pH.

10 8. The cosmetic composition of claim 1, wherein the cosmetic composition is a sunless tanning lotion and the cosmetically active agent comprises skin tinting agent.

15 9. The cosmetic composition of claim 1, wherein the cosmetic composition is an antiperspirant and the cosmetically active agent comprises aluminum chlorhydrate.

20 10. The cosmetic composition of claim 1, wherein the cosmetic composition is a shaving cream and the cosmetically active agent comprises an emollient and a foaming surfactant.

11. The cosmetic composition of claim 1, wherein the cosmetic composition is a face cosmetic and the cosmetically active agent comprises a pigment.

25 12. The cosmetic composition of claim 1 or 2, wherein the cosmetic agent comprises a hydrophobic material, wherein the cosmetically acceptable carrier stabilizes the hydrophobic material in the aqueous medium.

13. The cosmetic composition of claim 2, wherein said cosmetic agent selected to treat imperfections or disorders of the skin is selected from the group consisting of acidulents, antiacne agents, anti-aging agents, anti-inflammatories, anti-irritants, antioxidants, depilatories, detergents, disinfectants, emollients, exfoliants, humectants, lubricants, moisturizers, skin conditioners, skin protectants, skin lightening agents, skin soothing agents, sunscreens, tanning accelerators and mixtures thereof.

14. The composition of claim 4, wherein said composition further comprises a cosmetic agent selected from the group consisting of humectants and emollients.

15. The composition of claim 1 or 2, further comprising one or more additives selected from the group consisting of preservatives, abrasives, acidulents, antiacne agents, anti-aging agents, antibacterials, anticaking, anticaries agents, anticellulites, antidandruff, antifungal, anti-inflammatories, anti-irritants, antimicrobials, antioxidants, astringents, antiperspirants, antiseptics, antistatic agents, antrngents, binders, buffers, additional carriers, chelators, cell stimulants, cleansing agents, conditioners, deodorants, depilatories, detergents, dispersants, emollients, emulsifiers, enzymes, essential oils, exfoliants, fibers, film forming agents, fixatives, foaming agents, foam stabilizers, foam boosters, fungicides, gellants, glosser, hair conditioner, hair set resins, hair sheen agents, hair waving agents, humectants, lubricants, moisture barrier agents, moisturizers, ointment bases, opacifier, plasticizer, polish, polymers, powders, propellant, protein, refatting agents, sequestrant, silicones, skin calming agents, skin cleansers, skin conditioners, skin healing, skin lightening agents, skin protectants, skin smoothing agents, skin softening agents, skin soothing agents, stabilizers, sunscreen agents, surfactants, suspending agents, tanning accelerators, thickeners, vitamins, waxes, wetting agents, liquefiers, colors, flavors and/or fragrances

16. The composition of claim 1, wherein the cosmetic composition takes a form selected from the group consisting of lotions, creams, sticks, roll-on formulations, mousses, sprays, aerosols, pad-applied formulations and masks.

5 17. The composition of claim 1, wherein the viscosification occurs at a temperature in the range of about 27 to 40°C.

18. The composition of claim 1, wherein the viscosification occurs at a temperature in the range of about 30 to 37°C.

10 19. The composition of claim 1, wherein said composition is formulated as a product selected from the group consisting of baby products, baby shampoos, lotions, powders and creams; bath preparations, bath oils, tablets and salts, bubble baths, bath fragrances bath capsules; eye makeup preparations, eyebrow pencil,
15 eyeliner, eye shadow, eye lotion, eye makeup remover, mascara; fragrance preparations, colognes, toilet waters, powders and sachets; noncoloring hair preparations, hair conditioner, hair spray, hair straighteners, permanent waves, rinses, shampoos, tonics, dressings and other grooming aids; color cosmetics: hair coloring preparations, hair dye, hair tints, hair color sprays, hair lighteners and hair bleaches;
20 makeup preparations, face powders, foundations, leg and body paints, lipstick makeup bases, rouges and makeup fixatives; manicuring preparations, basecoats, undercoats, cuticle softeners, nail creams, nail extenders, nail polish and enamel, and remover; oral hygiene products, dentrifices, mouthwashes; personal cleanliness, bath soaps, detergents, deodorants, douches and feminine hygiene product; shaving preparations,
25 aftershave lotion, beard softeners, men's talcum, shaving cream, shaving soap, preshave lotions; skin care preparations, skin cleansing preparations, skin antiseptics, depilatories, face and neck cleansers, body and hand cleansers, foot powders; moisturizers, night preparations, paste masks, skin fresheners; and suntan preparations, suntan creams, gels and lotions, and indoor tanning preparations.

20. The cosmetic composition of claim 1 or 2, wherein the poloxamer component is present in an amount in the range of about 0.01 to 20 wt% and the poly(acrylic acid component) is present in the amount of about 0.01 to 20 wt%.

5 21. The cosmetic composition of claim 1, wherein the polymer network comprises a plurality of poloxamers.

22. The cosmetic composition of claim 1, wherein the polymer network comprises a plurality of poloxamer components randomly bonded to a poly(acrylic acid) backbone.
10

23. The cosmetic composition of claim 1, wherein the reversibly viscosifying polymer composition comprises a plurality of poly(acrylic acid) components randomly bonded to a poloxamer component.
15

24. The cosmetic composition of claim 1, wherein the aqueous-based medium is selected from the group consisting of water, salt solutions and water with water-miscible organic compound(s).

20 25. The cosmetic composition of claim 1, further comprising an additive selected to increase transition temperature and increase viscosity of the reversible viscosifying polymer network.

25 26. The cosmetic composition of claim 1, further comprising an additive selected to increase transition temperature and decrease viscosity of the reversible viscosifying polymer network.

27. The cosmetic composition of claim 1, further comprising an additive selected to increase transition temperature without affecting viscosity of the reversible viscosifying polymer network..

5 28. The cosmetic composition of claim 1, further comprising an additive selected to decrease transition temperature and increase viscosity of the reversible viscosifying polymer network.

10 29. The cosmetic composition of claim 1, further comprising an additive selected to decrease transition temperature and decrease viscosity of the reversible viscosifying polymer network.

15 30. The cosmetic composition of claim 1, further comprising an additive selected to decrease transition temperature without affecting viscosity of the reversible viscosifying polymer network.

20 31. The cosmetic composition of claim 1, further comprising an additive selected to increase viscosity without affecting transition temperature of the reversible viscosifying polymer network.

 32. The cosmetic composition of claim 1, further comprising an additive selected to decrease viscosity without affecting transition temperature of the reversible viscosifying polymer network.

25 33. The cosmetic composition of claim 1 or 2, characterized in that the gel remains translucent to light before and after response to the environmental stimulus.

34. The cosmetic composition of claim 1, wherein the poly(acrylic acid) is branched.

35. Method of making an cosmetic composition, comprising:

5 dissolving a poloxamer capable of aggregation in response to a change in temperature in acrylic acid monomer;

initiating polymerization of the monomer to form a poly(acrylic acid) randomly bonded to the poloxamer, so as to form a reversibly viscosifying polymer composition;

10 mixing the reversibly gelling polymer compositions with a cosmetic agent which imparts a desired cosmetic effect to the composition.

36. The method of claim 36, wherein a polymerization initiator is selected to provide the polymer network having a selected temperature of viscosification.

15 37. The method of claim 36, wherein one or more poloxamers are added.

38. The cosmetic composition of claim 1, wherein the reversibly viscosifying polymer network is present in an amount in the range of 0.01% 10%.




Figure 1.




Figure 2




Figure 3




Figure 4

5/28




Figure 5




Figure 6

7/28




Figure 7

8/28




Figure 8

9/28






Figure 9



11/28




Figure 11

12/28




Figure 12




Figure 13

14/28




Figure 14

15/28




Figure 15

16/28




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23






Figure 24

25/28



a



b

Figure 25




Figure 26

27/28




Figure 27




Figure 28

INTERNATIONAL SEARCH REPORT

International application No
PCT/US98/08931

A. CLASSIFICATION OF SUBJECT MATTER

IPC(G) :A61K 7/00, 7/021, 7/025, 7/06, 7/09, 7/16, 7/32, 7/42, 31/74

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/49, 59, 63, 64, 65, 70.1, 70.2, 70.7, 78.02, 70.08, 400, 401, 405

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS: COSMETIC, POLYACRYLIC ACID, POLYMER NETWORK, POLOXAMER

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US 5,662,892 A (BOLICH, JR. et al.) 02 September 1997, see entire document.	1-38
Y	US 5,106,609 A (BOLICH, JR et al.) 21 April 1992, see entire document.	1-38

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	* T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* I* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* Z* document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means	
* P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 AUGUST 1998

Date of mailing of the international search report

02 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230

Authorized officer

SHELLEY A. DODSON

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08931

A. CLASSIFICATION OF SUBJECT MATTER:

US CL : 424/49, 59, 63, 64, 65, 70.1, 70.2, 70.7, 78.02, 70.08, 400, 401, 405

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.